



ANTIOXIDANT AND ANTI-INFLAMMATORY ACTIVITIES OF PLANTS EXTRACTS OF ISRAEL AND PALESTINE. UNEXPLORED PARADISE

Abdullatif Azab^{[a]*}

Keywords: Antioxidant, anti-inflammatory, plant extracts, plant families, current research, future opportunities, systematic mapping.

Antioxidant and anti-inflammatory activities are among the most important properties of plant materials used by humans. For many medicinal plants and other natural products sources, there is clear relationship between these properties. Despite the fact that some approved, commercial drugs were developed from natural products that possess these properties, published literature scan reveals a disappointing image, in some geographical regions, with rich flora, the vast majority of these plants were never studied for antioxidant and/or anti-inflammatory activities. Expectedly, some plant families were extensively studied, while others, with some of the most common and widespread plant species, were almost totally ignored. In this review, we will introduce the current situation of studying medicinal properties of plants, especially antioxidant and anti-inflammatory activities, on the central part of the Eastern region of the Mediterranean basin. We will also present an overall view of future research opportunities and scientific collaborations. These opportunities and collaborations must be based on systematic mapping of current knowledge.

* Corresponding Authors

Fax: +972-4-6205906, +972-50-565-0025

E-Mail: eastern.plants@gmail.com

[a] Eastern Plants Company, Box 868, Arara, Israel 30026

INTRODUCTION

Antioxidant activity of various materials is one of most essential properties for prevention, inhibition and curing of many health disorders, as well as boosting healthy well being.¹ On the same level of importance or even higher, lies the anti-inflammatory activity of materials, and the search for anti-inflammatory agents is as old as humanity itself.² Many studies have indicated the interrelation between these two properties, especially for plant materials, such as extracts, essential oils, pure compounds, chemical modifications and approved drugs based on plant materials. This correlation is very clear when antioxidant activity results from the presence of polyphenolic compounds.³ Expectedly, this clear relationship can be observed when testing various plant and fungi extracts, water and alcoholic extracts found most active.⁴ This result can be understood on the basis of the high polarity of phenolic compounds that are soluble in polar solvents such as water and alcohols, more than in less polar solvents such as ethyl acetate, ether and hexane. But this correlation can be seen in wider view since it exists not only in plant matter, but also in other products derived from other living organisms, such as bee honey.⁵ In this case also, it was evident that the polar (methanolic) extract and its fractions, contained the highest amount of active polyphenols.

The region on the Eastern side of the Mediterranean shores, namely Israel and Palestine, is one of the richest plant habitats. Plant diversity in this area, despite and because of the fact that wide parts of it are various types of arid lands, is unique and vast. This is because except for rain forests and Tundra climates, all other climates and plant habitats are found. This region is the junction of three

continents: Asia, Africa and Europe. To give readers a sense of this richness, we can consider the area of this region (Israel and Palestine) 28292 km². The flora of this area consists 137 plant families that include 2652 identified, different plant species.⁶ Compared to that, the area of Europe is 10180000 km² and it is home for 24700 different plant species.⁷ This means that Europe is approximately 360 times Israel and Palestine are, and if the area index is applied, Europe should have 954720 plant species, while actually it is habitat for only less than 2.6 % of that theoretical number.

Literature and plant selection

Scanning the literature for published studies of antioxidant or anti-inflammatory activities of all 2652 plants of our region, was not a practical consideration, since this number is too large. So, we decided to search for publications about species that are included in plant families that consist of 10 or more plant species. At first look and thought, this might seem too few and arbitrary as plant families that consist of 10 or more species 44 out of 137, meaning around 32 %. But these families include 2355 species out of 2652, meaning around 89 %. Plant families and number of species they include are shown in Table 1.

Results of literature search

After the selection of plant families that we wanted to search for published studies of their antioxidant or anti-inflammatory activities, the selection of the plant species was a very interesting task. For some families, published studies emerged immediately, for many species and sometimes, many publications for each plant (see section 4, discussion). Some families were extensively studied. But these are in small in number, while the majority of families have been very partially studied and in some cases, not even that. Our findings are presented in Table 2.

DISCUSSION

Plants materials and their products, especially plant extracts are in use by humans since the dawn of humanity. Preparing drinks like tea, coffee or cacao, is actually one of the simplest and earliest methods of preparing plant extracts. In addition to nutritional uses, plants and their extracts were among the very first human remedies. Extractions were

prepared using almost all available liquids: water, wine (hydroalcoholic extraction), different plant oils and vinegar (plant made acetic acid solution). Research efforts and financial support of drug discovery from plants, are influenced by many factors, and consequently, they vary over time. Many review articles were published about this topic. One of the most comprehensive was published by Pan *et al*⁷⁵ and other by Shen.⁷⁶

Table 1. Plant families and species that grow in Israel and Palestine (Ref. 6)

Family	Species	Family	Species	Family	Species
Acanthaceae	2	Chenopodiaceae	73	Hypericaceae	9
Adoxaceae	3	Cistaceae	15	Iridaceae	34
Aizoaceae	11	Cleomaceae	4	Ixioliriaceae	1
Alismataceae	5	Colchicaceae	12	Juncaceae	9
Amaranthaceae	15	Convolvulaceae	35	Lamiaceae	115
Amaryllidaceae	51	Crassulaceae	13	Lauraceae	1
Anacardiaceae	8	Cucurbitaceae	7	Lentibulariaceae	2
Apiaceae	97	Cupressaceae	6	Liliaceae	17
Apocynaceae	17	Cymodoceaceae	2	Linaceae	8
Araceae	18	Cynomoriaceae	1	Loranthaceae	1
Araliaceae	1	Cyperaceae	40	Lythraceae	9
Arecaceae	3	Cytinaceae	1	Malvaceae	32
Aristolochiaceae	6	Dennstaedtiaceae	1	Marchantiophyta	20
Asparagaceae	48	Dioscoreaceae	2	Marsileaceae	1
Aspleniaceae	6	Dipsacaceae	15	Meliaceae	1
Asteraceae	279	Dryopteridaceae	1	Menispermaceae	1
Berberidaceae	3	Elaeagnaceae	1	Molluginaceae	1
Biebersteiniaceae	1	Elatinaceae	3	Moraceae	8
Boraginaceae	70	Ephedraceae	3	Moringaceae	2
Brassicaceae	127	Equisetaceae	2	Myrtaceae	3
Bryophyta	67	Ericaceae	1	Neuradaceae	1
Butomaceae	1	Euphorbiaceae	45	Nitrariaceae	3
Cactaceae	1	Fabaceae	274	Nyctaginaceae	4
Campanulaceae	19	Fagaceae	6	Nymphaeaceae	3
Cannabaceae	1	Frankeniaceae	2	Oleaceae	4
Capparaceae	8	Fumariaceae	1	Onagraceae	8
Caprifoliaceae	2	Gentianaceae	5	Ophioglossaceae	2
Caryophyllaceae	117	Geraniaceae	27	Orchidaceae	31
Casuarinaceae	1	Haloragaceae	1	Orobanchaceae	16
Ceratophyllaceae	2	Hydrocharitaceae	4	Oxalidaceae	2
Paeniaceae	1	Pteridaceae	5	Smilacaceae	1
Papaveraceae	36	Ranunculaceae	40	Solanaceae	24
Passifloraceae	1	Resedaceae	15	Styracaceae	1
Phytolaccaceae	1	Rhamnaceae	9	Tamaricaceae	10
Pinaceae	4	Rosaceae	27	Thelypteridiaceae	1
Plantaginaceae	63	Rubiaceae	45	Thymelaeaceae	2
Platanaceae	1	Ruppiaceae	1	Typhaceae	5
Plumbaginaceae	10	Rutaceae	5	Ulmaceae	1
Poaceae	235	Salicaceae	4	Urticaceae	8
Polygalaceae	2	Salvadoraceae	1	Valerianaceae	14
Polygonaceae	39	Salviniaceae	2	Verbenaceae	5
Polypodiaceae	1	Santalaceae	4	Violaceae	5
Pontederiaceae	1	Sapindaceae	4	Vitaceae	2
Portulacaceae	1	Saxifragaceae	2	Xanthorrhoeaceae	7
Potamogetonaceae	8	Scrophulariaceae	30	Zygophyllaceae	17
Primulaceae	8	Simaroubaceae	1		

Table 2. Selected published studies of antioxidant (AO) and anti-inflammatory (AI) of plants species extracts, from major plant families.

Plant family	Plant species	Extract ^a	Activity	Reference
Aizoaceae	<i>Aizoon hispanicum</i>	aq,met	AO	8
	<i>Trianthema portulacastrum</i>	but	AI	9
Amaranthaceae	<i>Amaranthus graecizans</i> ^b	met	AO	10
	<i>Amaranthus graecizans</i> ^b	met	AI	11
Amaryllidaceae	<i>Allium ampeloprasum</i>	met	AO	12
	<i>Allium ampeloprasum</i>	met	AI	13
Apiaceae	<i>Ammi majus</i>	met	AO	14
	<i>Ammi majus</i>	hex,met	AI	15
Apocynaceae	<i>Calotropis procera</i>	aq	AO	16
	<i>Calotropis procera</i>	et,met	AI	17
Araceae	<i>Tanacetum vulgare</i>	EO	AO, AI	18
Asparagaceae	<i>Scilla autumnalis</i>	et	AO	19
	<i>Agave americana</i>	ac	AI	20
Asteraceae	<i>Chrysanthemum coronarium</i>	het	AO	21
	<i>Chrysanthemum coronarium</i>	met	AI	22
Boraginaceae	<i>Anchusa-undulata</i>	meaq	AO	23
	<i>Anchusa-azurea</i>	aq,met	AI	24
Brassicaceae	<i>Sinapis nigra</i>	het	AO	25
	<i>Sinapis alba</i>	het	AI	26
Bryophyta	None (see discussion)	None	None	None
Campanulaceae	<i>Campanula-retrorsa</i>	aq,dcm,met	AO,AI	27
Caryophyllaceae	<i>Silene aegyptiaca</i>	aq,met	AO	28
	<i>Silene vulgaris</i>	et	AI	29
Cistaceae	<i>Cistus salviiifolius</i>	q,meaq	AO	30
	<i>Cistus salviiifolius</i>	aq	AI	31
Colchicaceae	None (see discussion)	None	None	None
Convolvulaceae	<i>Convolvulus arvensis</i>	et	AO	32
	<i>Convolvulus arvensis</i>	et	AI	33
Crassulaceae	<i>Sedum sediforme</i>	aq,met,pet,ac	AO	34
	<i>Sedum sediforme</i>	met	AI	35
Dipsacaceae	<i>Knautia bidens</i>	aq,met	AO	36
	None (see discussion)	None	AI	None
Euphorbiaceae	<i>Euphorbia hirta</i>	met	AO	37
	<i>Euphorbia hirta</i>	et	AI	38
Fabaceae	<i>Ceratonía siliqua</i>	aq (honey)	AO	39
	<i>Ceratonía siliqua</i>	aq	AI	40
Geraniaceae	<i>Erodium laciniatum</i>	hex,het	AO	41
	<i>Geranium robertianum</i>	aq,hex	AO,AI	42
Iridaceae	None (see discussion)	None	None	None
Lamiaceae	<i>Salvia fruticosa</i>	ch,eta,met,but	AI,AO	43
	<i>Salvia officinalis</i>	aq,but	AI	44
Liliaceae	<i>Tulipa systola</i>	pet,et	AO	45
	None (see discussion)	none	AI	none
Malvaceae	<i>Alcea setosa</i>	dcm,met,aq	AO	46
	<i>Malva sylvestris</i>	et	AI	47
Marchantiophyta	None (see discussion)	None	None	None
Orchidaceae	None (see discussion)	None	None	None
Orobanchaceae	<i>Cistanche tubulosa</i>	aq	AO	48
	<i>Cistanche tubulosa</i>	aq	AI	49
Papaveraceae	<i>Papaver somniferum</i>	het	AO	50
	<i>Fumaria capreolata</i>	et	AI	51
Plantaginaceae	<i>Plantago coronopus</i>	hex	AO	52
	<i>Veronica persica</i>	het	AI	53
Poaceae	<i>Hordeum-vulgare</i>	et	AO	54
	<i>Sorghum bicolor</i>	het	AI	55
Polygonaceae	<i>Rumex crispus</i>	aq	AO	56
	<i>Rumex crispus</i>	aq	AI	57
Ranunculaceae	<i>Ranunculus arvensis</i>	aq, met,ac,ch	AO	58
	<i>Ranunculus constantinopolitanus</i>	met	AI	59

Resedaceae	<i>Reseda luteola</i>	chex, het, het, dcm	AO	60
	<i>Reseda luteola</i>	aq	AI	61
Rosaceae	<i>Crataegus aronia</i>	aq	AO	62
	<i>Crataegus monogyna</i>	het	AI	63
Rubiaceae	<i>Galium aparine</i>	met	AO	64
	None (see discussion)	None	AI	None
Scrophulariaceae	<i>Scrophularia hypericifolia</i>	meaq	AO	65
	<i>Scrophularia hypericifolia</i>	het	AI	66
Solanaceae	<i>Datura stramonium</i>	met	AO	67
	<i>Solanum nigrum</i>	Hex,meaq	AI	68
Tamaricaceae	<i>Tamarix aphylla</i>	het	AO	69
	<i>Tamarix aphylla</i>	et	AI	70
Valerianaceae	<i>Centranthus longiflorus</i>	met	AO	71
	<i>Centranthus longiflorus</i>	et	AI	72
Zygophyllaceae	<i>Zygophyllum album</i>	aq	AO	73
	<i>Zygophyllum-simplex</i>	aq,ch	AI,AO	74
Ranunculaceae	<i>Ranunculus arvensis</i>	aq,met,ac,ch	AO	58
	<i>Ranunculus constantinopolitanus</i>	met	AI	59
Resedaceae	<i>Reseda luteola</i>	chex,het,het,dc m	AO	60
	<i>Reseda luteola</i>	aq	AI	61
Rosaceae	<i>Crataegus aronia</i>	aq	AO	62
	<i>Crataegus monogyna</i>	het	AI	63
Rubiaceae	<i>Galium aparine</i>	met	AO	64
	None (see discussion)	None	AI	None
Scrophulariaceae	<i>Scrophularia hypericifolia</i>	meaq	AO	65
	<i>Scrophularia hypericifolia</i>	het	AI	66
Solanaceae	<i>Datura stramonium</i>	met	AO	67
	<i>Solanum nigrum</i>	hex,meaq	AI	68
Tamaricaceae	<i>Tamarix aphylla</i>	het	AO	69
	<i>Tamarix aphylla</i>	et	AI	70
Valerianaceae	<i>Centranthus longiflorus</i>	met	AO	71
	<i>Centranthus longiflorus</i>	et	AI	72
Zygophyllaceae	<i>Zygophyllum album</i>	aq	AO	73
	<i>Zygophyllum simplex</i>	aq,ch	AI,AO	74

a) aq, water; et, ethanol; eta, ethyl acetate; met, methanol; hex, hexane; dcm, dichloromethane; ac, acetone; ch, chloroform; but, *n*-butanol; het, hydroethanol; pet, petroleum ether; EO, essential oil; meaq, methanol-water; chex, cyclohexane. b) There is a mistake in the plant name it should be *sylvestris* not *silvestris*.

The second states that after scientists won the medicine Nobel Prize, for the development of approved drugs based of modifications of natural products, there is a "golden age" for drug discovery.⁷⁶ Both the articles agree that the basic, initial steps are as shown in Figure 1.




Figure 1. First steps of drug discovery from plant sources.

Clinical trials can start with the first step of "plant material preparation", but in most cases they will start after initial extraction, which in many reports, followed by additional fractionation of extracts.

This was presented in many of the publications we have cited. The published activity was linked to certain natural products. For example, Rodrigues Adao *et al.* reported the isolation the steroidal saponin presented in Figure 2. The biological/medicinal importance, especially their anti-

inflammatory activity, of these compounds have been presented.¹³




Figure 2. Steroidal saponin isolated from *Allium ampeloprasum*.¹³




Figure 3. New anti-inflammatory coumarins isolated from *Ammi majus* extracts.¹⁵

The reported anti-inflammatory activity of *Ammi majus* that was reported by Selim and Ouf,¹⁵ is actually not of the plant extract, which was prepared by extraction with *n*-hexane followed by methanol. Two new coumarins were isolated from the extract and were found as active anti-inflammatory agents (Figure 3).

We found during preparing this review article that the vast majority of local plants of Israel and Palestine have never been studied for any biological activity, medicinal property or even for partial chemical composition. The number of examples is in thousands and some of them will be presented here. The presentation will be according to discussed plant family, not consecutively, but according to their appearance in Table 2.

Asparagaceae family includes 48 plants. Very few of them have been studied for antioxidant activity, and only the anti-inflammatory activity of *Agave americana* has been published.²⁰ The major genera of this family (*Bellevia*, 12 species *Ornithogalum*, 11 species) were never studied for either activity.

Yan-Fang *et al.*,²⁶ suggest that the anti-inflammatory activity of *Sinapis alba* results from the presence of sinapine, sinalbin (Figure 4) and the enzyme myrosinase, that act synergistically.




Figure 4. Structures of sinapine and sinalbin found in *Sinapis alba*²⁶

This joint effect of natural products is well known and has gained greater interest over time.⁷⁷ But in recent years, there is a growing recognition of the fact that synergism between plant extracts plays an important role in many biological activities.^{78,79}

The Bryophyta family of non-vascular plants include 67 species in the area of interest of this review article. None of them has been studied for either anti-inflammatory or antioxidant activity. Moreover, most of them have not been studied for any biological activity. *Bryum argenteum* (*n*-butanol extract, among four extracts) has considerable antibacterial activity.⁸⁰ Some non-local species have been studied for their antioxidant⁸¹ and anti-inflammatory activities.⁸²

The case of Colchicaceae family is one of the strangest and most interesting. Israel and Palestine is home to 12 very

beautiful plant species of this family, but none of them was studied for anti-inflammatory or antioxidant activity. Actually, some of them have been studied but most of the interest of researchers was focused on alkaloid content, identification and some biological properties of these alkaloids. Summary of these published studies is shown in Table 3.

Table 3. Summary of published medicinal and phytochemical research of plants of Colchicaceae family, native of Israel and Palestine.

Plant Species	Research Interest and Findings	Ref.
<i>Androcymbium palaestinum</i>	Alkaloids. Two new	83
<i>Colchicum ritchii</i>	Alkaloids: demecolcine, colchicine	84
<i>C. tunicatum</i>	Chemical composition,	85
<i>C. hierosolymitanum</i>	cytotoxicity	
<i>C. tauri</i>	Application of liquid chromatography methods.	86
<i>C. stevenii</i>	18	
<i>C. tunicatum</i>	Alkaloids were identified and their structures are presented	
<i>C. hierosolymitanum</i>	Increasing production of colchicine	87
<i>C. hierosolymitanum</i>	Effect of NPK fertilizer of the production of colchicine	88
<i>C. tunicatum</i>	Effect of NPK fertilizer of the production of colchicine	89
<i>C. hierosolymitanum</i>	Determination of colchicine using various analytical methods	90
<i>C. stevenii</i>		
<i>C. ritchii</i>	Five alkaloids. Two new.	91
<i>C. stevenii</i>	Cytotoxicity study of active compounds isolated using various methods	92
<i>C. tauri</i>	Isolation of nine alkaloids. One new.	93
<i>C. svovitsii</i>	Alkaloids. Structure of szovitsamine	94
<i>C. svovitsii</i>	Alkaloids. <i>O</i> -methylkreysigine	95
<i>C. svovitsii</i>	Alkaloids. Two phenethylisoquinolines	96
<i>C. svovitsii</i>	Analgesic activity of methanolic extract	97

The alkaloid that attracted most research interest is colchicine and most of the isolated alkaloids from *Colchicum* species are its derivatives. Interestingly enough, this compound was tested for anti-inflammatory activity and found active.^{98,99}




Figure 5. Structure of colchicine found in plants of Colchicaceae family.⁸⁴

Finally, it is important to mention that some non-local Colchicaceae species have been studied for anti-inflammatory and antioxidant activities.^{100,101}

Even though Dipsacaceae family includes only 15 local plant species, only *Knautia bidens* was tested for antioxidant activity,³⁶ before we started our experimental research. None of these plants was studied for anti-inflammatory activity despite two very important facts. First, some of the plants of this family (e.g. *Cephalaria joppensis*) are very common in the non-arid areas of Israel and Palestine. Secondly, some non-local species, we studied showed high activity, and these studies were conducted a decade ago.¹⁰²

Local plants of Iridaceae family belong to two major genera, *Crocus* and *Iris*. All plants of this family have very beautiful flowers, and some species are relatively very common. The genus *Iris* have some cultural and national aspects (*I. regis-uzziae* and *I. palaestina*). Despite all this, none of these plants, 34 total of the Iridaceae family, was investigated for antioxidant or anti-inflammatory activities, while some of the non-local species have been studied for both properties.^{103,104}

Three small genera consist the family of Liliaceae, which include a total number of 17 plants. As we have shown in Table 2, only *Tulipa systola* has been studied for antioxidant activity, while none of these species has been studied for anti-inflammatory activity. Non-local plants such as *Fritillaria cirrhosa* have been recently investigated for anti-inflammatory activity.¹⁰⁵

Local liverworts of the Marchantiophyta family belong mainly to the *Riccia* genus, 10 out of 20 species of the whole family. But none of the plants of this genus or the other Marchantiophyta family plants has been investigated for antioxidant or anti-inflammatory activities. As for non-local plants, *Riccia fluitans* and some others, have been extensively studied.^{106,107} Contrary to the ordinary look of Marchantiophyta, the flowers of the Orchidaceae family are among the most spectacular in nature. Yet, the local plants of this family (31) have been totally ignored in terms of antioxidant and anti-inflammatory research, so far. This is not the situation for non-local species.^{108,109}

The genus *Rumex* (Polygonaceae) includes 15 local plants. One of them, *Rumex pulcher* is one of the most important winter delicacies in the Palestinian society. In addition to its nutritional value, it has many uses in traditional medicine. In Lebanon and Syria, the most edible species is *R. acetosa*, and local communities use it as a medicinal plant also.¹¹⁰ New studies of non-local species reveal the health promoting phytochemicals that they contain.¹¹¹ The structures of these compounds are shown in Figure 6.

Strangely enough, the most common and edible local species (*R. pulcher*) has never been investigated for any medicinal, nutritional, phytochemical or any other related property.

Galium (not Gallium) is the genus that includes the largest number of local plants of the Rubiaceae family. As we presented in Table 2, none of the plants of this family was studied for anti-inflammatory activity. *Galium aparine* is one of the most studied plants of this genus, and some

publications claim that its anti-inflammatory activity has been published,¹¹² but we found no reliable support of these claims. Interestingly, asperulosidic acid (Figure 7), was isolated from this plant,¹¹³ and this phytochemical is reported to possess anti-inflammatory activity.¹¹⁴




Figure 6. Phytochemicals isolated from *Rumex cypricus*.¹¹¹




Figure 7. Asperulosidic acid from *Galium aparine* and has anti-inflammatory activity.¹¹³

But it is important to mention that while local species such as *Oldenlandia capensis* were not studied for anti-inflammatory activity, non-local species like *Oldenlandia diffusa* has been extensively studied for many medicinal properties.¹¹⁵

From traditional medicine to methodical science

The findings of this work are very important for us, despite the fact that the method of selecting plant families for our search can be considered inaccurate. We searched for published reports about antioxidant and anti-inflammatory properties of species included in local plant families that have 10 or more plant species. We explained the statistical relevance of our method in the section of Literature and Plant Selection. However, we know that this method suffers two major weaknesses

(1) It produces the impression of equality between families having published works about antioxidant and anti-inflammatory effects. Readers who are not familiar with

plants of the selected area, can not distinguish between the scarcity of studies of Scrophulariaceae on one hand, and Lamiaceae on the other hand, since both appeared in Table 2 as reported.

While reports that we cited about Scrophulariaceae (30 species) are the only ones we found, the Lamiaceae family (115 species) has been very extensively investigated. The difference is not a matter of plant species number only. It is mainly because the Lamiaceae family includes some of the most aromatic and/or medicinal and/or edible, local plant genera namely *Lamium*, *Lavandula*, *Majorana*, *Marrubium*, *Micromeria*, *Nepeta*, *Origanum*, *Phlomis*, *Salvia*, *Stachys*, *Teucrium* and *Ziziphra*. For this reason, there are dozens of reports about the properties and activities of this family.

The Solanaceae family is even a better example. It consisted of 24 plant species, less than the Scrophulariaceae family, but the medicinal activities of the Solanaceae family have been published in numerous studies. In this case, the main reason is that plants of the family are among those that contain the highest concentrations and types of alkaloids, some are very toxic. These plants were used since antiquity as powerful medicinal plants for very wide variety of purposes, including few spiritual applications.

(2) This selection method, on the contrary, ignores some very important medicinal plant families and plant species. Some families that include fewer than 10 plant species are among the most important and most studied. One of these is Oleaceae, which includes *Olea europaea*, common olive tree. Not only olive oil with its superb nutritional and medicinal properties, but all other parts of the tree, including wild species, are vastly researched for antioxidant and anti-inflammatory activities.^{116,117} In the first work, leaves were extracted with aqueous methanol after removing lipophilic fraction with *n*-hexane, and in the second work, anti-inflammatory activity was tested for methanolic and chloroform extracts.

In some cases, the omission is even more apparent. The Portulacaceae family, is locally presented by just one species, *Portulaca oleracea*. This plant has been extensively studied for almost all biological, medicinal and nutritional (widely edible) properties, including antioxidant (aqueous extract),¹¹⁸ and anti-inflammatory (hydroalcoholic extract).¹¹⁹

As we mentioned above, our results of literature search are important. This was revealed to us while preparing and writing our previous works. To link the knowledge of traditional medicine and herbalism of local communities, with modern systematic research, we initiated in 2016 a series of review articles. Each one of these articles reviewed a plant genus of the most known and useful in local and regional traditional medicine and herbalism. In each one of these articles we highlighted the ethnobotanical knowledge and uses of the plants, along with presenting the latest discoveries of medicinal and biological properties. These review articles are shown in Table 4.

But the impression of literature abundance about medicinal plants that these review articles might produce is misleading. While preparing a review article titled, "Anti-

inflammatory Activity of Natural Products",¹²⁵ we discovered the first fault lines of this assumption. Strangely enough, we found that some of the most commonly used plants by local populations, have never been studied for many biological activities. For example, *Eminium spiculatum* (Araceae) is closely related to *Arum* plants,¹²² and like *Arum palaestinum*, it is eaten by local populations and used for cancer treatment in ethnomedicine. We discovered that it was studied for some medicinal activities, such as antioxidant,¹²⁶ but to best of our knowledge, not researched for anti-inflammatory activity up to the time of writing this review.

Table 4. Our published plant-genus-specific review articles.

Genus	Major Species of Interest	Reference
<i>Micromeria</i>	<i>M. fruticosa</i>	120
<i>Alcea</i>	<i>A. setosa</i>	121
<i>Arum</i>	<i>A. palaestinum</i>	122
<i>Malva</i>	<i>M. sylvestris</i>	123
<i>Ceratonia</i>	<i>C. siliqua</i>	124

The Malvaceae family include two of the plant genera that presented in table 4, *Alcea* and *Malva*. Many of the plants of this family are edible and almost all of them are used as medicinal plants. But while some species were very widely studied (*Alcea setosa* and *Malva sylvestris*, see table 2) and such as *Corchorus olitorius* (Molokhia),^{127,128} other have been almost completely ignored. For example, *Malva sylvestris* is most common among the *Malva* species, and it is the largest (by size), it is also the major component of some traditional foods. *Malva nicaeensis* is slightly less common and it grows in hilly landscapes rather than plains like *M. sylvestris*. But *M. nicaeensis* has softer texture and it is considered more delicious. Despite this, while dozens of articles for almost every possible biological or medicinal property of *M. sylvestris* have been published, *M. nicaeensis* was never studied for anti-inflammatory or antioxidant activity. In fact, there are only few reports about heavy metal accumulation in it¹²⁹ and some reports of lipase inhibition.

The great diversity and quantities of local plants sometime results in the neglect of some remarkable species even in traditional medicine practice. One of such ignored plants in local ethnomedicine is *Lotus angustissimus*, a member of the Fabaceae family (274 local species). It is very common in most habitats of Israel and Palestine, with beautiful yellow flowers and a very distinctive smell. But the main reason that makes it potentially interesting for both traditional and modern medicinal research, is the fact that grazing livestock avoid it completely. This is an indication that it might contain toxic natural products, similar to most plant species of the Solanaceae family that have been extensively studied. Grazing animals naturally identify toxic plants and avoid them, and most poisoning cases occur when these plants are dried and mixed with other foods.^{130,131}

The case of *Notobasis syriaca* is even stranger. Locally, it is the only plant of the genus *Notobasis*, but this genus is one of the vast Asteraceae family (279 local plants). Except in desert parts of the reported area, *N. syriaca* is fairly widespread, high with pretty, unmistakable flowers, edible

(peeled, young, fresh stems) and its seeds are used as substitute of coffee beans. In traditional medicine, it is known for its anti-inflammatory use. While preparing the article on anti-inflammatory Activity of natural products^{1,125} we found that this property has never been investigated. As a result, we conducted a research that expectedly yielded positive results.¹³²

At this stage we prepared a list of locally very widespread plants that we planned to study, because each one of them has either never been studied before for various medicinal activities or the published studies are very incomplete or inconsistent. The major two properties that we investigated as a start are total phenolic content and antioxidant activity, but we also studied other activities of these plants extracts. A summary of these studies is presented in Table 5.

Table 5. Summary selected properties we studied of local medicinal plants.

Plant Species	Extracts ^a	Studied Properties	Reference
<i>Notobasis syriaca</i>	aq	Anti-inflammatory	132
<i>Carthamus tenuis</i>	aq, et, etac	Total phenolic content, antioxidant, antifungal	133
<i>Cephalaria joppensis</i>	aq, et, etac	Total phenolic content, antioxidant, antifungal, alkaloid content	134
<i>Notobasis syriaca</i>	aq, et, etac	Total phenolic content, antioxidant, antifungal, anti-termite	135
<i>Scolymus maculatus</i>	aq, et, etac, hex		
<i>Prosopis fatcta</i>	aq, et, etac, hex		

^aaq, aqueous; et, ethanol; etac, ethyl acetate; hex, *n*-hexane

It is important to indicate that anti-inflammatory tests of some extracts mentioned in Table 5, are currently being conducted and others are planned to be performed in the future. It is also important to indicate that our list highly exceeds this small number of partially studied plants.

CONCLUSIONS AND RECOMMENDATIONS

(1) Significant majority of the plant species of Israel and Palestine have never been studied for any biological or medicinal properties.

(2) Some of the plant families have been completely ignored by researcher of anti-inflammatory or antioxidant activities.

(3) Some of these ignored plants and/or families have known properties in traditional medicines.

(4) Researchers should collaborate to plan comprehensive studies of these plant species.

(5) As a start, there is a need for an immediate mapping and documenting published studies of biological and medicinal properties.

(6) There is a need for database of plants/properties that have not been studied yet.

(7) A very comprehensive effort is needed to study the properties of these plants.

(8) Collaboration between researchers from different disciplines in crucial.

(9) Collaboration between researchers from different countries in vital.

(10) Governments of Europe and the Middle East should support this joint effort, since this can bring up some breakthroughs in drug discovery and development.

REFERENCES

- ¹Kasotel, D.M, Katyare, S.S., Hegde, M.V, Bae, H. Significance of Antioxidant Potential of Plants and its Relevance to Therapeutic Applications, *Int. J. Biol. Sci.* **2015**, *11*, 982-991. doi: 10.7150/ijbs.12096
- ²Dinarello, C.A. Anti-inflammatory Agents: Present and Future, *Cell* **2010**, *140*, 935-950. doi: 10.1016/j.cell.2010.02.043
- ³Zhang, L., Ravipati, A.S., Koyyalamudi, S.R., Jeong, S.C., Reddy, N., Smith, P.T., Barlett, J., Shanmugam, K., Munch, G., Wu, M.J. Antioxidant and Anti-inflammatory Activities of Selected Medicinal Plants Containing Phenolic and Flavonoid Compounds, *J. Agric. Food Chem.* **2011**, *59*, 12361-12367. <http://dx.doi.org/10.1021/jf203146e>
- ⁴Diaz, P., Jeong, S.C., Lee, S., Khoo, C., Koyyalamudi, S.R. Antioxidant and anti-inflammatory activities of selected medicinal plants and fungi containing phenolic and flavonoid compounds, *Chin. Med.* **2012**, *7*:26. doi:10.1186/1749-8546-7-26
- ⁵Ruiz-Ruiz, J.C., Matus-Basto, A.J., Acereto-Escoffie, P., Segura-Campos, M.R. Antioxidant and anti-inflammatory activities of phenolic compounds isolated from *Melipona beecheii* honey, *Food Agric. Immunol.* **2017**, *28*, 1424-1437. <https://doi.org/10.1080/09540105.2017.1347148>
- ⁶Wild Flowers of Israel: <http://www.wildflowers.co.il/english/families.asp>
- ⁷Planta Europa: <https://rm.coe.int/16807462bd>
- ⁸Khettaf, A., Belloula, N., Dridi, S. Antioxidant activity, phenolic and flavonoid contents of some wild medicinal plants in southeastern Algeria, *Afr. J. Biotechnol.* **2016**, *15*, 524-530. doi: 10.5897/AJB2015.14459
- ⁹Yadav, E., Singh, D., Yadav, P., Verma, A. Attenuation of dermal wounds via downregulating oxidative stress and inflammatory markers by protocatechuic acid rich *n*-butanol fraction of *Trianthema portulacastrum* Linn. in wistar albino rats, *Biomed. Pharmacother.* **2017**, *96*, 86-97. <http://dx.doi.org/10.1016/j.biopha.2017.09.125>
- ¹⁰Ishtiaq, S., Ahmad, M., Hanif, U., Akbar, S., Mehjabeen, Kamran, S.H. Phytochemical and *in vitro* antioxidant evaluation of different fractions of *Amaranthus graecizans* subsp. *Silvestris* (Vill.) Brenan, *Asian Pac. J. Trop. Med.* **2014**, *7*, S342-S347. doi: 10.1016/S1995-7645(14)60256-X
- ¹¹Ishtiaq, S., Ali, T., Ahmad, B., Anwar, F., Khan Afridi, M.S., Shaheen, H. Phytochemical and Biological Evaluations of Methanolic Extract of *Amaranthus graecizans* subsp. *silvestris* (Vill.) Brenan, *Br. J. Pharm. Res.* **2017**, *15*, 1-11. doi: 10.9734/BJPR/2017/31698
- ¹²Garcia-Herrera, P., Morales, P., Fernandez-Ruiz, V., Sanchez-Mata, M.C., Camara, M., Carvalho, A.M., Ferreira, I.C.F.R., Pardo-de-Santayana, M., Molina, M., Tardio, J. Nutrients, phytochemicals and antioxidant activity in wild populations of *Allium ampeloprasum* L., a valuable underutilized vegetable, *Food Res. Int.* **2014**, *62*, 272-279. <https://doi.org/10.1016/j.foodres.2014.03.004>

- ¹³Rodrigues Adao, C., Pereira da Silva, B., Paz Parente, J. A new steroidal saponin with anti-inflammatory and antiulcerogenic properties from the bulbs of *Allium ampeloprasum* var. Porrum, *Fitoterapia*. **2011**, *82*, 1175-1180. doi:10.1016/j.fitote.2011.08.003
- ¹⁴Al-Hadhrami, R.M.S., Hossain, M. A. Evaluation of antioxidant, antimicrobial and cytotoxic activities of seed crude extracts of *Ammi majus* grown in Oman, *Egypt. J. Basic Appl. Sci.* **2016**, *3*, 329-334. <http://dx.doi.org/10.1016/j.ejbas.2016.08.001>
- ¹⁵Selim, Y.A., Ouf, N.H. Anti-inflammatory new coumarin from the *Ammi majus* L, *Org. Med. Chem. Lett.* **2012**, *2*, 4 pages. doi:10.1186/2191-2858-2-1
- ¹⁶Kumar, S., Gupta, A., Pandey, A.K. *Calotropis procera* Root Extract Has the Capability to Combat Free Radical Mediated Damage, *ISRN Pharmacol.* **2013**, Article ID 691372, 8 pages. <http://dx.doi.org/10.1155/2013/691372>
- ¹⁷Arya, S., Kumar, V.L. Antiinflammatory Efficacy of Extracts of Latex of *Calotropis procera* Against Different Mediators of Inflammation, *Mediators Inflamm.* **2005**, *4*, 228-232. doi: 10.1155/MI.2005.228
- ¹⁸Cote, H., Boucher, M.A., Pichette, A., Legault, J. Anti-Inflammatory, Antioxidant, Antibiotic, and Cytotoxic Activities of *Tanacetum vulgare* L. Essential Oil and Its Constituents, *Medicines*. **2017**, *4*, 34-42. doi:10.3390/medicines4020034
- ¹⁹Mammadov, R., Kaska, A., Ozay, C. Phenolic Composition, Antioxidant and Cytotoxic Activities of *Prospero autumnale*, *Indian J. Pharm. Sci.* **2017**, *79*, 585-590. doi: 10.4172/pharmaceutical-sciences.1000266
- ²⁰Monterrosas-Brisson, N., Arenas Ocampo, M.L., Jimenez-Ferrer, E., Jimenez-Aparicio, A.R., Zamilpa, A., Gonzalez-Cortazar, M., Tortoriello, J., Herrera-Ruiz, M. Anti-Inflammatory Activity of Different Agave Plants and the Compound *Cantalasaponin-1*, *Molecules*. **2013**, *18*, 8136-8146. doi:10.3390/molecules18078136
- ²¹Donia, A.M. Biological Activity of *Chrysanthemum coronarium* L. Extracts. *Annu. Res. Rev. Biol.* **2014**, *4*, 2617-2627. doi: 10.9734/ARRB/2014/10112
- ²²Amro, B.I., Haddadin, R.N., Tawaha, K., Mohammad, M., Mashallah, S., Assaf, A.M. *In vitro* antimicrobial and anti-inflammatory activity of Jordanian plant extracts: A potential target therapy for *Acne vulgaris*, *Afr. J. Pharm. Pharmacol.* **2013**, *7*, 2087-2099. doi: 10.5897/AJPP2013.3497
- ²³Taban, K., Eruygun, N., Ustun, O. Biological activity studies on the aqueous methanol extract of *Anchusa undulata* L. subsp. *hybrida* (Ten.) Coutinho, *Marmara Pharm. J.* **2018**, *22*, 357-364. <https://doi.org/10.12991/jrp.2018.75>
- ²⁴Kuruuzum-Uz, A., Suleyman, H., Cadirci, E., Guvenalp, Z., Demirezer, L.O. Investigation on Anti-Inflammatory and Antiulcer Activities of *Anchusa azurea* Extracts and their Major Constituent Rosmarinic Acid, *Z. Naturforsch. C.* **2012**, *67*, 360-366. <https://doi.org/10.1515/znc-2012-7-802>
- ²⁵Boscaro, V., Boffa, L., Binello, A., Amisano, G., Fornasero, S., Cravotto, G., Gallicchio, M. Antiproliferative, Proapoptotic, Antioxidant and Antimicrobial Effects of *Sinapis nigra* L. And *Sinapis alba* L. Extracts, *Molecules*. **2018**, *23*, 3004-3020. doi:10.3390/molecules23113004
- ²⁶Yan-Fang, X., Zhen, H., Siu-Po, I., Jian-Nan, C., Zi-Ren, S., Xiao-Pin, L., Zhi-Xiu, L. Comparison of the anti-inflammatory effects of *Sinapis alba* and *Brassica juncea* in mouse models of inflammation, *Phytomedicine*. **2018**, *50*, 196-204. <https://doi.org/10.1016/j.phymed.2018.05.010>
- ²⁷Alhaje, J., Elbitar, H., Taha, S., Benvegnu, T. *In vitro* assessment of antioxidant, antimicrobial, cytotoxic, anti-inflammatory, and antidiabetic activities of *Campanula retrorsa* crude extracts, *Pharmacogn. Res.* **2018**, *10*, 397-403. doi: 10.4103/pr.pr_73_18
- ²⁸Tawaha, K., Alali, F.Q., Gharaibeh, M., Mohammad, M., El-Elimat, T. Antioxidant activity and total phenolic content of selected Jordanian plant species, *Food Chem.* **2007**, *104*, 1372-1378. doi: 10.1016/j.foodchem.2007.01.064
- ²⁹Boukhira, S., El Mansouri, L., Bouarfa, M., Ouhammou, A., Achour, S., Khadhr, M., Bousta, D. Phytochemical Screening, Anti-Inflammatory and Analgesic Activities Of Formulation Cream of *Silene vulgaris*, *Res. J. Med. Plant.* **2016**, *10*, 150-158. doi: 10.3923/rjmp.2016.150.158
- ³⁰Sayah, K., Marmouzi, I., Mrabti, H.N., Cherrah, Y., Faouzi, M.E. Antioxidant Activity and Inhibitory Potential of *Cistus salviifolius* (L.) and *Cistus monspeliensis* (L.) Aerial Parts Extracts against Key Enzymes Linked to Hyperglycemia, *BioMed Res.Int.* **2017** Article ID 2789482, 7 pages. <https://doi.org/10.1155/2017/2789482>
- ³¹Sayah, K., Chemlal, L., Marmouzi, I., El Jemli, Y., Cherrah, Y., Faouzi, M.E. *In vivo* anti-inflammatory and analgesic activities of *Cistus salviifolius* (L.) and *Cistus monspeliensis* (L.) aqueous extracts, *S. Afr. J. Bot.* **2017**, *113*, 160-163. <http://dx.doi.org/10.1016/j.sajb.2017.08.015>
- ³²Azman, N.A. Gallego, M.G., Julia, L., Fajari, L., Almajano, M.P. The Effect of *Convolvulus arvensis* Dried Extract as a Potential Antioxidant in Food Models, *Antioxidants*. **2015**, *4*, 170-184. doi:10.3390/antiox4010170
- ³³Ali, M., Qadir, M.I., Saleem, M., Janbaz, K.H., Gul, H., Hussain, L., Ahmad, B. Hepatoprotective potential of *Convolvulus arvensis* against paracetamol induced hepatotoxicity, *Bangladesh J. Pharmacol.* **2013**, *8*, 300-304. doi: 10.3329/bjp.v8i3.15165
- ³⁴Ertas, A., Boga, M., Yilmaz, M.A., Yesil, Y., Hasimi, N., Kaya, M.S., Temel, H., Kolak, U. Chemical Compositions by Using LC-MS/MS and GC-MS and Biological Activities of *Sedum sediforme* (Jacq.) Pau, *J. Agric. Food Chem.* **2014**, *62*, 4601-4609. doi: 10.1021/jf500067q
- ³⁵Winekenst dde, D., Angelis, A., Waltenberger, B., Schwaiger, S., Tchoumtchoua, J., K nig, S., Werz, O., Aligiannis, N., Skaltsounis, A.L., Stuppner, H. Phytochemical profile of the aerial parts of *Sedum sediforme* and anti-inflammatory activity of myricitrin, *Nat. Prod. Commun.* **2015**, *10*, 83-88. <https://doi.org/10.1177/1934578X1501000122>
- ³⁶Alali, F.Q., Tawaha, K., El-Elimat, T., Syouf, M., El-Fayad, M., Abulaaila, K., Nielsen, S.J., Wheaton, W.D., Falkinham, J.O., Oberlies, N.H. Antioxidant activity and total phenolic content of aqueous and methanolic extracts of Jordanian plants: an ICBG project, *Nat. Prod. Res.* **2007**, *21*, 1121-1131. doi: 10.1080/14786410701590285
- ³⁷Abu Arra, B., Zuraini, Z., Lacimanan, Y.L., Sasidharan, S. Antioxidant activity and phytochemical screening of the methanol extracts of *Euphorbia hirta* L, *Asian Pac. J. Trop. Med.* **2011**, *4*, 386-390. [https://doi.org/10.1016/s1995-7645\(11\)60109-0](https://doi.org/10.1016/s1995-7645(11)60109-0)
- ³⁸Xia, M., Liu, L., Qiu, R., Li, M., Huang, W., Ren, G., Zhang, J. Anti-inflammatory and anxiolytic activities of *Euphorbia hirta* extract in neonatal asthmatic rats, *AMB Express*. **2018**, *8*:179, 11 pages. <https://doi.org/10.1186/s13568-018-0707-z>
- ³⁹El-Haskoury, R., Kriaa, W., Lyoussi, B., Makni, M. *Ceratonia siliqua* honeys from Morocco: Physicochemical properties, mineral contents, and antioxidant activities, *J. Food Drug Anal.* **2018**, *26*, 67-73. <http://dx.doi.org/10.1016/j.jfda.2016.11.016>
- ⁴⁰Rtibi, K., Jabri, M.A., Selmi, S., Sebai, H., Marie, J.C., Amri, M., Marzouki, L., El-Benna, J. Preventive effect of carob (*Ceratonia siliqua* L.) in dextran sulfate sodium-induced ulcerative colitis in rat, *RSC Adv.* **2016**, *6*, 19992-20000. doi: 10.1039/c5ra21388f
- ⁴¹Jaradat, N., Almasri, M., Zaid, A., Othman, D.G. Pharmacological and phytochemical screening of Palestinian traditional medicinal plants *Erodium laciniatum* and *Lactuca orientalis*, *J. Complement. Integr. Med.* **2017**, *15*, 16 pages. doi: 10.1515/jcim-2017-0059

- ⁴²Catarino, M.D., Silva, A.M.S., Cruz, M.T., Cardoso, S.M. Antioxidant and anti-inflammatory activities of *Geranium robertianum* L. decoctions, *Food Funct.* **2017**, *20*, 3355-3365. doi: 10.1039/c7fo00881c
- ⁴³Boukhary, R., Raafat, K., Ghoneim, A.I., Aboul-Ela, M., El-Lakany, A. Anti-Inflammatory and Antioxidant Activities of *Salvia fruticosa*: An HPLC Determination of Phenolic Contents, *Evid. Based Complementary Altern. Med.* **2016**, Article ID 7178105, 6 pages. <http://dx.doi.org/10.1155/2016/7178105>
- ⁴⁴Qnais, E.Y., Abu-Dieyeh, M., Abdulla, F.A., Abdalla, S.S. The antinociceptive and anti-inflammatory effects of *Salvia officinalis* leaf aqueous and butanol extracts, *Pharm. Biol.* **2010**, *48*, 1149-1156. doi: 10.3109/13880200903530763
- ⁴⁵Ibrahim, M.F., Hussain, F.H.S., Zaroni, G., Vidari, G. The main constituents of *Tulipa systola* Stapf. roots and flowers, their antioxidant activities, *Nat. Prod. Res.* **2017**, *31*, 2001-2007. doi: 10.1080/14786419.2016.1272107
- ⁴⁶Alhaje, J., Elbitar, H. In vitro screening for antioxidant and antimicrobial properties of three Lebanese medicinal plants crude extracts, *Pharmacogn. Res.* **2019**, *11*, 127-133. doi: 10.4103/pr.pr_171_18
- ⁴⁷Martins, C.A.F., Campos, M.L., Irioda, A.C., Stremel, D.P., Trindade, A.C.L.B., Pontarolo, R. Anti-Inflammatory Effect of *Malva sylvestris*, *Sida cordifolia*, and *Pelargonium graveolens* Is Related to Inhibition of Prostanoid Production, *Molecules.* **2017**, *22*, 1883-1847. doi: 10.3390/molecules22111883
- ⁴⁸Zhang, W., Huang, J., Wang, W., Li, Q., Chen, Y., Feng, W., Zheng, D., Zhao, T., Mao, G., Yang, L., Wu, X. Extraction, purification, characterization and antioxidant activities of polysaccharides from *Cistanche tubulosa*, *Int. J. Biol. Macromol.* **2016**, *93*, 448-458. doi: 10.1016/j.jbiomac.2016.08.079
- ⁴⁹Kyung, J., Kim, D., Park, D., Yang, Y.H., Choi, E.K., Lee, S.P., Kim, T.S., Lee, Y.B., Kim, Y.B. Synergistic anti-inflammatory effects of *Laminaria japonica* fucoidan and *Cistanche tubulosa* extract, *Lab. Anim. Res.* **2012**, *28*, 91-97. <http://dx.doi.org/10.5625/lar.2012.28.2.91>
- ⁵⁰Krosiak E., Maliar, T., Nemecek, P., Viskupicova, J., Maliarova, M., Havrlentova, M., Kraic, J. Antioxidant and Proteinase Inhibitory Activities of Selected Poppy (*Papaver somniferum* L.) Genotypes, *Chem. Biodivers.* **2017**, *14*, 10 pages. doi: 10.1002/cbdv.201700176
- ⁵¹Bribi, N., Algieri, F., Rodriguez-Nogales, A., Garrido-Mesa, J., Vezza, T., Maiza, F., Utrilla, M.P., Rodriguez-Cabezas, M.E., Galvez, J. Antinociceptive and Anti-Inflammatory Effects of Total Alkaloid Extract from *Fumaria capreolata*, *Evid. Based Complementary Altern. Med.* **2015**, Article ID 736895, 7 pages. <http://dx.doi.org/10.1155/2015/736895>
- ⁵²Pereira, C.G., Custodio, L., Rodrigues, M.J., Neng, N.R., Nogueira, J.M.F., Carlier, J., Costa, M.C., Varela, J., Barreira, L. Profiling of antioxidant potential and phytoconstituents of *Plantago coronopus*, *Braz. J. Biol.* **2017**, *77*, 632-641. <http://dx.doi.org/10.1590/1519-6984.02416>
- ⁵³Fierascu, R.C., Georgiev, M.I., Fierascu, I., Ungureanu, C., Avramescu, S.M., Ortan, A., Georgescu, M.I., Sutan, A.N., Zandirescu, A., Dinu-Pirvu, C.E., Velescu, B.S., Anuta, V. Mitodepressive, antioxidant, antifungal and anti-inflammatory effects of wild-growing Romanian native *Arctium lappa* L. (Asteraceae) and *Veronica persica* Poiret (Plantaginaceae), *Food Chem. Toxicol.* **2018**, *111*, 44-52. <https://doi.org/10.1016/j.fct.2017.11.008>
- ⁵⁴Omwamba, M., Feng Li, F., Sun, G., Hu, Q. Antioxidant Effect of Roasted Barley (*Hordeum vulgare* L.) Grain Extract towards Oxidative Stress *In Vitro* and *In Vivo*, *Food Nutr. Sci.* **2013**, *4*, 139-146. <http://dx.doi.org/10.4236/fns.2013.48A017>
- ⁵⁵Burdette, A., Garner, P.L., Mayer, E.P., Hargrove, J.L., Hartle, D.K., Greenspan, P. Anti-Inflammatory Activity of Select Sorghum (*Sorghum bicolor*) Brans, *J. Med. Food.* **2010**, *13*, 1-9. doi: 10.1089/jmf.2009.0147
- ⁵⁶Yildirim, A., Mavi, A., Kara, A.A. Determination of Antioxidant and Antimicrobial Activities of *Rumex crispus* L. Extracts, *J. Agric. Food Chem.* **2001**, *49*, 4083-4089. doi: 10.1021/jf0103572
- ⁵⁷Park, E.S., Song, G.H., Kim, S.H., Lee, S.M., Kim, Y.G., Lim, Y.L., Kang, S.A., Park, K.Y. *Rumex crispus* and *Cordyceps militaris* Mixture Ameliorates Production of Pro-Inflammatory Cytokines Induced by Lipopolysaccharide in C57BL/6 Mice Splenocytes, *Prev. Nutr. Food Sci.* **2018**, *23*, 374-381. <https://doi.org/10.3746/pnf.2018.23.4.374>
- ⁵⁸Bhatti, A.Z., Ali, A., Ahmad, A., Saeed, A., Malik, S.A. Antioxidant and phytochemical analysis of *Ranunculus arvensis* L. extracts, *BMC Res. Notes.* **2015**, *8*:279, 8 pages. doi: 10.1186/s13104-015-1228-3
- ⁵⁹Akkol, E.K., Sutar, I., Erdogan, T.F., Keles, H., Gonenc, T.M., Kivcak, B. Wound healing and anti-inflammatory properties of *Ranunculus pedatus* and *Ranunculus constantinopolitanus*: A comparative study, *J. Ethnopharmacol.* **2012**, *139*, 478-484. doi: 10.1016/j.jep.2011.11.037
- ⁶⁰Burger, P., Monchot, A., Bagarri, O., Chiffolleau, P., Azoulay, S., Fernandez, X., Michel, T. Whitening Agents from *Reseda luteola* L. and Their Chemical Characterization Using Combination of CPC, UPLC-HRMS and NMR, *Cosmetics.* **2017**, *4*, 51-65. doi:10.3390/cosmetics4040051
- ⁶¹Kim, S.S., Seo, J.Y., Lim, S.S., Suh, H.J., Kim, L., Kim, J.S. Neuroprotective Effect of *Reseda luteola* L. Extract in a Mouse Neuronal Cell Model, *Food Sci. Biotechnol.* **2015**, *24*, 333-339. doi: 10.1007/s10068-015-0044-9
- ⁶²Ljubuncic, P., Portnaya, I., Cogan, U., Azaizeh, H., Bomzon, A. Antioxidant activity of *Crataegus aronia* aqueous extract used in traditional Arab medicine in Israel, *J. Ethnopharmacol.* **2005**, *101*, 153-161. doi:10.1016/j.jep.2005.04.024
- ⁶³Tadic, V.M., Dobric, S., Markovic, G.M., Dordevic, S.M., Arsic, I., Menkovic, N.R., Stevic, T. Anti-inflammatory, Gastroprotective, Free-Radical-Scavenging, and Antimicrobial Activities of Hawthorn Berries Ethanol Extract, *J. Agric. Food Chem.* **2008**, *56*, 7700-7709. doi: 10.1021/jf801668c
- ⁶⁴Bokhari, J., Khan, M.R., Shabbir, M., Rashid, U., Jan, S., Zai, J.A. Evaluation of diverse antioxidant activities of *Galium aparine*, *Spectrochim. Acta A.* **2013**, *102*, 24-29. <http://dx.doi.org/10.1016/j.saa.2012.09.056>
- ⁶⁵Shahat, A.A., Ibrahim, A.Y., Ezzeldin, E., Alsaid, M.S. Acetylcholinesterase Inhibition and Antioxidant Activity of Some Medicinal Plants for Treating Neuro Degenerative Disease, *Afr. J. Tradit Complement. Altern. Med.* **2015**, *12*, 97-103. <http://dx.doi.org/10.4314/ajtcam.v12i3.12>
- ⁶⁶Alqasoumi, S.I., Evaluation of the hepatoprotective and nephroprotective activities of *Scrophularia hypericifolia* growing in Saudi Arabia, *Saudi Pharm. J.* **2014**, *22*, 258-263. <http://dx.doi.org/10.1016/j.jsps.2013.12.001>
- ⁶⁷Sharma, P., Bhardwaj, R., Yadav, A., Sharmam R.A. Study of Antioxidant Activity of *Datura stramonium* Linn, *Res. J. Phytochem.* **2014**, *8*, 112-118. doi: 10.3923/rjphyto.2014.112.118
- ⁶⁸Wang, Y., Xiang, L., Yi, X., He, X. Potential Anti-inflammatory Steroidal Saponins from the Berries of *Solanum nigrum* L. (European Black Nightshade), *J. Agric. Food Chem.* **2017**, *65*, 4262-4272. doi: 10.1021/acs.jafc.7b00985
- ⁶⁹Yusufoglu, H.S., Alqasoumi, S.I. Anti-inflammatory and Wound Healing Activities of Herbal Gel Containing an Antioxidant *Tamarix aphylla* Leaf Extract, *Int. J. Pharmacol.* **2011**, *7*, 829-835. doi: 10.3923/ijp.2011.829.835
- ⁷⁰Ali, S., Alam, A., Ahmad, S., Ali, M., Ahsan, W., Siddiqui, M.R., Ansari, S., Shamim, M., Ali, D. Wound Healing Activity of Alcoholic Extract of *Tamarix aphylla* L. on Animal Models, *Biomed. Pharmacol. J.* **2019**, *12*, 41-48. <http://dx.doi.org/10.13005/bpi/1611>

- ⁷¹Aliyazicioglu, R., Korkmaz, N., Akkaya, S., Sener, S.O. Phenolic components, antioxidant and antimicrobial activities of *Centranthus longiflorus* L., *Int. J. Adv. Res. Biol. Sci.* **2016**, *3*, 80-87. doi: 10.22192/ijarbs
- ⁷²Askin, H., Yilmaz, B., Bakirci, S., Ayar, A. Simultaneous determination of α -amyrin and β -sitosterol in *Centranthus longiflorus* Stev. Subsp. *longiflorus* Stev and *Iris taochia* Woronow ex Grossh by GC-MS method, *Prog. Nutr.* **2018**, *20*, 209-217. doi: 10.23751/pn.v20i1-S.6744
- ⁷³El Ghoul, J., Smiri, M., Ghrab, S., Boughattas, N.A., Ben-Attia, M. Antihyperglycemic, antihyperlipidemic and antioxidant activities of traditional aqueous extract of *Zygophyllum album* in streptozotocin diabetic mice, *Pathophysiology.* **2012**, *19*, 35-42. doi: 10.1016/j.pathophys.2011.12.001
- ⁷⁴Abdallah, H.M., Esmat, A. Antioxidant and anti-inflammatory activities of the major phenolics from *Zygophyllum simplex* L., *J. Ethnopharmacol.* **2017**, *205*, 51-56. <http://dx.doi.org/10.1016/j.jep.2017.04.022>
- ⁷⁵Pan, S.Y., Zhou, S.F., Gao, S.H., Yu, Z.L., Zhang, S.F., Tang, M.K., Sun, J.N., Ma, D.L., Han, Y.F., Fong, W.F., Ko, K.M. New Perspectives on How to Discover Drugs from Herbal Medicines: CAM's Outstanding Contribution to Modern Therapeutics, *Evid. Based Complementary Altern. Med.* **2013**, Article ID 627375, 25 pages. <http://dx.doi.org/10.1155/2013/627375>
- ⁷⁶Shen, B. A New Golden Age of Natural Products Drug Discovery, *Cell.* **2015**, *163*, 1297-1304. doi:10.1016/j.cell.2015.11.031.
- ⁷⁷Capitani, C.D., Carvalho, A.C.L., Botelho, P.B., Carrapeiro, M.M., Castro, I.A. Synergism on antioxidant activity between natural compounds optimized by response surface methodology, *Eur. J. Lipid Sci. Tech.* **2009**, *111*, 1100-1110. <https://doi.org/10.1002/ejlt.200800258>
- ⁷⁸Cheung, D.W.S., Koon, C.M., Ng, C.F., Leung, P.C., Fung, K.P., Poon, S.K.S., Lau, C.B.S. The roots of *Salvia miltiorrhiza* (Danshen) and *Pueraria lobata* (Gegen) inhibit atherogenic events: A study of the combination effects of the 2-herb formula, *J. Ethnopharmacol.* **2012**, *143*, 859-866. <http://dx.doi.org/10.1016/j.jep.2012.08.011>
- ⁷⁹Caesar, L.K., Cech, N.B. Synergy and antagonism in natural product extracts: when 1 + 1 does not equal 2, *Nat. Prod. Rep.* **2019**, *36*, 869-888. doi: 10.1039/c9np00011a
- ⁸⁰Singh, M., Singh, S., Nath, V., Sahu, V., Kumar, A.S.R. Antibacterial activity of some bryophytes used traditionally for the treatment of burn infection, *Pharm. Biol.* **2011**, *49*, 526-530. doi: 10.3109/13880209.2010.523007
- ⁸¹Erturk, O., Sahin, H., Erturk, E.Y., Hotaman, H.E., Koz, B., Ozdemir, O. The antimicrobial and antioxidant activities of extracts obtained from some moss species in Turkey, *Herba Pol.* **2015**, *61*, 52-65. <https://doi.org/10.1515/hepo-2015-0031>
- ⁸²Tosun, A., Akkol, E.K., Suntar, I., Kiremit, H.O., Asakawa, Y. Phytochemical investigations and bioactivity evaluation of liverworts as a function of anti-inflammatory and antinociceptive properties in animal models, *Pahrm. Biol.* **2013**, *51*, 1008-1013. doi: 10.3109/13880209.2013.774028
- ⁸³Tojo, E., Abu Zarga, M.H., Sabri, S.S., Freyer, A.J., Shamma, M. The Homoaporphine Alkaloids of *Androcymbium palaestinum*, *J. Nat. Prod.* **1989**, *52*, 1055-1059. <https://doi.org/10.1021/np50065a023>
- ⁸⁴Alali, F.Q., Tawaha, K., El-Elimat, T. Determination of (-)-demecolcine and (-)-colchicine content in selected Jordanian *Colchicum* species, *Pharmazie.* **2007**, *62*, 739-742. doi: 10.1691/ph.2007.10.7032
- ⁸⁵Alali, F., Tawaha, K., El-Elimat, T., Qassaymeh, R., Li, C., Burgess, J.P., Nakanishi, Y., Kroll, D.J., Wani, M.C., Oberlies, N. H. Phytochemical studies and cytotoxicity evaluations of *Colchicum tunicatum* Feinbr and *Colchicum hierosolymitanum* Feinbr (Colchicaceae): Two native Jordanian meadow saffrons, *Nat. Prod. Res.* **2006**, *20*, 558-566. doi: 10.1080/14786410500183381
- ⁸⁶Gharaibeh, A.A., Al-Serini, A., Qasaymeh, R. M., Maayah, A.S., Tawaha, K., El-Elimat, T., Alali, F.Q. Liquid Chromatography-Mass Spectroscopy and Liquid Chromatography Ultraviolet/Visible Photodiode Array Analysis of Selected *Colchicum* Species, *Z. Naturforsch. C.* **2012**, *67*, 451-460. doi: <https://doi.org/10.1515/znc-2012-9-1002>
- ⁸⁷Daradkeh, N.Q., hibli, R.A., Makhadmeh, I.M., Alali, F., Al-Qudah, T.S. Cell Suspension and *In Vitro* Production of Colchicine in Wild *Colchicum Hierosolymitanum* Feib, *Open Conf. Proc. J.* **2012**, *3*, 52-59. doi: 10.2174/1876326X01203020052
- ⁸⁸Al-Fayyad, M., Alali, F., Al-Tell, A. Effect of NPK Fertilizer Levels on Morphological Characteristics and Productivity of *Colchicum hierosolymitanum* and *Colchicum tunicatum*, *J. Herbs Spices Med Plants.* **2004**, *10*, 11-17. doi: 10.1300/J044v10n04_02
- ⁸⁹Al-Fayyad, M., Alali, F., Alkofahi, A., Tell, A. Determination of colchicine content in *Colchicum hierosolymitanum* and *Colchicum tunicatum* under cultivation, *Nat. Prod. Lett.* **2002**, *16*, 395-400. doi: 10.1080/10575630290033178
- ⁹⁰Alali, F., Tawaha, K., Qasaymeh, R.M. Determination of colchicine in *Colchicum stevenii* and *C. hierosolymitanum* (Colchicaceae): comparison between two analytical methods, *Phytochem. Anal.* **2004**, *15*, 27-29. doi: 10.1002/pca.738
- ⁹¹Freyer, A.J., Abu Zarga, M.H., Firdous, S., Guinaudeau, H., Shamma, M. Five New Alkaloids from *Colchicum ritchei*, *J. Nat. Prod.* **1987**, *50*, 684-689. <https://doi.org/10.1021/np50052a018>
- ⁹²Al-Mahmoud, M.S., Alali, F.Q., Tawaha, K., Qasaymeh, R.M. Phytochemical study and cytotoxicity evaluation of *Colchicum stevenii* Kunth (Colchicaceae): a Jordanian meadow saffron, *Nat. Prod. Res.* **2006**, *20*, 153-60. doi: 10.1080/14786410500046224
- ⁹³Alali, F.Q., Maayaa, A.S., Alkofahi, A., Qandil, A., Li, C., Burgess, J., Wani, m.c., Oberlies, N.H. A New Colchicinoid from *Colchicum tauri*, an Unexplored Meadow Saffron Native to Jordan, *Nat. Prod. Commun.* **2006**, *1*, 95-99. <https://doi.org/10.1177/1934578x0600100203>
- ⁹⁴Yusupov, M.K., Ngo, D., Aslanov, K.A., Sadykov, A.S. Alkaloids of *Colchicum szovitsii*. Structure of szovitsamine, *Chem. Nat. Compd.* **1975**, *11*, 127-128. <https://doi.org/10.1007/bf00567063>
- ⁹⁵Yusupov, M.K., Ngo, D., Aslanov, K.A.. O-methylkreysigine from *Colchicum szovitsii*, *Chem. Nat. Compd.* **1975**, *11*, 555-556. <https://doi.org/10.1007/bf00566822>
- ⁹⁶Tojo, E., Onur, M.A., Freyer, A.J., Shamma, M. Two Trioxxygenated Phenethylisoquinoline Alkaloids from *Colchicum szovitsii*, *J. Nat. Prod.* **1990**, *53*, 634-637. <https://doi.org/10.1021/np50069a015>
- ⁹⁷Heidari, M.R., Vahedian, M., Moamenzadeh, S., Abbasi, M.M. The Analgesic Effect and Possible Mechanism of *Colchicum Szovitsii* Methanolic Extract in Mouse, *J. Rafsanjan Uni. Med. Sci.* **2005**, *4*, 25-33. http://journal.rums.ac.ir/browse.php?a_id=6&sid=1&slc_lang=en
- ⁹⁸Ben-Chetrit, E., Bergmann, S., Sood, R. Mechanism of the anti-inflammatory effect of colchicine in rheumatic diseases: a possible new outlook through microarray analysis, *Rheumatology.* **2006**, *45*, 274-282. doi:10.1093/rheumatology/kei140
- ⁹⁹Deftereos, S., Giannopoulos, G., Angelidis, C., Alexopoulos, N., Filippatos, G., Papoutsidakis, N., Sianos, G., Goudevenos, J., Alexopoulos, D., Pyrgakis, V., Cleman, M.W., Antonis S. Anti-Inflammatory Treatment With Colchicine in Acute Myocardial Infarction A Pilot Study, *Circulation.* **2015**, *132*, 1395-1403. doi: 10.1161/CIRCULATIONAHA.115.017611
- ¹⁰⁰Karagoz, A., Artun, F.T., Ozcan, G., Melikoglu, G., Anil, S., Kultur, S., Sutlupinar, N. *In vitro* evaluation of antioxidant activity of some plant methanol extracts, *Biotechnol.*

- Biotechnol. Equip.* **2015**, *29*, 1184-1189, doi: 10.1080/13102818.2015.1080600
- ¹⁰¹Nair, V., Kumar, R., Singh, S., Gupta, Y.K. Investigation into the anti-inflammatory and Antigranuloma activity of *Colchicum luteum* Baker in experimental models, *Inflammation*. **2012**, *35*, 881-888. doi: 10.1007/s10753-011-9389-2
- ¹⁰²Zhang, L., Hu, J.J.J., Lin, J.W., Fang, W.S., Du, G.H. Anti-inflammatory and analgesic effects of ethanol and aqueous extracts of *Pterocephalus hookeri* (C.B. Clarke) Höeck, *J. Ethnopharmacol.* **2009**, *23*, 510-514. doi: 10.1016/j.jep.2009.01.039
- ¹⁰³Ullah, F., Ayaz, M., Sadiq, A., Hussain, A., Ahmad, S., Imran, M., Zeb, A. Phenolic, flavonoid contents, anticholinesterase and antioxidant evaluation of *Iris germanica* var. florentina, *Nat. Prod. Res.* **2015**, *30*, 1440-1444. doi: 10.1080/14786419.2015.1057585
- ¹⁰⁴Rahman, A., Nasim, S., Baig, I., Jalil, S., Orhan, I., Sener, B., Choudhary, M.I. Anti-inflammatory isoflavonoids from the rhizomes of *Iris germanica*, *J. Ethnopharmacol.* **2003**, *86*, 177-180. doi:10.1016/S0378-8741(03)00055-2
- ¹⁰⁵Wang, D., Du, Q., Li, H., Wang, S. The Isosteroid Alkaloid Imperialine from Bulbs of *Fritillaria cirrhosa* Mitigates Pulmonary Functional and Structural Impairment and Suppresses Inflammatory Response in a COPD-Like Rat Model, *Mediators Inflamm.* **2016**, Article ID 4192483, 17 pages. <http://dx.doi.org/10.1155/2016/4192483>
- ¹⁰⁶Turkoglu, S., Parlak, A.E. Determination of total phenolic and total flavonoid contents and antioxidant capacities of an aquatic plant (*Riccia fluitans*), *Ege. J. Fish Aqua. Sci.* **2014**, *31*, 35-40. doi: 10.12714/egejfas.2014.31.1.06
- ¹⁰⁷Tosun, A., Akkol, E.K., Suntar, I., Kiremit, H.O., Asakawa, Y. Phytochemical investigations and bioactivity evaluation of liverworts as a function of anti-inflammatory and antinociceptive properties in animal models, *Pharm. Biol.* **2013**, *51*, 1008-1013. doi: 10.3109/13880209.2013.774028
- ¹⁰⁸Stajner, D., Popovic, B.M., Kapor, A., Boza, P., Stajner, M. Antioxidant and scavenging capacity of *Anacamptis pyramidalis* L.-pyramidal orchid from Vojvodina, *Phytother. Res.* **2010**, *24*, 759-763. doi: 10.1002/ptr.3041.
- ¹⁰⁹Chinsamy, M., Finnie, J.F., J. Van Staden, J. Anti-inflammatory, antioxidant, anti-cholinesterase activity and mutagenicity of South African medicinal orchids, *S. Afr. J. Bot.* **2014**, *91*, 88-98. <http://dx.doi.org/10.1016/j.sajb.2013.12.004>
- ¹¹⁰A'rmoosh, H., Al-Omari, M. Chapter Three. *The Plants in a Book*, 1st ed., Dar-Alnafa'is, Damascus, Syria, 2004, 391-392.
- ¹¹¹Abdelwahab, M.F., Sangi, S., Arafat, H.H., Ragab, E.A. New Phytochemical Constituent and Bioactivities of *Horwoodia dicksoniae* and *Rumex cyprius*, *Pharmacogn. Mag.* **2016**, *12*, 165-170. doi: 10.4103/0973-1296.186348
- ¹¹²Ali Esmail Al-Snafi, A.E. Chemical Constituents and Medical Importance of *Galium Aparine* – A Review, *Indo Am. j. pharm. Sci.* **2018**, *5*, 1739-1744. doi: 10.5281/zenodo.1210517
- ¹¹³Deliorman, D., Calis, I., Ergun, F. Iridoids from *Galium aparine*, *Pharm. Biol.* **2001**, *39*, 234-235. doi: 10.1076/phbi.39.3.234.5928
- ¹¹⁴He, J., Lu, X., Wei, T., Dong, Y., Cai, Z., Tang, L., Liu, M. Asperuloside and Asperulosidic Acid Exert an Anti-Inflammatory Effect via Suppression of the NF-κB and MAPK Signaling Pathways in LPS-Induced RAW 264.7 Macrophages, *Int. J. Mol. Sci.* **2018**, *19*, 2027-2038. doi: 10.3390/ijms19072027
- ¹¹⁵Zhu, H., Liang, Q.H., Xiong, X.G., Wang, Y., Zhang, Z.H., Sun, M.J., Lu, X., Wu, D. Anti-Inflammatory Effects of p-Coumaric Acid, a Natural Compound of *Oldenlandia diffusa*, on Arthritis Model Rats, *Evid. Based Complementary Altern. Med.* **2018**, Article ID 5198594, 9 pages. <https://doi.org/10.1155/2018/5198594>
- ¹¹⁶Goldschmidt Lins, P., Piccoli Pugine, S.M., Scatolini, A.M., Pires de Melo, M. *In vitro* antioxidant activity of olive leaf extract (*Olea europaea* L.) and its protective effect on oxidative damage in human erythrocytes, *Heliyon*. **2018**, *4*, e00805 (26 pages). doi: 10.1016/j.heliyon.2018.e00805
- ¹¹⁷Mahjoub, R.C., Khemiss, M., Dhidah, M., Dellai, A., A. Bouraoui, A., Khemiss, F. Chloroformic and Methanolic Extracts of *Olea europaea* L. Leaves Present Anti-Inflammatory and Analgesic Activities, *ISRN Pharmacology*. **2011**. Article ID 564972, 5 pages. doi: 10.5402/2011/564972
- ¹¹⁸Gallo, M., Conte, E., Naviglio, D. Analysis and Comparison of the Antioxidant Component of *Portulaca Oleracea* Leaves Obtained by Different Solid-Liquid Extraction Techniques, *Antioxidants*. **2017**, *6*, 64-72. doi:10.3390/antiox6030064
- ¹¹⁹Allahmoradi, E., Taghiloo, S., Omrani-Nava, V., Shobeiri, S.S., Tehrani, M., Ebrahimzadeh, M.A., Asgarian-Omran, H. Anti-inflammatory effects of the *Portulaca oleracea* hydroalcoholic extract on human peripheral blood mononuclear cells, *Med. J. Islam. Repub. Iran.* **2018**, *32*, 6 pages. <https://doi.org/10.14196/mjiri.32.80>
- ¹²⁰Azab, A. *Micromeria*: Chemistry and Medicinal Activities, *Eur. Chem. Bull.* **2016**, *5*, 300-307. doi: 10.17628/ECB.2016.5.300
- ¹²¹Azab, A. *Alcea*: Traditional Medicine, Current Research and Future Opportunities, *Eur. Chem. Bull.* **2016**, *5*, 505-514. doi: 10.17628/ECB.2016.5.505
- ¹²²Azab, A. *Arum*: a Plant Genus with Great Medicinal Potential, *Eur. Chem. Bull.* **2017**, *6*, 59-68. doi: 10.17628/ecb.2017.6.59-68
- ¹²³Azab, A. *Malva*: Food, Medicine, Chemistry, *Eur. Chem. Bull.* **2017**, *6*, 295-320. doi: 10.17628/ecb.2017.6.295-320
- ¹²⁴Azab, A. Carob (*Ceratonia siliqua*): Health, Medicine, Chemistry, *Eur. Chem. Bull.* **2017**, *6*, 456-469. doi: 10.17628/ecb.2017.6.456-469
- ¹²⁵Azab, A., Nassar, A., Azab, A.N., Anti-inflammatory Activity of Natural Products, *Molecules*. **2016**, *21*, 1321-1330. doi: 10.3390/molecules21101321
- ¹²⁶Alkofahi, A.S., Alzoubi, K.H., Omar F. Khabour, O.F., Mhaidat, N.M. Screening of selected medicinal plants from Jordan for their protective properties against oxidative DNA damage, *Ind. Crop. Prod.* **2016**, *88*, 106-111. <http://dx.doi.org/10.1016/j.indcrop.2016.02.059>
- ¹²⁷Azuma, K., Nakayama, M., Koshioka, M., Ippoushi, K., Yamaguchi, Kohata, K., Yamauchi, Y., Ito, H., Higashio, H. Phenolic Antioxidants from the Leaves of *Corchorus olitorius* L., *J. Agric. Food Chem.* **1999**, *47*, 3963-3966. doi: 10.1021/jf990347p
- ¹²⁸Owoyele, B.V., Oyewole, A.L., Alimi, M.L., Sanni, S.A., Oyeleke, S.A. Anti-inflammatory and antipyretic properties of *Corchorus olitorius* aqueous root extract in Wistar rats, *J. Basic Clin. Physiol. Pharmacol.* **2015**, *26*, 363-368. doi: 10.1515/jbcpp-2013-0166.
- ¹²⁹Del Rio, M., Font, R., Almela, C., Velez, D., Montoro, R., De Haro Bailon, A. Heavy metals and arsenic uptake by wild vegetation in the Guadiamar river area after the toxic spill of the Aznalcollar mine, *J. Biotechnol.*, **2002**, *98*, 125-137. doi: 10.1016/s0168-1656(02)00091-3
- ¹³⁰Molyneux, R.J., Ralphs, M.H. Plant Toxines and Palatability to Herbivores. *J. Range Manage.* **1992**, *45*, 13-18. doi: 10.2307/4002519
- ¹³¹Cortinovis, C., Caloni, F. Alkaloid-Containing Plants Poisonous to Cattle and Horses in Europe, *Toxins*. **2015**, *7*, 5301-5307. doi:10.3390/toxins7124884
- ¹³²Azab, A., Nassar, A., Kaplanski, J., Mahajneh, R., Agam, G., Azab, A. N. Effects of aqueous extract of *Notobasis syriaca* on lipopolysaccharide-induced inflammation in rats, *Asian Pac. J. Trop. Med.* **2018**, *11*, 48-52. doi: 10.4103/1995-7645.225353

¹³³Azab, A. Total Phenolic Content, Antioxidant Capacity and Antifungal Activity of Extracts of *Carthamus tenuis* and *Cephalaria joppensis*, *Eur. Chem. Bull.* **2018**, *7*, 156-161. doi: 10.17628/ecb.2018.7.156-161

¹³⁴Azab, A. A Facile Method for Testing Antioxidant Capacity and Total Phenolic Content of *Notobasis syriaca* and *Scolymus maculatus* extracts and Their Antifungal Activity, *Eur. Chem. Bull.* **2018**, *7*, 210-217. doi: 10.17628/ecb.2018.7.210-217

¹³⁵Azab, A. Antifungal and anti-termite activities, total phenolic content of *Prosopis Farcta* extracts, and attempts to develop weed biocontrol method against it. *Eur. Chem. Bull.* **2018**, *7*, 293-302. doi: 10.17628/ecb.2018.7.293-302

Received: 08.07.2019.

Accepted: 04.08.2019.