



PREVALENCE OF DENTURE STOMATITIS AND ITS ASSESSMENT TOWARDS MANAGEMENT -A RETROSPECTIVE STUDY

Vatika Agarwal^[a], Subhabrata Maiti^{[b]*}

Article History: Received: 26.02.2022

Revised: 25.03.2022

Accepted: 24.04.2022

Abstract: Denture stomatitis is a common oral condition of multifactorial etiology. Factors such as age, gender, oral and denture hygiene may play a role in predisposition to denture stomatitis. The aim of this study was to evaluate these factors and understand the various modalities of treatment for effective management of denture stomatitis with respect to the South Indian population. Case sheets of around 26,000 patients were reviewed from March 2019 to June 2020 out of which a sample size of 56 patients were included. To eliminate bias all patients affected by the disease, who had completed treatment were included in the study. Epidemiological data of the patient along with their ongoing treatment was collected and tabulated in MS Excel sheet. The data was then analysed using IBM SPSS software version 23. The prevalence of denture stomatitis in females was 57.1% while in males was 42.9%. Age group of 51-60 years and 61-70 years was found to be equally and most commonly affected. Palate was found to be most commonly affected. The most common treatment modality used was pharmacotherapy using antifungal drugs. The increasing prevalence of denture stomatitis amongst the older population is a matter of great concern. The present study shows the prevalence of denture stomatitis in females especially in the old age population. Thus it is important to increase awareness amongst the patients regarding the condition and effective maintenance of oral and denture hygiene and to increase awareness amongst the dentists regarding the various treatment modalities to effectively prevent and clinically manage such conditions.

Keywords: Denture stomatitis, Candida Albicans, Denture hygiene, oral hygiene, pharmacotherapy, innovative.

[a]. Saveetha Dental College and Hospitals. Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-600077, Tamil Nadu, India

[b]. Senior Lecturer, Department of Prosthodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-600077, Tamil Nadu, India

*Corresponding Author

Email: 151909003.sdc@saveetha.com, subhabratamaiti.sdc@saveetha.com.

DOI: 10.31838/ecb/2022.11.02.006

INTRODUCTION

Inflammation of the mucosa underlying a complete denture is called denture stomatitis (Yarborough *et al.*, 2016). It is a disorder frequently seen in elderly patients (Cunha-Cruz, 2006), having a prevalence of 15%-70% amongst all denture wearers (Gendreau and Loewy, 2011). Frequent signs associated with denture stomatitis may be inflamed red mucosa, soft tissue hyperplasia, angular cheilitis, glossitis, burning sensation in the palate (Ritchie *et al.*, 1969) and less frequently bone resorption (Jeganathan and Lin, 1992). It is rarely associated with severe pain and discomfort (Yarborough *et al.*, 2016) (Ariga *et al.*, 2018). Denture stomatitis has a characteristic appearance which can be classified using the Newton classification system (Newton, 1975). It is classified into 3 types:

Type 1- localized inflammation or pinpoint hyperemia

Type 2 -diffuse erythema associated with the denture contacting mucosa

Type 3 - papillary hyperplasia of the keratinized mucosa

If untreated, the disease usually progresses from Type I through Type II to Type III. Type I is usually related to denture trauma only, as compared with Type II and III which have been associated with a multi-factorial aetiology (Budtz-Jørgensen and Bertram, 1970) (Jyothi *et al.*, 2017). Multiple factors are associated with denture stomatitis, however microbial pathogenesis is most frequently implicated as its cause (Girard, Landry and Giasson, 1996) (Duraisamy *et al.*, 2019). Multiple organisms like *C. albicans*, *C. glabrata*, *C. tropicalis*, and other fungi from the *Candida* species have been associated with denture stomatitis (Budtz-Jørgensen, Stenderup and Grabowski, 1975; Girard, Landry and Giasson, 1996), however *Candida albicans* is considered as its primary cause (Allison and Douglas, 1973). Other factors like trauma due to ill-fitting dentures (Nyquist, 1953) (Selvan and Ganapathy, 2016), xerostomia, poor dental hygiene, compromised immunity, nocturnal use of dentures too plays a significant role (Altarawneh *et al.*, 2013) (Ganapathy *et al.*, 2016).

Denture stomatitis is a chronic disorder, treatment of which is essential for the well being of patients. If not treated with care, denture stomatitis can cause severe systemic disorders like deep tissue infections especially in patients with comorbidities like HIV-AIDS, nutritional deficiency, uncontrolled diabetes (Shepherd, 1986) (Subasree, Murthykumar and Dhanraj, 2016). According to (Budtz-Jørgensen and Bertram, 1970; Abelson, 1981) ill fitting dentures, poor oral and denture hygiene and trauma were the main causes of denture stomatitis hence these factors had to be taken into consideration during its treatment. Commonly used treatment modalities currently include maintenance of denture and oral hygiene (Fisher, 1956) (Ranganathan, Ganapathy and Jain, 2017), correction of denture faults or change of faulty denture and pharmaceutical management using anti-fungal medication and mouth washes. (Preti *et al.*,

1996)(Vijayalakshmi and Ganapathy, 2016) suggested that preventive measures like implant supported dentures can be considered to allow even distribution of force, thereby causing less trauma to the underlying tissue and allowing even distribution of force. Treatment planned should be carried out properly as relapse of disease is very high in immunocompromised individuals (Greenspan and Greenspan, 1991)(Ganapathy, Kannan and Venugopalan, 2017). Previously our team has a rich experience in working on various research projects across multiple disciplines (Ramesh Kumar *et al.*, 2011; Jain, Kumar and Manjula, 2014; Krishnan, Pandian and Kumar S, 2015; Keerthana and Thenmozhi, 2016; Sivamurthy and Sundari, 2016; Felicita, 2017a, 2017b; Kumar, 2017; Sekar *et al.*, 2019; Johnson *et al.*, 2020). Now the growing trend in this area motivated us to pursue this project. The aim of this research is to study the prevalence and treatment modality of denture stomatitis in a well reputed hospital in Chennai.

Data of patients visiting Saveetha Oral medicine department June 2019- March 2020 was collected by method of simple random sampling. A total of 26,000 case sheets were reviewed out of which 61 patients had denture stomatitis. 5 patients were eliminated from the study as 1 was not a regular denture wearer, 2 only wore upper dentures and treatment of 2 patients wasn't completed. Cross verification of error in data was done via telephonic conversation. Measure taken to minimise sampling bias was that all case sheets of patients undergoing treatment for denture stomatitis were included. External validity is that it is applicable to the South Indian population.

The data collected was entered in MS Excel spreadsheet and tabulated. The data was imported in spss software version 26 and variables were defined. Statistical analysis of data was carried out using unpaired t test. Independent variable was denture stomatitis and dependent variable was age and sex. Data transfer was done using spss software version 26.

MATERIALS AND METHOD

This study was carried out in a university setting at Saveetha Dental College and hospital, Chennai, India by the department of Prosthodontics and Implantology. The study consisted of a data collector and 1 data reviewer. The disadvantage of the study was its geographical limitation. This study was approved by the institute ethical board committee.

RESULTS AND DISCUSSION

A total of 26,000 case sheets were reviewed out of which 61 were found to be suffering from Denture stomatitis and 56 were included in the study as 5 patients were eliminated from the study. Amongst the patients 1 was not a regular denture wearer, 2 only wore upper dentures and treatment of 2 patients wasn't completed.



Figure 1. Pie Chart depicting prevalence of denture stomatitis in males and females. The prevalence is greater in females than males. Of all the patients affected by denture stomatitis 57.14% are females and 47.88% are males.




Figure 2. Bar Graph depicting prevalence of denture stomatitis in various age groups. X-axis represents the age of the individual and Y-axis represents the frequency of individuals affected by denture stomatitis. It is observed that the most common age group affected by the disease was 51-60 and 61-70 years (26.79% each)




Figure 3. Bar Graph depicting the prevalence of denture stomatitis according to region. The X-axis represents the various regions affected, while the Y-axis represents the percentage of individuals affected in the given regions. It was observed that the upper arch was most commonly affected (92.85%).




Figure 4. Bar graph depicting popularity of various treatment modalities used in the management of Denture Stomatitis. X-axis represents the various treatment modalities used in management of denture stomatitis. Y-axis represents the percentage of patients undergoing the given treatment modalities. It is observed that the most common treatment was the use of antifungal drugs(51.69%).




Figure 5. Bar graph depicting the correlation between Age and Gender in Denture stomatitis patients. The X-axis represents the age groups and the Y-axis the gender. The blue represents male and green female. Chi square test showed Pearson Chi Square value 5.56 and P value- 0.339. Since the p value >0.05 there is no statistically significant correlation between Age and Gender in Denture Stomatitis patients, however it is observed that the most predominantly affected group is females of age group (41-50 years)

It is observed that the prevalence of denture stomatitis was greater in females compared to males. Amongst all patients suffering from denture stomatitis 57.1% were female while 42.9 were males(Figure 1).

In the present study it is observed that there is equal prevalence present in the age groups 51-60 years and 61-70 years of age. These two age groups showed the highest prevalence. While the least affected group was of patients 31-40 years.(Figure 2)It was also observed that 92.9% of denture stomatitis cases affected only the upper arch, while the rest affected both upper and lower. No cases were reported affecting only the lower arch. individuals affected by denture stomatitis(Figure 3)It is observed that the popularity of various treatment modalities in the management of Denture Stomatitis, was pharmacotherapy using Antifungals like clotrimazole and nystatin.. (Figure 4)

In the present study it was observed that the prevalence of denture stomatitis was greater in females compared to male. Similar results were also seen in a study conducted by (Junior *et al.*, 1991)(Arendorf and Walker, 1987). The reason may be that women have a higher life expectancy than men, and undergo dental extractions more frequently than men due to higher caries index(Shaffer *et al.*, 2015)(Ashok and Suvitha, 2016), hence their need for a dental prosthesis may be higher, resulting in higher cases of denture related stomatitis. According to (Nyquist, 1952)(Ashok *et al.*, 2014) greater prevalence in women can also be explained due to hormonal changes women undergo during menopausal period predisposing them to infections, this hypothesis may be supported as a great number of female complete denture patients are found to be undergoing menopause. It has been suggested that increased prevalence may also be due to the fact that female patients wear their dentures more often and perhaps for longer periods of time for esthetic purposes(Cutright, 1974)(Venugopalan *et al.*, 2014). Many other investigators too have concluded that denture stomatitis is more common in females(Nyquist, 1952; Fisher, 1956) (Davenport, 1970; Budtz-Jørgensen and Løe, 1972)(Nater *et al.*, 1978) However, there have been a few reports in which the two sexes were equally affected(Bergman, Carlsson and Hedegard, 1964; Grabowski and Bertram, 1975)(Bergman, Carlsson and Hedegard, 1964; Ettinger, 1975)(Ettinger, 1975). Denture stomatitis is typically present in the older age groups.In the present study most patients affected by it were present within the age group 51-60 years and 61-70 years. Similar results were found in a study conducted by (Axéll, 1976)(Kannan and Venugopalan, 2018). This may be due to the fact that the elderly are the most likely population to wear dentures and are at particular risk of poor denture hygiene as they have limited financial resources ,manual dexterity, information and poorer vision. Poverty also influences the ability to visit a dentist regularly and replace ill-fitting dentures, which causes trauma to the underlying tissue predisposing them to the disease. Decreased immunity due to age may also play an important role. However the number of patients affected by denture stomatitis greater than 70 years of age, were less as the average lifespan in India is 68.7 years(National Health Report-2019), hence the number of patients in the age group greater than 70 would be lesser. Denture stomatitis was most commonly seen affecting the palate in the present study, similar results were found in a study conducted by(Figueiral *et al.*, 2007)(Basha, Ganapathy and Venugopalan, 2018). Palate is most commonly affected probably due to the large contact area between denture and oral mucosa, while the mandible is less affected due to the washing action of saliva.

(Girard, Landry and Giasson, 1996) had conducted a study in which it was observed that the primary cause of denture stomatitis is microbial pathogenesis, and the most commonly present species found was *Candida albicans*. Even if the cause of denture stomatitis may be prolonged hours of denture wear, or the use of old worn out dentures the primary line of treatment is usually pharmacotherapy using antifungals due to the increased presence of candida species in such patients. In the present study too the most popular line of treatment was the use of antifungals amongst which the use of clotrimazole was more popular than that of nystatin as it had better taste and less gastrointestinal adverse reactions(Kaur and Kakkar, 2010), however many still advocate the use of nystatin over clotrimazole due its high efficacy, low cost, and less side effects(Sklenár *et al.*, 2013)(Ajay *et al.*, 2017). Although most cases resolve in a span of 12-14 days with the use of topical antifungals the use of systemic antifungals can be recommended to immunocompromised individuals or patients with poor compliance(Dias, Samaranayake and Lee, 1997). Denture hygiene plays a significant role in the prevention of denture stomatitis(Sadamori *et al.*, 1990)(Marinoski, Bokor-Bratić and Čanković, 2014)(Kulak-Ozkan, Kazazoglu and Arikan, 2002), hence it is required to use denture cleaners such as hypochlorite solution and chlorhexidine to disinfect the dentures and decrease plaque and microbial accumulation on the denture surface.In the present study we observe the use of hypochlorite solution for the disinfection of dentures and chlorhexidine for the maintenance of oral hygiene. Hypochlorite solution although is an effective disinfecting agent(Orsi *et al.*, 2011) its use is limited because it whitens acrylic resins(Paranhos *et al.*, 2013)and corrodes metal components of prostheses(Paranhos *et al.*, 2014). A study done by (Pavarina *et al.*, 2003) had proved the efficacy of chlorhexidine as a denture cleaner, hence chlorhexidine solution can be used as a denture disinfectant at a concentration of 4% while as a mouthwash for oral hygiene at a concentration of 0.12%. (Nyquist, 1952) along with several other workers(Nyquist, 1952; Turrell, 1966)(Niesluchowska *et al.*, 1980) had claimed the role of trauma and ill fitting dentures in progression of denture stomatitis. (Samaranayake, McCourtie and MacFarlane, 1980) suggested that inflammatory exudate leaking from a traumatized palatal mucosa promotes candidal colonization of the denture surface. Hence it was essential to either replace ill fitting dentures or rebase them. In the present study it is observed that replacement of ill fitting dentures was more advisable than their rebasing, to overcome problems such as crazing and attrited teeth seen in old dentures. The limitations of this study were that it was geographically limited. The sample size was also less. Hence studies with larger sample size and a longer duration of follow up are required to achieve conclusive data.

Denture stomatitis is a chronic disorder, which if not managed well may cause deep tissue infections. It also has high recurrence in patients with comorbidities. Hence this study was important to understand the prevalence of the disorder and educate clinicians regarding its effective management. Since prevention is better than cure it is advised to carry out effective prevention programs which include(Hoad-Reddick, Grant and Griffiths, 1990) routine inspection of the oral cavity disorders,educating patients regarding proper denture wearing, denture sanitation and oral hygiene habits. Patients with partial dentures should undergo periodic professional plaque control procedures. Prosthodontists should ensure that the dentures they deliver have good fit and polish so as to reduce trauma to denture

bearing tissues and to decrease microbial load on denture.viii. Patients who are susceptible should be given softliners only with incorporation of antifungals like nystatin as soft-liners are more porous than acrylic and may have increased adherence to the fungi. However if precautionary methods ix. fail, it is essential to manage the disease well to allow the patient to enjoy a good quality of life. The present study helps x. us understand the prevalence of the disease to allow prevention, as well as explains the advantages and disadvantages of various popularly used treatment modalities to allow effective management of the disease, xi. increasing clinician and patient satisfaction. Our institution is passionate about high quality evidence based research and has excelled in various fields (Pc, Marimuthu and Devadoss, 2018; Ramesh *et al.*, 2018; Ezhilarasan, Apoorva and Ashok xii. Vardhan, 2019; Ramadurai *et al.*, 2019; Sridharan *et al.*, 2019; Vijayashree Priyadharsini, 2019; Mathew *et al.*, 2020). We hope this study adds to this rich legacy.

CONCLUSION

Denture stomatitis is a chronic, debilitating disease effective xiv. management of which is important. In present study prevalence of denture stomatitis is greater in females than males, and is commonly found in the older population. The high prevalence of this disease is quite alarming hence it is xv. essential to increase awareness. General dentists should be knowledgeable and familiar with the etiopathogenesis, clinical presentation, diagnosis and management of this condition, while prosthodontists should xvi. take care in the fabrication of dentures and should educate the patient well regarding its use.

REFERENCES

- i. Abelson, D.C. (1981) 'Denture plaque and denture cleansers', *The Journal of prosthetic dentistry*, 45(4), pp. 376–379.
- ii. Ajay, R. *et al.* (2017) 'Effect of Surface Modifications on the Retention of Cement-retained Implant Crowns xix. under Fatigue Loads: An In vitro Study', *Journal of pharmacy & bioallied sciences*, 9(Suppl 1), pp. S154–S160.
- iii. Allison, R.T. and Douglas, W.H. (1973) 'Micro-colonization of the denture-fitting surface by Candida xx. albicans', *Journal of dentistry*, 1(5), pp. 198–201.
- iv. Altarawneh, S. *et al.* (2013) 'Clinical and histological findings of denture stomatitis as related to intraoral colonization patterns of Candida albicans, salivary xxi. flow, and dry mouth', *Journal of prosthodontics: official journal of the American College of Prosthodontists*, 22(1), pp. 13–22.
- v. Arendorf, T.M. and Walker, D.M. (1987) 'Denture stomatitis: a review', *Journal of Oral Rehabilitation*, pp. 217–227. doi:10.1111/j.1365-2842.1987.tb00713.x. xxii.
- vi. Ariga, P. *et al.* (2018) 'Determination of Correlation of Width of Maxillary Anterior Teeth using Extraoral and Intraoral Factors in Indian Population: A Systematic Review', *World Journal of Dentistry*, pp. 68–75 xxiii. doi:10.5005/jp-journals-10015-1509.
- vii. Ashok, V. *et al.* (2014) 'Lip Bumper Prosthesis for an Acromegaly Patient: A Clinical Report', *Journal of Indian Prosthodontic Society*, 14(Suppl 1), pp. 279–282. xxiv.
- Ashok, V. and Suvitha, S. (2016) 'Awareness of all ceramic restoration in rural population', *Research Journal of Pharmacy and Technology*, p. 1691. doi:10.5958/0974-360x.2016.00340.1.
- Axéll, T. (1976) *A prevalence study of oral mucosal lesions in an adult Swedish population.*
- Basha, F.Y.S., Ganapathy, D. and Venugopalan, S. (2018) 'Oral Hygiene Status among Pregnant Women', *Research Journal of Pharmacy and Technology*, p. 3099. doi:10.5958/0974-360x.2018.00569.3.
- Bergman, B., Carlsson, G.E. and Hedegard, B. (1964) 'A LONGITUDINAL TWO-YEAR STUDY OF A NUMBER OF FULL DENTURE CASES', *Acta odontologica Scandinavica*, 22, pp. 3–26.
- Budtz-Jørgensen, E. and Løe, H. (1972) 'Chlorhexidine as a denture disinfectant in the treatment of denture stomatitis', *Scandinavian journal of dental research*, 80(6), pp. 457–464.
- Budtz-Jørgensen, E., Stenderup, A. and Grabowski, M. (1975) 'An epidemiologic study of yeasts in elderly denture wearers', *Community dentistry and oral epidemiology*, 3(3), pp. 115–119.
- Budtz-Jørgensen, E. and Bertram, U. (1970) 'Denture Stomatitis: II. The Effect of Antifungal and Prosthetic Treatment', *Acta Odontologica Scandinavica*, pp. 283–304. doi:10.3109/00016357009032036.
- Cunha-Cruz, J. (2006) 'One in 3 removable denture users in the United States has denture stomatitis', *The journal of evidence-based dental practice*, pp. 197–198.
- Cutright, D.E. (1974) 'The histopathologic findings in 583 cases of epulis fissuratum', *Oral surgery, oral medicine, and oral pathology*, 37(3), pp. 401–411.
- Davenport, J.C. (1970) 'The oral distribution of candida in denture stomatitis', *British Dental Journal*, pp. 151–156. doi:10.1038/sj.bdj.4802540.
- Dias, A.P., Samaranayake, L.P. and Lee, M.T. (1997) 'Miconazole lacquer in the treatment of denture stomatitis: clinical and microbiological findings in Chinese patients', *Clinical oral investigations*, 1(1), pp. 47–52.
- Duraisamy, R. *et al.* (2019) 'Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments', *Implant dentistry*, 28(3), pp. 289–295.
- Ettinger, S. (1975) 'Übersetzen aus Übersetzungen?', *Babel Revue internationale de la traduction / International Journal of Translation*, pp. 187–189. doi:10.1075/babel.21.4.13ett.
- Ezhilarasan, D., Apoorva, V.S. and Ashok Vardhan, N. (2019) 'Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(2), pp. 115–121.
- Felicita, A.S. (2017a) 'Orthodontic management of a dilacerated central incisor and partially impacted canine with unilateral extraction - A case report', *The Saudi dental journal*, 29(4), pp. 185–193.
- Felicita, A.S. (2017b) 'Quantification of intrusive/retraction force and moment generated during en-masse retraction of maxillary anterior teeth using mini-implants: A conceptual approach', *Dental press journal of orthodontics*, 22(5), pp. 47–55.
- Figureiral, M.H. *et al.* (2007) 'Denture-related

- stomatitis: identification of aetiological and predisposing factors - a large cohort', *Journal of oral rehabilitation*, 34(6), pp. 448–455.
- xxv. Fisher, A.A. (1956) 'Allergic sensitization of the skin and oral mucosa to acrylic resin denture materials', *The Journal of Prosthetic Dentistry*, pp. 593–602. doi:10.1016/0022-3913(56)90003-8.
- xxvi. Ganapathy, D. et al. (2016) 'Effect of Resin Bonded Luting Agents Influencing Marginal Discrepancy in All Ceramic Complete Veneer Crowns', *Journal of clinical and diagnostic research: JCDR*, 10(12), pp. ZC67–ZC70.
- xxvii. Ganapathy, D.M., Kannan, A. and Venugopalan, S. (2017) 'Effect of Coated Surfaces influencing Screw Loosening in Implants: A Systematic Review and Meta-analysis', *World Journal of Dentistry*, pp. 496–502. doi:10.5005/jp-journals-10015-1493.
- xxviii. Gendreau, L. and Loewy, Z.G. (2011) 'Epidemiology and etiology of denture stomatitis', *Journal of prosthodontics: official journal of the American College of Prosthodontists*, 20(4), pp. 251–260.
- xxix. Girard, B., Jr, Landry, R.G. and Giasson, L. (1996) '[Denture stomatitis: etiology and clinical considerations]', *Journal*, 62(10), pp. 808–812.
- xxx. Grabowski, M. and Bertram, U. (1975) 'Oral health status and need of dental treatment in the elderly Danish population', *Community dentistry and oral epidemiology*, 3(3), pp. 108–114.
- xxxi. Greenspan, D. and Greenspan, J.S. (1991) 'Management of the oral lesions of HIV infection', *Journal of the American Dental Association*, 122(8), pp. 26–32.
- xxxii. Hoad-Reddick, G., Grant, A.A. and Griffiths, C.S. (1990) 'Investigation into the cleanliness of dentures in an elderly population', *The Journal of Prosthetic Dentistry*, pp. 48–52. doi:10.1016/0022-3913(90)90152-3.
- xxxiii. Jain, R.K., Kumar, S.P. and Manjula, W.S. (2014) 'Comparison of intrusion effects on maxillary incisors among mini implant anchorage, j-hook headgear and utility arch', *Journal of clinical and diagnostic research: JCDR*, 8(7), pp. ZC21–4.
- xxxiv. Jeganathan, S. and Lin, C.C. (1992) 'Denture stomatitis - a review of the aetiology, diagnosis and management', *Australian Dental Journal*, pp. 107–114. doi:10.1111/j.1834-7819.1992.tb03046.x.
- xxxv. Johnson, J. et al. (2020) 'Computational identification of MiRNA-7110 from pulmonary arterial hypertension (PAH) ESTs: a new microRNA that links diabetes and PAH', *Hypertension research: official journal of the Japanese Society of Hypertension*, 43(4), pp. 360–362.
- xxxvi. Junior, J.J. et al. (1991) 'Oral mucosal health and disease in institutionalized elderly in Brazil', *Community Dentistry and Oral Epidemiology*, pp. 173–175. doi:10.1111/j.1600-0528.1991.tb00136.x.
- xxxvii. Jyothi, S. et al. (2017) 'Periodontal health status of three different groups wearing temporary partial denture', *Research Journal of Pharmacy and Technology*, 10(12), pp. 4339–4342.
- xxxviii. Kannan, A. and Venugopalan, S. (2018) 'A systematic review on the effect of use of impregnated retraction cords on gingiva', *Research Journal of Pharmacy and Technology*, p. 2121. doi:10.5958/0974-360x.2018.00393.1.
- xxxix. Kaur, I.P. and Kakkar, S. (2010) 'Topical delivery of antifungal agents', *Expert opinion on drug delivery*, 7(11), pp. 1303–1327.
- Keerthana, B. and Thenmozhi, M.S. (2016) 'Occurrence of foramen of huschke and its clinical significance', *Research Journal of Pharmacy and Technology*, 9(11), pp. 1835–1836.
- Krishnan, S., Pandian, S. and Kumar S, A. (2015) 'Effect of bisphosphonates on orthodontic tooth movement-an update', *Journal of clinical and diagnostic research: JCDR*, 9(4), pp. ZE01–5.
- Kulak-Ozkan, Y., Kazazoglu, E. and Arikan, A. (2002) 'Oral hygiene habits, denture cleanliness, presence of yeasts and stomatitis in elderly people', *Journal of oral rehabilitation*, 29(3), pp. 300–304.
- Kumar, S. (2017) 'The emerging role of botulinum toxin in the treatment of orofacial disorders: Literature update', *Asian journal of pharmaceutical and clinical research*, 10(9), p. 21.
- Marinoski, J., Bokor-Bratić, M. and Čanković, M. (2014) 'Is denture stomatitis always related with candida infection? A case control study', *Medicinski glasnik: official publication of the Medical Association of Zenica-Doboj Canton, Bosnia and Herzegovina*, 11(2), pp. 379–384.
- Mathew, M.G. et al. (2020) 'Evaluation of adhesion of Streptococcus mutans, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary ...', *Clinical oral investigations* [Preprint]. Available at: <https://link.springer.com/article/10.1007/s00784-020-03204-9>.
- Nater, J.P. et al. (1978) 'Etiologic factors in denture sore mouth syndrome', *The Journal of Prosthetic Dentistry*, pp. 367–373. doi:10.1016/0022-3913(78)90114-2.
- Newton, A.V. (1975) 'The difficult denture patient. A review of psychological aspects', *British Dental Journal*, pp. 93–97. doi:10.1038/sj.bdj.4803388.
- Niesluchowska, M. et al. (1980) '[Some parameters of saliva in cases of denture stomatitis infected with Candida albicans]', *Protetyka stomatologiczna*, 30(4-5), pp. 299–302.
- Nyquist, G. (1952) 'A study of denture sore mouth; an investigation of traumatic, allergic and toxic lesions of the oral mucosa arising from the use of full dentures', *Acta odontologica Scandinavica. Supplementum*, 10(9), pp. 1–154.
- Nyquist, G. (1953) 'The influence of denture hygiene and the bacterial flora on the condition of the oral mucosa in full denture cases', *Acta Odontologica Scandinavica*, pp. 24–60. doi:10.3109/00016355308997700.
- Orsi, I.A. et al. (2011) 'Evaluation of the efficacy of chemical disinfectants for disinfection of heat-polymerised acrylic resin', *Gerodontology*, pp. 253–257. doi:10.1111/j.1741-2358.2010.00400.x.
- Paranhos, H. de F.O. et al. (2013) 'Color Stability, Surface Roughness and Flexural Strength of an Acrylic Resin Submitted to Simulated Overnight Immersion in Denture Cleansers', *Brazilian Dental Journal*, pp. 152–156. doi:10.1590/0103-6440201302151.
- Paranhos, H. de F.O. et al. (2014) 'Effect of cleanser solutions on the color of acrylic resins associated with titanium and nickel-chromium alloys', *Brazilian Oral Research*, pp. 1–7. doi:10.1590/1807-3107bor-2014.vol28.0017.
- Pavarina, A.C. et al. (2003) 'Effects of chemical disinfectants on the transverse strength of denture base acrylic resins', *Journal of Oral Rehabilitation*, pp.

- 1085–1089. doi:10.1046/j.1365-2842.2003.01150.x. lxxv.
- iv. Pc, J., Marimuthu, T. and Devadoss, P. (2018) 'Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study', *Clinical implant dentistry and related research* [Preprint]. Available at <https://europepmc.org/article/med/29624863>. atlxvi.
- lvi. Preti, G. *et al.* (1996) 'Histological changes in edentulous oral mucosa under implant-supported overdentures', *Journal of oral rehabilitation*, 23(10)lxvii. pp. 651–654.
- lvii. Ramadurai, N. *et al.* (2019) 'Effectiveness of 2% Articaine as an anesthetic agent in childreklxviii. randomized controlled trial', *Clinical oral investigations*, 23(9), pp. 3543–3550.
- lviii. Ramesh, A. *et al.* (2018) 'Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study', *Journal oflxix. periodontology*, 89(10), pp. 1241–1248.
- lix. Ramesh Kumar, K.R. *et al.* (2011) 'Depth of resin penetration into enamel with 3 types of enamellxx. conditioning methods: a confocal microscopic study', *American journal of orthodontics and dentofacial orthopedics: official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics*, 140(4), pp. 479–485. lxxi.
- lx. Ranganathan, H., Ganapathy, D.M. and Jain, A.R. (2017) 'Cervical and Incisal Marginal Discrepancy in Ceramic Laminate Veneering Materials: A SEM Analysis', *Contemporary clinical dentistry*, 8(2), pp. 272–278.
- lxi. Ritchie, G.M. *et al.* (1969) 'The etiology, exfoliative cytology, and treatment of denture stomatitis', *Thelxiii. Journal of Prosthetic Dentistry*, pp. 185–200. doi:10.1016/0022-3913(69)90246-7.
- lxii. Sadamori S. *et al.* (1990) '[Clinical survey on denturelxiv. stomatitis. 2. The relation between the maintenance of denture and denture stomatitis]', *Nihon Hotetsu Shika Gakkai zasshi*, 34(1), pp. 202–207. lxxv.
- lxiii. Samaranayake, L.P., McCourtie, J. and MacFarlane, T.W. (1980) 'Factors affecting the in-vitro adherence of *Candida albicans* to acrylic surfaces', *Archives of Oral Biology*, pp. 611–615. doi:10.1016/0003-9969(80)90076-x. lxxvi.
- lxiv. Sekar, D. *et al.* (2019) 'Methylation-dependent circulating microRNA 510 in preeclampsia patients', *Hypertension research: official journal of the Japanese Society of Hypertension*, 42(10), pp. 1647–1648.
- Selvan, S.R. and Ganapathy, D. (2016) 'Efficacy of fifth generation cephalosporins against methicillin-resistant *Staphylococcus aureus*-A review', *Research Journal of Pharmacy and Technology*, 9(10), pp. 1815–1818.
- Shaffer, J.R. *et al.* (2015) 'Caries Experience Differs between Females and Males across Age Groups in Northern Appalachia', *International journal of dentistry*, 2015, p. 938213.
- Shepherd, M.G. (1986) 'The pathogenesis and host defence mechanisms of oral candidosis', *The New Zealand dental journal*, 82(369), pp. 78–81.
- Sivamurthy, G. and Sundari, S. (2016) 'Stress distribution patterns at mini-implant site during retraction and intrusion—a three-dimensional finite element study', *Progress in orthodontics*, 17(1), pp. 1–11.
- Sklenár, Z. *et al.* (2013) 'Compounded preparations with nystatin for oral and oromucosal administration', *Acta poloniae pharmaceutica*, 70(4), pp. 759–762.
- Sridharan, G. *et al.* (2019) 'Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(4), pp. 299–306.
- Subasree, S., Murthykumar, K. and Dhanraj (2016) 'Effect of Aloe Vera in Oral Health-A Review', *Research Journal of Pharmacy and Technology*, p. 609. doi:10.5958/0974-360x.2016.00116.5.
- Turrell, A.J. (1966) 'Aetiology of inflamed upper denture-bearing tissues', *British dental journal*, 120(11), pp. 542–546.
- Venugopalan, S. *et al.* (2014) 'Magnetically retained silicone facial prosthesis', *Nigerian journal of clinical practice*, 17(2), pp. 260–264.
- Vijayalakshmi, B. and Ganapathy, D. (2016) 'Medical management of cellulitis', *Research Journal of Pharmacy and Technology*, 9(11), pp. 2067–2070.
- Vijayashree Priyadharsini, J. (2019) 'In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens', *Journal of periodontology*, 90(12), pp. 1441–1448.
- Yarborough, A. *et al.* (2016) 'Evidence Regarding the Treatment of Denture Stomatitis', *Journal of prosthodontics: official journal of the American College of Prosthodontists*, 25(4), pp. 288–301.