

CRUCIFERÆ INDEHISCENTES
(LOMENTACEÆ & NUCAMENTACEÆ) FLORÆ EUROPÆÆ

Auctore VICTORBE DE JANKA.

1. (MORISIA) Herba deprese rosulata subacaulis; scapi radicales 1-flori, fructiferi deflexi, contorti; flores proportione magni, flavi; folia subruncinato-pinnatiloba scapis multo longiora; siliquæ transverse 2-articulatæ, articuli æquales vel superior minor; stylus crassiusculus leviter declinatus *Morisia hypogaea* J. Gay
Manifeste cauliferæ --- --- --- --- --- --- --- --- --- --- --- --- 2

2. (STERIGMA) Herba stellato-tomentosa; siliqua elongata anguste linearis, stylus crassiusculus; *stigma divaricato-bilobum*: *Sterigma tomentosum* Willd.
Tomentum stellipilum nullum; stigma indivisum vel (in Chorispora) erecte subbilobum --- --- --- --- --- --- --- --- --- --- 3

3. Pedicelli (saltem pro parte) foliaceobracteati; siliquæ angustæ linear-i-elongatæ transverse 2-articulatæ, articulis multilocellatis --- --- --- --- --- --- --- --- --- --- 4

Pedicelli ebracteati --- --- --- --- --- --- --- --- --- --- 5

4. (ENARTHROCARPUS) Pedicelli fere omnes bracteati; siliquæ articulus *inferior elongatus*; flores parvi *Enarthrocarpus lyratus* Forsk.
Pedicelli inferiores solum bracteati; siliquæ articulus *inferior brevissimus* v. *subnullus*; flores majusculi --- --- *E. arcuatus* Labill.

5. Siliquæ angustæ, tenuiter elongatae, pedicellis (saltem inferne) haud vel vix crassiores) --- --- --- --- --- --- --- --- --- 6

Siliquæ abbreviatæ --- --- --- --- --- --- --- --- --- 7

6. (CHORISPORA) Siliquæ (inarticulatæ, incurvæ) apicem versus subulato-attenuatæ (flores lilacini, folia superiora indivisa): *Chorispora tenella* DC.
(ERUCARIA) Siliquæ articulatæ pars inferior cylindricus, superior (brevior) oblongo-cylindricus (flores albi, folia linear-i-pinnatisceta) --- --- --- --- --- --- --- --- --- *Erucaria aleppica* Gærtn.

7. Siliquæ inarticulatæ	---	---	---	---	---	---	---	8
Siliquæ articulatæ	---	---	---	---	---	---	---	18
8. Siliquæ abrupte pedicellatæ i. e. pedicellis fructiferis elongatis gracilibus multo latiores	---	---	---	---	---	---	---	9
Siliquæ basi pedicellis fructiferis abbreviatis incrassatisve æquilate vel sessiles	---	---	---	---	---	---	---	17
9. (NESLIA) Siliquæ exacte globulosæ filiformi-stylatae (folia caulinæ indivisa sagittato-sessilia, pilis ramosis vestita):								
								<i>Neslia paniculata Desv.</i>
Siliquæ oblique ovales vel longiores	---	---	---	---	---	---	---	10
10. Herbæ glaberrimæ foliis caulinis sagittato-vel cordato-sessilibus; flores minuti	---	---	---	---	---	---	---	11
Herbæ \pm pilis exasperatæ; flores majusculi	---	---	---	---	---	---	---	12
11. (CALEPINA) Siliqua (parva) ovoidea; stigma sessile; pedicelli fructiferi erecto-patuli	---	---	---	---	---	---		<i>Calepina Corvini Desv.</i>
(GOLDBACHIA) Siliqua (majuscula) curva tetragona in stylum compressum attenuata; pedicelli fructiferi recurvi:								
								<i>Goldbachia lœvigata DC.</i>
12. (BUNIAS) Siliqua oblique ovoideo-acutata i. e. in stylum breviter conicum attenuata	---	---	---	---	---	---		<i>Bunias orientalis L.</i>
Siliqua quadrialata vel cylindracea subulato-stylata	---	---	---	---	---	---	---	13
13. (ERUCAGO) Siliqua erosio-quadrialata medio strangulata:								<i>Erucago runcinata Horn.</i>
Siliqua teretiuscula	---	---	---	---	---	---	---	14
14. (RAPHANUS) Siliquæ spongioso-inflatæ oblongo-conicæ vix toru- losæ indehiscentes	---	---	---	---	---	---		<i>Raphanus sativus L.</i>
Siliquæ moniliformes articulatim secedentes	---	---	---	---	---	---	---	15
15. (RAPHANISTRUM) Stylus siliquæ internodio supremo vix lon- gior (foliorum segmenta approximatissima alternatim valde dis. æqualia)	---	---	---	---	---	---		<i>Raphanistrum maritimum Rchb.</i>
Stylus magis elongatus (foliorum segmenta inter se \pm remota)	---	---	---	---	---	---	---	16
16. Stylus siliquæ articulo supremo 3—4-plo longior; (foliorum lyra- torum laciniae regulariter oppositæ)	---	---	---	---	---	---		<i>R. innocuum Med.</i>
Stylus articulo proximo 2-plo longior; folia interrupte lyrata i. e. laciniae alternantes	---	---	---	---	---	---	---	<i>R. Landra Rchb.</i>
17. (EUCLIDIUM) Siliqua oblique ovato-globosa arcuatim recurvo- rostrata (herba pilis ramosis aspera, foliis basi attenuatis):								<i>Euclidium syriacum R. Br.</i>
(MYAGRUM) Siliqua incudiformis i. e. anceps e basi angusta tringulari-dilatata, apice truncata breviter conoideo-stylata, (herba glaberrima foliis auriculato-sessilibus)	---	---	---	---	---	---	---	<i>Myagrum perfoliatum L.</i>

18. Siliquæ articuli ambeduo subæquilongi, plerumque crassi, ampli- tudine vix conspicue difformes ; stylus conicus vel subulatus	19
Siliquæ articulus inferior brevissimus, superiore multo brevior, demum fere oblitteratus, pedicelliformis ; stigma sessile	26
19. (CAKILE) Siliquæ articulus superior ensiformis, compressus, infe- rior obconicus ; tota siliqua obconico-pyramidalo-quadrangularis :	
<i>Cakile maritima Scop.</i>	
Siliquæ articulus superior ovoideus vel subglobosus	20
20. (GUIRAOA) Siliquæ articulus superior (inæqualiter) 8-alato-cos- tatus	
<i>Guiraoa arvensis Coss.</i>	
Siliquæ haud alari-costatae	21
21. Flores flavi (semen in articulo superiore erectum)	22
Flores albi (semina omnia pendula ; stylus subulatus)	25
22. (RAPISTRUM) Stylus brevissimus	23
Stylus articulo superiore æquilongus vel longior	24
23. Pedicelli fructiferi (incrassati) articulo inferiore cylindrico te- nuiores	
<i>Rapistrum orientale DC.</i>	
Pedicelli fructiferi articulo inferiore oblongo angustiores :	
<i>R. perenne Berg.</i>	
24. Articulus siliquæ inferior obconicus v. oblongus ; pedicelli crassi articulo inferiore æquilongi vel longiores ;	
<i>R. rugosum Berg.</i>	
Articulus inferior cylindricus ; pedicelli filiformes articulo infe- riore (vix tenuiore) 2—3-plo longiores	
<i>R. Linnæanum B. et R.</i>	
25. (DIDESMUS) Adpressissime aspero-puberulus ; folia indivisa vel lyrata	
<i>Didesmus ægyptius L.</i>	
Hirtus ; folia in lacinias lineares breves pinnatisecta :	
<i>D. tenuifolius Sibth.</i>	
26. (CRAMBE) Siliquæ crasse ovoideæ ; folia indivisa undulata, si- nuata vel lobata ; tota herba crassa glaberrima	
<i>Crambe maritima L.</i>	
Siliquæ exakte globosæ	27
27. Folia basilaria simpliciter pinnatipartita ; tota glabra :	
<i>Cr. pinnatifida R. Br.</i>	
Folia basilaria diversimodo nunc palmatum inordinate multi- partita, supradecomposita, nunc lyrato-pinnatisecta ; herbæ ± hispidae vel strigosaæ	28
28. Folia inordinate multipartita supradecomposita ; herbæ robustæ, ramis crassis	
<i>Cr. pinnatifida R. Br.</i>	
Folia lyrato-pinnatisecta ; herbæ graciles, caule ramisque virgatis	29
29. Siliquæ (pisi magnitudine) præter costas quatuor longitudinales	
30	

tota superficie nervis secundariis tertiarisque elevatis creber-
rime arguteve reticulato-rugulosæ — — — — — *Cr. Tataria Sebeck.*
(Cr. aspera MaB.)

Siliquæ (grani piperis magnitudine!) præter costas longitudinaliter quadrantes nervis primariis elevatis obscure scrobiculatæ, lævigatae — Cr. Biebersteinii Janka.
 (Cr. Tataria MaB. et autor. transsilv.—
 Cr. aspera Janka pl. exs. a. 1876,

30. Pedicelli fructu breviores; foliorum segmenta lateralia 4—6:

Cr. reniformis Desf.

Pedicelli siliqua 2-3-plo longiores; foliorum segmenta lateralia solum 2 parva ... Cr. hispanica L.