

LIGNINOLYTIC ENZYME SYSTEM IN ECOLOGICAL ADAPTATION OF LIGNICOLOUS MACROFUNGI

BALAES, T.^{1*} – PETRE, C. V.¹ – UNGUREANU, C.² – MARDARI, C.¹ – TĂNASE, C.³

¹*Anastasiu Fatu Botanical Garden, Alexandru Ioan Cuza University of Iasi
Dumbrava Rosie 7-9, 700487, Iasi, Romania*

²*Dunarea de Jos University, Faculty of Food Science and Engineering
47 Domneasca St., 800008 Galati, Romania*

³*Faculty of Biology, Alexandru Ioan Cuza University of Iasi
Carol I 20A, 700505, Iasi, Romania*

**Corresponding author*

e-mail: tiberius.balaes@gmail.com; phone: +40749789156

(Received 13th Aug 2016; accepted 10th Oct 2016)

Abstract. Lignicolous macrofungi are the most important group of wood decomposers. Among the enzymes involved in wood decomposition, ligninases play an important role in this process and species that produce those enzymes degrade both cellulose and lignin. In this study we hypothesized that ligninases are influencing the ecological success of lignicolous macrofungi. Three hypotheses have been tested: h1 – the species producing several types of ligninases have a broader spectrum of hosts / substrates; h2 – the obligate saprotrophic species have a greater potential in degrading lignin and related compounds than the other groups; and h3 – the frequencies of lignicolous species is related to the production of highly active enzymes. Scientific data compiled from literature and completed with our own experimental results have been statistically interpreted using XLStat and MaxLite Software. Our results confirm the first two hypotheses, proving that ligninases play a direct role in colonizing a wide range of wood types, with chemical particularities. The third hypothesis should be rejected as no correlation has been observed. The present study offers new insights into ecological meanings of ligninases, and is the first attempt to connect the ligninolytic enzyme system to host range.

Keywords: *ligninases, wood decomposers, lignin degradation, white-rot fungi, laccases*

Introduction

Lignicolous macrofungi form a large group of fungi involved in the degradation of wood, with remarkable adaptations to different ecological niches.

Saprotrophic, parasitic or sapro-parasitic lignicolous macrofungi possess different strategies for colonizing wood and using it as a substrate and a source of nutrients. Depending on their ecology and type of nutrition, these species decay the wood, decomposing cellulose, hemicelluloses and sometimes lignin, which is one of the most abundant biomacromolecule on Earth (Knežević et al., 2013). Degradation of lignin might offer fungi advantages such as eliminating the barrier for the degradation of cellulose and increasing the availability of nitrogen in a substrate with a very high C : N ratio, through degradation of other wood constituents (Deacon, 2006).

This study is aimed at testing if the number of ligninases produced is influencing the spectrum of substrate / wood type / hosts, thus having ecological meanings. For this purpose three hypothesis have been formulated: h1 – the species producing several types of ligninases are able to colonize more types of wood compared to the species that produce a smaller number of ligninases; h2 – the obligate saprotrophic species (OS)

have a greater potential in degrading wood (by degrading lignin) than sapro-parasitic (SP) and obligate parasitic species (OP); and h3 – competitive species produce highly active enzymes. To test these hypothesis data concerning the types of substrata, ligninase production and enzyme activity, for 69 lignicolous basidiomycetes, have been collected from literature and completed with personal experimental results.

Our interpretation suggests that there is a positive correlation between the number of ligninases and the number of hosts and OS are better wood decomposers, while the third hypothesis should be rejected. These findings offer new insights over the biochemical adaptation for colonizing a substratum with particular ecological characteristics such as wood. From our knowledge, this is the first study trying to search possible directly connections between the ligninolytic enzyme system and host range of lignicolous macrofungi.

Review of Literature

Wood decay takes place through different biochemical mechanisms, and three types of rot can occur: a – soft rot, when fungi produce cellulases and H_2O_2 ; b – brown rot, when a cellulolytic enzyme system is involved and the cellulose / hemicelluloses are degraded while the lignin persists in the wood giving it a characteristic brown color; and c- white rot, in this case cellulose / hemicelluloses and lignin are degraded by particular enzymes. The concept of classifying wood rot is currently in revision, as some species “mimic” the white rot production (Nagy et al., 2015).

Ligninases are enzymes that produce the breakdown of lignin during wood decay. These enzymes play an important role in the nutrition of lignicolous fungi, frequently associated and acting synergistically.

There are several types of ligninases, and a particular species of macrofungi might produce one, few or all types. Many species produce two or three isomorphs of the same enzyme. The most common type of ligninases is represented by laccases, described for a large number of lignicolous macrofungi, including the ones that produce brown rot although the synthesis of laccases have been recorded in these cases as traces.

The occurrence of laccases (EC 1.10.3.2) and their role in the ecology of lignicolous fungi have been reviewed by several authors (Baldrian, 2006; Valderrama et al., 2003). Other common ligninases are Lignin Peroxidase-LiP (E.C. 1.11.1.14), Manganese dependent Peroxidase-MnP (E.C. 1.11.1.13), Manganese Independent Peroxidase-MiP (EC 1.11.1.16), while less frequent are Aryl Alcohol Oxidase-AAO, Versatile Peroxidase-VP, Dye-degrading Peroxidase-DyP (Anastasi et al., 2010; Graž and Jarosz-Wilkolazka, 2011; Palmieri et al., 2005). In an extensive study involving the screening of genes encoding ligninases in a broad sense, Nagy and collaborators (2015) suggest that in the degradation of lignin, a much larger group of ligninases is involved, although some classes of enzymes play a secondary role. The production of a particular ligninase vary largely from one species to another in terms of amounts and the enzyme activity and the required conditions as well.

Materials and Methods

The available literature has been reviewed in order to collect information regarding the host's spectrum, number of ligninases produced, frequency of species, type of nutrition and the efficiency of ligninolytic enzyme system for 69 species of lignicolous basidiomycetes. Most of the species listed in *Table 1* have also been isolated and stored

in our laboratory collection. When data were missing from literature for particular species, the information has been completed with experimental data.

Table 1. List of investigated species of macrofungi, their enzymatic properties, ecological features and data source (NE – number of enzymes, EA – enzyme activity, DE – dye degradation efficiency, CH – common host genera – TH, F – frequency, NT – nutrition type)

| Species | NE | EA | DE | CH | TH | F | NT | References |
|-----------------------------------|----|----|----|----|----|---|----|--|
| <i>Abortiporus biennis</i> | 3 | 3 | 4 | 18 | 26 | 1 | OS | Aggelis et al., 2002; ARS; Bernicchia, 2005; Casieri et al., 2010; Ryvarden and Gilbertson, 1993 |
| <i>Armillaria mellea</i> | 3 | 3 | 1 | 9 | 82 | 2 | OP | Balaş et al., 2013; Casieri et al., 2010; Diamandis and Perlerou, 2001; Matsushita and Suzuki, 2005; Stoytchev and Nerud, 2000; Otieno et al., 2003; PKB; Qin et al., 2007; Szczepkowski, 2007 |
| <i>Auricularia auricula-judae</i> | 4 | 4 | 1 | 2 | 24 | 2 | OS | ARS; Balaş et al., 2013; Liers et al., 2010; Negrean and Anastasiu, 2004 |
| <i>Bjerkandera adusta</i> | 6 | 7 | 5 | 25 | 32 | 2 | OS | Anastasi et al., 2010; Balaş et al., 2013; Balaş and Tănase, 2013; Bernicchia, 2005; Eichlerova et al., 2006a; Erkkilä and Niemelä, 1986; Guillén et al., 2011; Robinson and Nigam, 2008; Ryvarden and Gilbertson, 1993; Tinoco et al., 2007 |
| <i>Bjerkandera fumosa</i> | 1 | 1 | 3 | 15 | 20 | 1 | OS | ARS; Balaş et al., 2013; Bernicchia, 2005; Graz and Jarosz-Wilkolazka 2011; Ryvarden and Gilbertson, 1993 |
| <i>Cerrena unicolor</i> | 4 | 4 | 4 | 17 | 21 | 2 | OS | Bernicchia, 2005; Elisashvili et al., 2010; Erkkilä and Niemelä, 1986; Hattori, 2005; Ranadive et al., 2011; Souza-Ticlo et al., 2009 |
| <i>Coriolopsis gallica</i> | 2 | 2 | 4 | 2 | 20 | 2 | SP | ARS; Balaş et al., 2013; Bernicchia, 2005; Hansent and Knudsen, 1997; Robinson et al., 2001; Ryvarden and Gilbertson, 1993 |
| <i>Cyathus striatus</i> | 1 | 1 | 2 | 1 | 3 | 2 | OS | ARS; Balaş et al., 2013; Casieri et al., 2010 |
| <i>Daedalea quercina</i> | 2 | 2 | 1 | 1 | 15 | 2 | OS | Balaş et al., 2013; Baldrian 2004; Bernicchia, 2005; Ranadive et al., 2011; Ryvarden and Gilbertson, 1993 |
| <i>Daedaleopsis confragosa</i> | 1 | 1 | 1 | 1 | 19 | 2 | SP | Balaş et al., 2013; Bernicchia, 2005; Erkkilä and Niemelä, 1986; Orth et al., 1993; Ranadive et al., 2011; Ryvarden and Gilbertson, 1993 |
| <i>Datronia caperata</i> | 1 | 1 | 3 | 1 | 3 | 1 | SP | Abrahao et al., 2008; ARS; Gilbert and Sousa, 2002; Minter et al., 2001 |
| <i>Dichomitus squalens</i> | 6 | 5 | 5 | 2 | 7 | 1 | OS | Aggelis et al., 2002; ARS; Bernicchia, 2005; Eichlerova et al., 2006c; Novotný et al., 2012; Orth et al., 1993; Ryvarden and Gilbertson, 1993; Šušla et al., 2007 |
| <i>Flammulina velutipes</i> | 1 | 1 | 2 | 2 | 16 | 2 | OS | ARS; Balaş et al., 2013; Eichlerova et al., 2006a; Gerault, 2005; Petersen et al., 1999; Szczepkowski, 2007 |

| | | | | | | | | |
|--------------------------------|---|---|---|----|----|---|----|--|
| <i>Fomes fomentarius</i> | 2 | 2 | 2 | 3 | 18 | 2 | SP | Balaeş et al., 2013; Bernicchia, 2005; Casieri et al., 2010; Elisashvili et al., 2009; Erkkilä and Niemelä, 1986; Ryvarden and Gilbertson, 1993 |
| <i>Fomitopsis pinicola</i> | 1 | 1 | 1 | 5 | 23 | 2 | SP | Balaeş et al., 2013; Casieri et al., 2010; Erkkilä and Niemelä, 1986; Hattori, 2005; Ryvarden and Gilbertson, 1993 |
| <i>Ganoderma adspersum</i> | 1 | 1 | 3 | 1 | 9 | 2 | OP | ARS; Balaeş et al., 2013; De Simone and Annesi, 2012; Elisashvili and Kachlishvili 2009 |
| <i>Ganoderma applanatum</i> | 2 | 2 | 2 | 9 | 53 | 2 | SP | Balaeş et al., 2013; Bernicchia, 2005; Casieri et al., 2010; Elisashvili and Kachlishvili 2009; Elisashvili et al., 2009; Erkkilä and Niemelä, 1986; Levin et al., 2004; Ryvarden and Gilbertson, 1993; Sankaran et al., 2005; SIPMP; Szczepkowski, 2007 |
| <i>Ganoderma lucidum</i> | 4 | 4 | 2 | 13 | 45 | 2 | OS | Asgher et al., 2010; Balaeş et al., 2013; Bernicchia, 2005; Elisashvili and Kachlishvili 2009; Erkkilä and Niemelä, 1986; Orth et al., 1993; Ranadive et al., 2011; Ryvarden and Gilbertson, 1993; Sankaran et al., 2005; SIPMP |
| <i>Ganoderma resinaceum</i> | 1 | 1 | 3 | 1 | 16 | 2 | SP | Balaeş et al., 2013; Bernicchia, 2005; Casieri et al., 2010; Ranadive et al., 2011; Ryvarden and Gilbertson, 1993 |
| <i>Gloeophyllum odoratum</i> | 3 | 3 | 3 | 1 | 4 | 2 | SP | Anastasi et al., 2010; Casieri et al., 2010; Bernicchia, 2005; Erkkilä and Niemelä, 1986; Ryvarden and Gilbertson, 1993 |
| <i>Grifola frondosa*</i> | 2 | 1 | 1 | 1 | 7 | 1 | SP | Bernicchia, 2005; Hansent and Knudsen, 1997; Orth et al., 1993; Ryvarden and Gilbertson, 1993 |
| <i>Gymnopilus junonius</i> | 2 | 2 | 2 | 4 | 14 | 1 | OS | ARS; Balaeş et al., 2013; Rees and Strid, 2001; Valentin et al., 2009 |
| <i>Hemipholiota populnea</i> | 0 | 0 | 1 | 1 | 1 | 1 | OP | ARS; Balaeş et al., 2013; Szczepkowski, 2007 |
| <i>Hymenopelis radicata*</i> | 1 | 1 | 2 | 1 | 4 | 2 | OS | ARS; Balaraju et al., 2010 |
| <i>Hypholoma fasciculare</i> | 2 | 2 | 2 | 2 | 7 | 2 | OS | Abrahao et al., 2008; Angelini et al., 2012; Balaeş et al., 2013; Casieri et al., 2010; Gramss et al., 1999 |
| <i>Inonotus hispidus</i> | 3 | 3 | 1 | 3 | 23 | 2 | OP | Aggelis et al., 2002; ARS; Balaeş et al., 2013; Bernicchia, 2005; Casieri et al., 2010; Gerault, 2006; Nerud and Mišurcova 1996; Ryvarden and Gilbertson, 1993 |
| <i>Irpex lacteus</i> | 6 | 6 | 5 | 14 | 20 | 2 | OS | Aggelis et al., 2002; Balaeş et al., 2013; Bernicchia and Gorjón, 2010; Bernicchia, 2005; Hattori, 2005; Levin et al., 2004; Novotný et al., 2009; Ryvarden and Gilbertson, 1993 |
| <i>Kuehneromyces mutabilis</i> | 1 | 1 | 1 | 1 | 3 | 2 | OS | Balaeş et al., 2013; Hansent and Knudsen, 1992; Steffen et al., 2007 |
| <i>Laetiporus sulphureus*</i> | 1 | 1 | 2 | 3 | 23 | 2 | SP | ARS; Bernicchia, 2005; Casieri et al., 2010; Erkkilä and Niemelä, 1986; Hansent and Knudsen, 1997; Hattori, 2005; Ryvarden and Gilbertson, 1993 |

| | | | | | | | | |
|-----------------------------------|---|---|---|----|----|---|----|--|
| <i>Lentinula edodes</i> | 3 | 3 | 4 | 3 | 12 | 2 | OS | ARS; Bisen et al., 2010; Boer et al., 2004; Kalmış et al., 2008; Orth et al., 1993 |
| <i>Lentinus neostrigosus</i> * | 2 | 2 | 3 | 3 | 20 | 2 | OS | ARS; Hansent and Knudsen, 1992; Vaithanomsat et al., 2012 |
| <i>Lentinus tigrinus</i> * | 3 | 3 | 4 | 3 | 12 | 2 | OS | Aggelis et al., 2002; ARS; Hansent and Knudsen, 1992; Moreira et al., 2000 |
| <i>Lenzites betulina</i> | 5 | 5 | 5 | 8 | 21 | 2 | OS | Anastasi et al., 2010; Balaes et al., 2013; Balaes et al., 2014; Bernicchia, 2005; Erkkilä and Niemelä, 1986; Guillén et al., 2011; Hansent and Knudsen, 1997; Hattori, 2005; Moturi et al., 2009; Ryvarden and Gilbertson, 1993 |
| <i>Lycoperdon pyriforme</i> | 1 | 1 | 1 | 1 | 10 | 2 | OS | Angelini et al., 2012; ARS; Balaes et al., 2013; Casieri et al., 2010; Pegler et al., 1995 |
| <i>Megacollybia platyphylla</i> * | 1 | 1 | 1 | 1 | 2 | 2 | OS | ARS; Casieri et al., 2010 |
| <i>Meripilus giganteus</i> * | 2 | 2 | 1 | 2 | 24 | 2 | SP | ARS; Bernicchia, 2005; Kalmış et al., 2008; Ryvarden and Gilbertson, 1994 |
| <i>Merulius tremellosus</i> | 2 | 2 | 1 | 1 | 18 | 2 | OS | ARS; Balaes et al., 2013; Bernicchia and Gorjón, 2010; Kum et al., 2011; Szczepkowski, 2007 |
| <i>Oudemansiella mucida</i> * | 2 | 1 | 1 | 2 | 5 | 2 | OS | ARS; Daniel et al., 1994; Gerault, 2005 |
| <i>Panellus stypticus</i> * | 3 | 3 | 3 | 4 | 17 | 2 | OS | Aggelis et al., 2002; ARS; Bernicchia and Gorjón, 2010; Casieri et al., 2010; Hansent and Knudsen, 1997; Nerud and Mišurcova 1996 |
| <i>Peniophora quercina</i> * | 1 | 1 | 2 | 2 | 14 | 2 | OS | ARS; Balaes et al., 2013; Bernicchia, 2005; Erkkilä and Niemelä, 1986; Hansent and Knudsen, 1997; Ryvarden and Gilbertson, 1994 |
| <i>Phellinus igniarius</i> | 1 | 1 | 2 | 3 | 39 | 2 | SP | ARS; Balaes et al., 2013; Bernicchia, 2005; Casieri et al., 2010; Erkkilä and Niemelä, 1986; Hansent and Knudsen, 1997; Ryvarden and Gilbertson, 1994 |
| <i>Phellinus pomaceus</i> | 1 | 1 | 2 | 1 | 24 | 2 | OP | ARS; Balaes et al., 2013; Bernicchia, 2005; Casieri et al., 2010; Ranadive et al., 2011; Ryvarden and Gilbertson, 1994 |
| <i>Phellinus torulosus</i> | 1 | 1 | 4 | 28 | 64 | 2 | OP | Casieri et al., 2010; Nakasone and Burdsall, 1995 |
| <i>Phlebia floridensis</i> * | 3 | 3 | 4 | 2 | 6 | 2 | OS | Arora and Gill, 2005; ARS; Bernicchia and Gorjón, 2010; Szczepkowski, 2007 |
| <i>Phlebia radiata</i> * | 6 | 7 | 5 | 2 | 15 | 2 | OS | Anastasi et al., 2010; ARS; Casieri et al., 2010; Hilden et al., 2005; Mäkelä et al., 2006; Smith and Hesler, 1968 |
| <i>Pholiota aurivella</i> | 2 | 2 | 1 | 3 | 13 | 2 | OP | Balaes et al., 2013; Eichlerova et al., 2006a |
| <i>Piptoporus betulinus</i> * | 1 | 1 | 1 | 1 | 1 | 2 | SP | Bernicchia, 2005; Casieri et al., 2010; Erkkilä and Niemelä, 1986; Gerault, 2006; Hansent and Knudsen, 1997; Ryvarden and Gilbertson, 1994 |
| <i>Pleurotus florida</i> | 1 | 1 | 1 | 1 | 1 | 2 | OS | ARS; Gbolagade et al., 2006; Pant and Adholeya, 2009 |
| <i>Pleurotus dryinus</i> | 2 | 2 | 2 | 10 | 18 | 1 | OS | ARS; Eichlerova et al., 2006b; EOL |
| <i>Pleurotus</i> | 7 | 7 | 2 | 11 | 43 | 2 | OS | Aggelis et al., 2002; Anastasi et al., 2010; |

| | | | | | | | | |
|--------------------------------|---|---|---|----|----|---|----|--|
| <i>ostreatus</i> | | | | | | | | Balaes et al., 2013; Casieri et al., 2008; Eichlerova et al., 2006a; Hansent and Knudsen, 1997; Palmieri et al., 2005; Pozdnyakova et al., 2010 |
| <i>Pleurotus pulmonarius</i> | 3 | 3 | 3 | 5 | 7 | 2 | OS | ARS; Bernicchia, 2005; Eichlerova et al., 2006c; Hattori, 2005; Orth et al., 1993; Rigas et Dritsa, 2006; Ryvarden and Gilbertson, 1994 |
| <i>Polyporus alveolaris*</i> | 2 | 2 | 5 | 9 | 20 | 1 | OS | Barassa et al., 2009; Bernicchia, 2005; Erkkilâ and Niemelâ, 1986; Lee et al., 2010; Ryvarden and Gilbertson, 1994 |
| <i>Polyporus brumalis</i> | 3 | 3 | 2 | 22 | 27 | 2 | OS | ARS; Bernicchia, 2005; Erkkilâ and Niemelâ, 1986; Kum et al., 2011; Rigas si Dritsa, 2006; Ryu et al., 2008; Ryvarden and Gilbertson, 1994 |
| <i>Polyporus squamosus</i> | 2 | 2 | 1 | 25 | 28 | 2 | SP | ARS; Casieri et al., 2010; Eichlerova et al., 2006a |
| <i>Postia caesia</i> | 1 | 1 | 1 | 7 | 30 | 2 | OS | ARS; Balaes et al., 2013; Eichlerova et al., 2006a; Gerault, 2006 |
| <i>Postia stiptica</i> | 1 | 1 | 1 | 3 | 22 | 2 | OS | ARS; Balaes et al., 2013; Bernicchia, 2005; Casieri et al., 2010; Hansent and Knudsen, 1997; Ryvarden and Gilbertson, 1994 |
| <i>Pycnoporus cinnabarinus</i> | 2 | 2 | 4 | 12 | 28 | 2 | OS | ARS; Bernicchia, 2005; Casieri et al., 2010; Fedrova et al., 2013; Hansent and Knudsen, 1997; Orth et al., 1993; Ryvarden and Gilbertson, 1994 |
| <i>Trametes gibbosa</i> | 3 | 3 | 5 | 24 | 34 | 2 | OS | ARS; Balaes et al., 2013; Balaes et al., 2014; Bernicchia, 2005; Elisashvili and Kachlishvili 2009; Elisashvili et al., 2009; Fedrova et al., 2013; Hansent and Knudsen, 1997; Orth et al., 1993; Ryvarden and Gilbertson, 1994; Szczepkowski, 2007 |
| <i>Trametes hirsuta</i> | 5 | 5 | 5 | 38 | 52 | 2 | OS | Aggelis et al., 2002; Balaes et al., 2013; Bernicchia, 2005; Casieri et al., 2010; Erkkilâ and Niemelâ, 1986; Haibo et al., 2009; Hansent and Knudsen, 1997; Hattori, 2005; Orth et al., 1993; Ryvarden and Gilbertson, 1994; Szczepkowski, 2007; Tomšovský and Homolka 2003 |
| <i>Trametes ochracea</i> | 3 | 3 | 5 | 19 | 23 | 2 | OS | ARS; Balaes et al., 2013; Bernicchia, 2005; Casieri et al., 2010; Elisashvili and Kachlishvili 2009; Gerault, 2006; Ryvarden and Gilbertson, 1994; Tomšovský and Homolka 2003 |
| <i>Trametes pubescens</i> | 6 | 7 | 5 | 15 | 20 | 2 | OS | Anastasi et al., 2010; ARS; Balaes et al., 2013; Bernicchia, 2005; Casieri et al., 2008, 2010; Erkkilâ and Niemelâ, 1986; Nikitina et al., 2005; Ryvarden and Gilbertson, 1994 |
| <i>Trametes suaveolens</i> | 2 | 2 | 4 | 2 | 9 | 2 | SP | ARS; Balaes et al., 2013; Bernicchia, 2005; Erkkilâ and Niemelâ, 1986; Hansent and Knudsen, 1997; Knežević et al., 2013; Ryvarden and Gilbertson, 1994 |
| <i>Trametes trogii*</i> | 6 | 6 | 5 | 2 | 9 | 2 | OS | ARS; Bernicchia, 2005; Dhoub et al., 2005; Levin et al., 2002, 2004; Ryvarden |

| | | | | | | | | |
|------------------------------|---|---|---|----|----|---|----|---|
| | | | | | | | | and Gilbertson, 1993 |
| <i>Trametes versicolor</i> | 8 | 9 | 4 | 40 | 61 | 2 | OS | Anastasi et al., 2010; Balaes et al., 2013; Bernicchia, 2005; Casieri et al., 2010; Erkkilä and Niemelä, 1986; Guillén et al., 2011; Hattori, 2005; Levin et al., 2004; Liu et al., 2004; Lorenzo et al., 2006; Moturi et al., 2009; Orth et al., 1993; Ryvarden and Gilbertson, 1994; Tomšovský and Homolka 2003; Ungureanu et al., 2015 |
| <i>Trichaptum abietinum</i> | 2 | 2 | 2 | 5 | 22 | 2 | OS | ARS; Bernicchia, 2005; Eichlerova et al., 2006a; Erkkilä and Niemelä, 1986; Ryvarden and Gilbertson, 1994 |
| <i>Trichaptum bifforme*</i> | 2 | 2 | 2 | 7 | 18 | 2 | OS | ARS; Bernicchia, 2005; Elisashvili et al., 2009; Hansent and Knudsen, 1997 |
| <i>Schizophyllum commune</i> | 4 | 3 | 2 | 6 | 42 | 2 | SP | ARS; Balaes et al., 2013 ; Bhatti et al., 2008; Hansent and Knudsen, 1997; Levin et al., 2004; Li et al., 2009; Szczepkowski, 2007 |
| <i>Stereum hirsutum</i> | 1 | 1 | 1 | 2 | 11 | 2 | OS | Balaes et al., 2013; Guillén et al., 2011; Orth et al., 1993; Szczepkowski, 2007 |
| <i>Xylobolus frustulatus</i> | 1 | 1 | 1 | 1 | 3 | 1 | OS | Balaes et al., 2013 ; Cookson, 1995; Hansent and Knudsen, 1997; Ranadive et al., 2011 |

*experimental results from the present research are included

Analytical procedures

For assessing the efficiency of dye degradation, the protocol described in Balaes and collaborators (2013) has been used. Pure fungal cultures were grown on solid and liquid media and used as sources of inoculum. The presence / absence of laccase, and the enzyme activity has been highlighted through a modified protocol (Guo et al., 2011), using liquid medium supplemented with wheat bran (per L⁻¹: 6 g glucose, 3 g (NH₄)₂SO₄, 6 g wheat bran, 0.01 g CuSO₄), and inoculated with pelletized mycelium (10 mL inoculum L⁻¹). As a specific substrate, 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid (ABTS) has been used. The reaction mixture contained 0.25 mL 50mM ABTS, 0.5 mL 0.1M acetate buffer, 0.25 mL enzyme extract. Substrate oxidation was monitored by measuring the increase of absorbance at 420 nm (ϵ_{420} , 3.6•10⁴ M⁻¹ cm⁻¹), after 1 min at 25° C. A unit of laccase activity was defined as the amount of enzyme oxidizing 1 µmol of ABTS per minute in the considered analysis conditions (Bourbonnais et al., 1995).

Data processing

Compiled dataset included 6 variables (*Table 1*): number of ligninases, enzyme activity (expressed as classes from 1 – poor enzyme activity / traces, to 7 – very strong activity), dye degradation efficiency (expressed as classes from 1 – very weak degradation, few dyes or no degradation at all, to 5 – very efficient degradation of different dyes), number of common host genera, number of total host genera and frequency (expressed as classes, 1 – less frequent, 2 – frequent or very frequent). A seventh variable refers to the type of nutrition – obligate saprotrophic (OS), sapro-

parasitic (SP) and obligate parasitic species (OP). For particular tests, values from previous datasets were grouped into three categories corresponding to the nutrition type.

Statistical interpretations

For all statistical calculations and analysis XLStat (trial version) and MaxLite (freeware) softwares have been used. As data do not follow normal distribution, non-parametric tests were used. For assessing the correlations between different variables, Spearman correlation coefficients were used, at a confidence interval of 95%. Mann-Whitney test has been used to compare variables related to the three types of fungi according to their nutrition. A Principal Component Analysis was run for a better view of correlations between all variables, considering the nutrition type as a qualitative supplementary variable.

Results and Discussions

The present investigation is an attempt to elucidate the role of ligninolytic enzyme system in ecological adaptation of lignicolous macrofungi in terms of diversity of substrata (hosts colonization). For this purpose, three hypotheses have been formulated.

Hypothesis number 1: the species producing several types of ligninases are able to colonize many types of wood compared to the species that produce a smaller number of ligninases.

One first step of our investigation was to analyze the correlation between the number of ligninolytic enzymes and the number of hosts. In the table of correlation (Table 2) it can be seen that there is a positive correlation ($r=0.5529$, $p < 0.0001$) between the number of enzymes and the number of common host genera for all 69 species studied, while the correlation is weaker when considering the total host genera ($r=0.3738$, $p < 0.0001$). In the latter case there are also included species of trees / lignicolous substrate that are accidentally colonized by fungi and not as a normal behavior of a particular fungal species (e.g. *Stereum hirsutum* usually grows on deciduous trees, but it has been reported also on coniferous wood, although one may not expect to find *S. hirsutum* on this substrate in the field). Hence, using the number of common hosts for assessing the correlation between the number of ligninolytic enzymes and host diversity gives a better understanding of how the ligninolytic enzymes system is influencing the ecology of these fungi.

Table 2. Correlations between ligninolytic enzyme system and number of hosts – Spearman correlation coefficients (r values in upper, and p values, two-tailed, in lower matrix; confidence interval 95%; with bold are marked the relevant results)

| | Number of enzymes | Enzyme activity | Dye degradation efficiency | Common host genera | Total host genera |
|----------------------------|-------------------|-----------------|----------------------------|--------------------|-------------------|
| Number of enzymes | | 0.9898 | 0.5712 | 0.5529 | 0.3738 |
| Enzyme activity | < 0.0001 | | 0.5902 | 0.5604 | 0.3883 |
| Dye degradation efficiency | < 0.0001 | < 0.0001 | | 0.4559 | 0.182 |
| Common host genera | < 0.0001 | < 0.0001 | < 0.0001 | | 0.759 |
| Total host genera | 0.0016 | 0.001 | 0.1344 | < 0.0001 | |

We have included in our tests one other variable: the general enzyme activity, expressed as classes of efficiency (data collected from literature and completed with our own results). These data have the disadvantage of being organized as classes: the quantitative interpretation is thus affected by subjectivity to some degree. Raw quantitative data could not be used due to very different experimental conditions of various researchers. Including of this variable had the role of verifying if there is still a positive correlation between enzymes and hosts number. The correlation in this case was slightly stronger ($r=0.5604$, $p < 0.0001$), re-confirming h1.

A second attempt to verify this hypothesis consisted in using a new variable, derived from the first one – the efficiency of dye degradation as a measure of enzyme versatility. Ligninolytic enzymes are very versatile and attack a large spectrum of chemical compounds with structure similar to lignin. This property confers the ability to degrade different human made aromatic compounds, hydrocarbons, dyes with heterocycles (Balaes et al., 2014), while in natural conditions these fungi degrade lignin and also other chemical compounds from wood (Deacon, 2006). Considering these facts, we hypothesize that the versatility of ligninolytic enzymes could also be a factor that influence the ecological adaptation in terms of colonizing very different types of wood (varied chemical composition). However, we found weaker correlation compared to previous variables ($r=0.4559$, $p < 0.0001$).

The Principal Component Analysis (PCA) is an explanatory statistical tool used for visualization of data and to a lesser extent for testing hypotheses. Running PCA lead to confirm the results from previous tests, offering, in the same time, a largely view over the connections between variable. The procedure is based on the projection of all variables (six + one in our case) onto two axes, considering the first two factors that are contributing to the variability. We have considered the first six variable as principal quantitative variables and the seventh one (*Nutrition type*) as a supplementary qualitative variable.

In *Fig. 1* we can observe on the correlation circle that the first two components explain 73% of the data variability, making the projection a trusting one. A very strong positive correlation is found between the *Number of enzymes* and *Enzyme activity*, as it would have been expected. A good positive correlation is seen between these two variables and *Common host genera* on a side and *Dye degradation efficiency* on the other side. The position of projection points at a particular distance from the circle prove that the projection should be considered with caution for these two variables.




Figure 1. Circle of correlation between factors and variables after Principal Component Analysis (numbers on axes represent the correlation coefficients)

As seen in *Table 3*, factor loadings for these variables have the highest values on F1 dimension and second values (close to the first ones) on F3 and F4 dimensions respectively. The *Total host genera* variable is better projected on the chart (*Table 3*) and it can be observed that it is not correlated with *Number of enzymes*. The highest values of factor loadings for *Frequency* are on the F2 and F3 dimensions and are almost equal, and therefore no explanations can be assumed based on the projection.

Table 3. Factor loadings for the six quantitative variables in Principal Component Analysis (with bold are marked the highest values for each variable)

| Variables\Factors | F1 | F2 | F3 | F4 | F5 | F6 |
|----------------------------|--------------|--------------|---------------|--------------|--------|--------|
| Number of enzymes | 0.900 | -0.199 | 0.263 | -0.273 | -0.015 | -0.085 |
| Common host genera | 0.815 | 0.206 | -0.443 | 0.067 | 0.303 | 0.000 |
| Frequency | 0.205 | 0.687 | 0.664 | 0.205 | 0.060 | 0.000 |
| Enzyme activity | 0.907 | -0.209 | 0.253 | -0.250 | -0.022 | 0.087 |
| Total host genera | 0.657 | 0.574 | -0.415 | -0.007 | -0.259 | -0.001 |
| Dye degradation efficiency | 0.685 | -0.463 | 0.046 | 0.554 | -0.080 | -0.003 |

Hypothesis number 2 – OS species have a greater potential in degrading wood (by degrading lignin) than SP and OP species.

OS macrofungi use products derived from wood decomposition as sources of nutrients: cellulose and hemicelluloses are primary sources of C and energy, while lignin is degraded especially for making cellulose and other constituents of wood more accessible (Deacon, 2006). Since OS are dependent on the dead wood constituents, it was assumed that these species tend to have a highly versatile and active enzymes system compared to SP and OP macrofungi, which are using additional or mainly other strategies for nutrition. In other words, the second hypothesis assumed that OS have more active enzyme and degrade a broader spectrum of synthetic dyes (different chemical structure), being more versatile.

When applying the Mann-Whitney test for two by two datasets (*Table 4*) we observed that there is a significant superiority of dye degradation efficiency of OS compared to SP (one tailed *p* value 0.047) and to OP (one tailed *p* value 0.048), while between SP and OP there are no significant differences for this property. Concerning enzyme activity, OS produce more highly active enzymes than SP (one tailed *p* 0.008) and OP (one tailed *p* 0.046). These results are confirming our hypothesis.

Table 4. Mann-Whitney test (right-tailed test) for comparing the enzyme activity and dye degradation efficiency for the three type of macrofungi (the Mann-Whitney's *U* is normalized and tested against the normal distribution)

| | Dye degradation efficiency | | | Enzyme activity | | |
|--------------------|----------------------------|----------|---------|-----------------|----------|---------|
| | OS-SP | OS-OP | SP-OP | OS-SP | OS-OP | SP-OP |
| U | 469.000 | 222.500 | 64.000 | 514.000 | 223.500 | 57.000 |
| U (expected value) | 368.000 | 161.000 | 56.000 | 368.000 | 161.000 | 56.000 |
| U (variance) | 3651.782 | 1369.084 | 200.316 | 3627.943 | 1381.644 | 193.233 |
| Z (observed value) | 1.671 | 1.662 | 0.565 | 2.424 | 1.681 | 0.072 |
| Z (critical value) | 1.645 | 1.645 | 1.645 | 1.645 | 1.645 | 1.645 |
| One-tailed p-value | 0.047 | 0.048 | 0.286 | 0.008 | 0.046 | 0.471 |
| Alpha | 0.05 | 0.05 | 0.05 | 0.05 | 0.05 | 0.05 |

Lignin is considered a barrier for wood decomposition, but woody species also produce tannins, alkaloids, resins and other types of chemical compounds with defensive and/or structural role. Lignin is formed from three different monomers: *p*-coumaryl alcohol, coniferyl alcohol and sinapyl alcohol, and their varied ratio give spatial different structures for each species. Broad leaves trees (dicotyledons) have lignine made predominantly from *p*-coumaryl, while coniferous trees contain lignin composed mainly from coniferyl alcohol. Ligninases are versatile enzymes, able to breakdown the lignin with its different structure from a species to another, as well as oxidize other compounds polyphenols and heterocyclic compounds (Elisashvili et al., 2010). For OS macrofungi ligninases play important roles in their nutrition.

The Principal Component Analysis also offered information for a better view of ecological features of the three groups of macrofungi separated based on their nutrition type. Fig. 2 shows the projection of each observation (species characteristics) on the chart, grouped according to the nutrition type (three colors for OS, SP and OP). There is a distinct projection of the three groups, with only OS sharing some features with other two groups and with a higher dispersal on the chart, meaning there is a higher variability inside group, in terms of ligninases synthesis and host spectrum. OP group is more homogenous, but three species have particular characteristics, with *Armillaria mellea* (AM) and *Phellinus torulosus* (PT) having a very large host spectrum and producing ligninases to some degree, while *Hemipholiota populnea* (HP) has a completely distinct projection with a reduced host range (only *Populus*) and does not produce known ligninases.




Figure 2. The projections of observations (species characteristics) on the first two factors axes (AM – *A. mellea*, BA – *B. adusta*, DC – *D. caperata*, DS – *D. squalens*, GF – *G. frondosa*, HP – *H. populnea*, PT – *P. torulosus*, TH – *Trametes hirsute*, XF – *Xylobolus frustulatus*; legend: OP – obligate parasitic, OS – obligate saprotrophic, SP – sapro-parasitic macrofungi)

Among the SP species, *Grifola frondosa* (GF) and *Datronia caperata* (DC) possess features distinct from the other species in the group, with reduced host range, less frequent and producing only laccase in small amounts. Four OS species have characteristics different from other species in the group: *Bjerkandera adusta* (BA) and *Trametes hirsuta* (TH) produce many enzymes with strong activity and have a large host spectrum, while *Xylobolus frustulatus* (XF) produces few enzymes and has a reduced host spectrum (only *Quercus*) and *Dichomitus squalens* (DS) with strong enzyme activity but reduced host spectrum.

The projection of the species based on the nutrition type offers an interesting view of their ecological behaviors. The higher variability for OS fungi can be explained through their very different strategies for survival and development, some species growing on wood not affected by decay, while other grow on rotten wood; some species manifest antagonism against competitors, while other expand their mycelium in soil and litter, being more opportunistic. In contrast, SP and OP appear to be more specialized on their ecological niche, probably their strategies being oriented toward “fighting” their hosts.

Hypothesis number 3 – competitive species (frequent species) produce highly active enzymes.

To test this assumption, we verified if there is a positive correlation between frequency and enzyme activity / dye degradation efficiency. As it can be observed in Table 5, there is no correlation, at least in the case of our limited data, and we might assume that other factors are being involved in the process (hosts number, host frequency, other ecological factors). The third hypothesis has been rejected.

Table 5. Correlations between frequencies and enzyme activity / dye degradation efficiency – Spearman correlation coefficients (*r* values in upper, and *p* values, two-tailed, in lower matrix; confidence interval 95%)

| | Enzyme activity | Dye degradation efficiency | Frequency |
|----------------------------|------------------------|-----------------------------------|------------------|
| Enzyme activity | | 0.5902 | 0.1650 |
| Dye degradation efficiency | < 0.0001 | | -0.0383 |
| Frequency | 0.1755 | 0.7544 | |

An attempt to include data for distribution of fungal species has been abandoned since the distribution is strongly dependent on their hosts and production of ligninases is irrelevant in this case. The limitations of our study consist in the reduced availability of data concerning ligninase production and activity for many species of macrofungi, forcing us to take into consideration only 69 species. Different macrofungi might produce a particular ligninase in different amounts or with a different activity, and a reasonable assumption is that the quantity and quality of ligninases also play roles in wood degradation efficiency. In these circumstances is difficult to compare the ligninolytic activity of certain species as different authors are using varied protocols and particular conditions to test the enzyme activity. However, using classes of efficiency, we overcome this limitation and tested the formulated hypotheses.

In summary, we have investigated the influence of ligninolytic enzyme system over ecological success of lignicolous fungi in terms of hosts' spectrum. We assumed that fungal species producing several types of ligninases are cosmopolitan and colonize a broader spectrum of tree hosts and we found positive correlation that sustain this

hypothesis ($r=0.5529$, $p < 0.0001$). Our analysis suggests that obligate saprotrophic fungi tend to degrade several types of wood and more efficiently due to higher enzyme activity and versatility. It appears that this group of macrofungi is more heterogeneous in terms of ligninases production and host range, probably due to other ecological factors involved in their adaptation strategies than in the case of parasitic species.

It was assumed that fungal species with a larger host spectrum produce highly active and versatile ligninases, but this hypothesis has been rejected.

Future studies might contribute further to the understanding of the ecological importance of ligninases.

Aknoledgements. This study was funded by Alexandru Ioan Cuza University of Iasi, Romania, through the project 13/3.12.2015, code GI-2015-10, Grant Competition for young researchers of UAIC.

REFERENCES

- [1] Abrahao M.C., Gugliotta A.M., Silva R.D., Fujieda R.J.Y., Boscolo M., Gomes E. (2008): Ligninolytic activity from newly isolated basidiomycete strains and effect of these enzymes on the azo dye orange II decolourisation. – *Annals of Microbiology* 58: 427–432.
- [2] Aggelis G., Ehalotis C., Nerud F., Stoychev I., Lyberatos G., Zervakis G.I. (2002): Evaluation of white-rot fungi for detoxification and decolorization of effluents from the green olive debittering process. – *Applied Microbiology and Biotechnology* 59: 353–360.
- [3] Anastasi A., Prigione V., Varese G.C. (2010): Industrial dye degradation and detoxification by basidiomycetes belonging to different eco-physiological groups. – *Journal of Hazardous Materials* 177: 260–267.
- [4] Angelini P., Bistocchi G., Arcangeli A., Venanzoni R. (2012): Preliminary checklist of the macromycetes from Collestrada forest ecosystems in Perugia (Italy). – *Mycotaxon* 120: 505.
- [5] Arora D.S., Gill P.K. (2005): Production of ligninolytic enzymes by *Phlebia floridensis*. – *World Journal of Microbiology & Biotechnology* 21: 1021–1028.
- [6] ARS (Agricultural Research Service, United State Department of Agriculture, <http://nt.ars-grin.gov/fungaldatabases/fungushost/FungusHost.cfm>).
- [7] Asgher M., Noreen S., Bhatti H.N. (2010): Decolorization of dye containing textile industry effluents using *Ganoderma lucidum* IBL-05 in still culture. – *Water Environment Research* 82(4): 357–361.
- [8] Balaeş T., Mangalagiu I.I., Tănase C. (2013): Lignicolous macromycetes: potential candidates for bioremediation of synthetic dyes. – *Revista de Chimie* 64(9): 790–795.
- [9] Balaeş T., Tănase C., Butnariu C.D. (2014): The use of heavy metals in mycoremediation of synthetic dyes. – *Central European Journal of Biology* 9(7): 659–667.
- [10] Balaeş T., Tănase C. (2013): Optimization of nutritional conditions for the mycoremediation of synthetic dyes. – *Romanian Biotechnological Letters* 18(6): 8804–8811.
- [11] Balaraju K., Park K., Jahagirdar S., Kaviyaran V. (2010): Production of cellulase and laccase enzymes by *Oudemansiella radicata* using agro wastes under solid-state and submerged conditions. – *Research in Biotechnology* 1: 21–28.
- [12] Baldrian P. (2004): Purification and characterization of laccase from the white-rot fungus *Daedalea quercina* and decolorization of synthetic dyes by the enzyme. – *Applied Microbiology and Biotechnology* 63: 560–563.
- [13] Baldrian P. (2006): Fungal laccases occurrence and properties. – *FEMS Microbiology Reviews* 30: 215–242.

- [14] Barrasa J.M., Martínez A.T., Martínez M.J. (2009): Isolation and selection of novel basidiomycetes for decolorization of recalcitrant dyes. – *Folia Microbiologica* 54(1): 59–66.
- [15] Bernicchia A. (2005): *Fungi Europaei*, Vol. X: Polyporaceae s.l. – Edizioni Candusso, Bologna.
- [16] Bernicchia A., Gorjón S.P. (2010): *Corticaceae s.l. Fungi Europaei*, n°12. – Edizioni Candusso, Bologna.
- [17] Bhatti H.N., Akram N., Asgher M. (2008): Optimization of culture conditions for enhanced decolorization of Cibacrom Red FN-2BL by *Schizophyllum commune* IBL-6. – *Applied Biochemistry and Biotechnology* 149: 255–264.
- [18] Bisen P.S., Baghel R.K., Sanodiya B.S., Thakur G.S., Prasad G.B.K.S. (2010): *Lentinus edodes*: A macrofungus with pharmaceutical activities. – *Current Medicinal Chemistry* 17: 2419–2430.
- [19] Boer C.G., Obici L., Marques De Souza C.G., Peralta R.M. (2004): Decolorization of synthetic dyes by solid state cultures of *Lentinula (Lentinus) edodes* producing manganese peroxidase as the main ligninolytic enzyme. – *Bioresource Technology* 94: 107–112.
- [20] Bourbonnais R., Paice M.G., Reid I.D., Lanthier P., Yaguchi M. (1995): Lignin oxidation by laccase isozymes from *Trametes versicolor* and role of the mediator 2,2-azinobis (3-ethylbenzthiazoline-6-sulphonate) in kraft lignin depolymerisation. – *Applied Environmental Microbiology* 61: 1876–1880.
- [21] Casieri L., Anastasi A., Prigione V., Varese G. C. (2010): Survey of ectomycorrhizal, litter-degrading, and wood-degrading Basidiomycetes for dye decolorization and ligninolytic enzyme activity. – *Antonie van Leeuwenhoek* 98: 483–504.
- [22] Casieri L., Varese G.C., Anastasi A., Prigione V., Svobodova K., Marchisio V.F., Novoný Č. (2008): Decolorization and detoxication of reactive industrial dyes by immobilized fungi *Trametes pubescens* and *Pleurotus ostreatus*. – *Folia Microbiologica* 53(1): 44–52.
- [23] Cookson L.J. (1995): Reliability of poly B-411, a polymeric anthraquinone-based dye, in determining the rot type caused by wood-inhabiting fungi. – *Applied Environmental Microbiology* 61(2): 801–803.
- [24] D'Souza-Ticlo D., Sharma D., Raghukumar C. (2009): A thermostable metal-tolerant laccase with bioremediation potential from a marine-derived fungus. – *Marine Biotechnology* 11: 725–737.
- [25] Daniel G., Volc I.J., Kubatova E. (1994): Pyranose oxidase, a major source of H₂O₂ during wood degradation by *Phanerochaete chrysosporium*, *Trametes versicolor*, and *Oudemansiella mucida*. – *Applied and Environmental Microbiology* 60(7): 2524–2532.
- [26] Deacon J. W. (2006): *Fungal Biology, 4-th edition*. – Blackwell Publishing, Cornwell.
- [27] Dhoub A., Hamza M., Zouari H., Mechichi T., Hmidi R., Labat M., Martinez M.J., Sayadi S. (2005): Screening for ligninolytic enzyme production by diverse fungi from Tunisia. – *World Journal of Microbiology & Biotechnology* 21: 1415–1423.
- [28] Diamandis S., Perlerou C. (2001): The mycoflora of the chestnut ecosystem in Greece. – *Forest Snow and Landscape Research* 76(3): 499–504.
- [29] Eichlerová I., Homolka L., Lisá L., Nerud F. (2006a): The influence of extracellular H₂O₂ production on decolorization ability in fungi. – *Journal of Basic Microbiology* 46(6): 449–455.
- [30] Eichlerová I., Homolka L., Nerud F. (2006b): Ability of industrial dyes decolorization and ligninolytic enzymes production by different *Pleurotus* species with special attention on *Pleurotus calyptratus*, strain CCBAS 461. – *Process Biochemistry* 41: 941–946.
- [31] Eichlerová I., Homolka L., Nerud F. (2006c): Synthetic dye decolorization capacity of white rot fungus *Dichomitus squalens*. – *Bioresource Technology* 97: 2153–2159.

- [32] Elisashvili V., Kachlishvili E. (2009): Physiological regulation of laccase and manganese peroxidase production by white-rot basidiomycetes. – *Journal of Biotechnology* 144: 37–42.
- [33] Elisashvili V., Kachlishvili E., Khardziani T., Agathos S.N. (2010): Effect of aromatic compounds on the production of laccase and manganese peroxidase by white-rot basidiomycetes. – *Journal of Industrial Microbiology and Biotechnology* 37: 1091–1096.
- [34] Elisashvili V., Kachlishvili E., Tsiklauri N., Metreveli E., Khardziani T., Agathos S.N. (2009): Lignocellulose-degrading enzyme production by white-rot Basidiomycetes isolated from the forests of Georgia. – *World Journal of Microbiology and Biotechnology* 25: 331–339.
- [35] EOL (Enciclopedia of Life): <http://eol.org/>
- [36] Erkkilä R., Niemelä T. (1986): Polypores in the parks and forests of the city of Helsinki. – *Karstenia* 26: 1–40.
- [37] Fedorova T.V., Shakhova N.V., Klein O. I., Glazunova O. A., Maloshenok L. G., Kulikova N. A., Psurtseva N. V., Koroleva O. V. (2013): Comparative analysis of the ligninolytic potential of basidiomycetes belonging to different taxonomic and ecological groups. – *Applied Biochemistry and Microbiology* 49(6): 570–580.
- [38] Gbolagade J., Ajayi A., Oku I., Wankasi D. (2006): Nutritive value of common edible mushrooms from South Nigeria. – *Global Journal of Biotechnology & Biotechnology* 1(1): 16–21.
- [39] Gerault A. (2005): Florule évolutive des Basidiomycotina du Finistere. Heterobasidiomycetes. Tricholomatales. Version 2.1.
- [40] Gerault A. (2006): Florule évolutive des Basidiomycotina du Finistere. Homobasidiomycetes. Hymenochaetales, Lachnocladiales, Ganodermatales, Poriales, Polyporales, Schizophyllales, Version 1.0.
- [41] Gilbert G.S., Sousa W.P. (2002): Host specialization among wood-decay polypore fungi in a Caribbean mangrove forest. – *Biotropica* 34(3): 396–404.
- [42] Gramss G., Voigt K.-D., Kirsche B. (1999): Degradation of polycyclic aromatic hydrocarbons with three to seven aromatic rings by higher fungi in sterile and unsterile soils. – *Biodegradation* 10: 51–62.
- [43] Graż M., Jarosz-Wilkolazka A. (2011): Oxalic acid, versatile peroxidase secretion and chelating ability of *Bjerkandera fumosa* in rich and limited culture conditions. – *World Journal of Microbiology and Biotechnology* 27: 1885–1891.
- [44] Guillén Y., Palfner G., Machuca A. (2011): Screening for lignocellulolytic enzymes and metal tolerance in isolates of wood-rotting fungi from Chile. – *Intersciencia* 36(11): 860–868.
- [45] Guo L.Q., Lin S.X., Zheng X.B., Huang Z.R., Lin J.F. (2011): Production, purification and characterization of a thermostable laccase from a tropical white-rot fungus. – *World Journal of Microbiology and Biotechnology* 27: 731–735.
- [46] Haibo Z., Yinglong Z., Feng H., Peiji G., Jiachuan C. (2009): Purification and characterization of a thermostable laccase with unique oxidative characteristics from *Trametes hirsuta*. – *Biotechnology Letters* 31: 837–843.
- [47] Hansent L., Knudsen H. (1992, 1997): *Nordic Macromycetes, Vol II: Polyporales, Boletales, Agaricales, Russulales*. Nordsvamp, Copenhagen, 474 p; *Vol. III: Heterobasidioid Aphyllophoroid and Gastromycetoid Basidiomycetes*. – Nordsvamp, Copenhagen, 444 p.
- [48] Hattori, T. (2005): Diversity of wood-inhabiting polypores in temperate forest with different vegetation types in Japan. – *Fungal Diversity* 18: 73–88.
- [49] Hildén K., Martínez A.T., Hatakka A., Lundell T. (2005): The two manganese peroxidases Pr-MnP2 and Pr-MnP3 of *Phlebia radiata*, a lignin-degrading basidiomycete, are phylogenetically and structurally divergent. – *Fungal Genetics and Biology* 42: 403–419.

- [50] Kalmış E., Yaşa I., Kalyoncu F., Pazarbaşı B., Koçyiğit, A. (2008): Ligninolytic enzyme activities in mycelium of some wild and commercial mushrooms. – *African Journal of Biotechnology* 7 (23): 4314–4320.
- [51] Knežević A., Milovanović I., Stajić M., Vukojević J. (2013): *Trametes suaveolens* as ligninolytic enzyme producer. – *Matica Srpska Journal of Natural Sciences* 124: 437–444.
- [52] Kum H., LEE S., Ryu S., Choi H.T. (2011): Degradation of endocrine disrupting chemicals by genetic transformants with two lignin degrading enzymes in *Phlebia tremellosa*. – *The Journal of Microbiology* 49 (5): 824–827.
- [53] Lee J.S., Woo E.J., Oh K.H., Kim J.-J., Lim Y.W. (2010): The first report of two species of *Polyporus* (Polyporaceae, Basidiomycota) from South Korea. – *The Journal of Microbiology* 48(6): 748–753.
- [54] Levin L., Papinutti L., Forchiassin F. (2004): Evaluation of Argentinian white rot fungi for their ability to produce lignin-modifying enzymes and decolorize industrial dyes. – *Bioresource Technology* 94: 169–176.
- [55] Levin L., Ramos A., Forchiassin F. (2002): Copper induction of lignin-modifying enzymes in the white-rot fungus *Trametes trogii*. – *Mycologia* 94(3): 377–383.
- [56] Li X., Jia R., Li P., ANG S. (2009): Response surface analysis for enzymatic decolorization of Congo red by manganese peroxidase. – *Journal of Molecular Catalysis B: Enzymatic* 56: 1–6.
- [57] Liers C., Bobeth C., Pecyna M., Ullrich R., Hofrichter M. (2010): DyP-like peroxidases of the jelly fungus *Auricularia auricula-judae* oxidize nonphenolic lignin model compounds and high-redox potential dyes. – *Applied Microbiology and Biotechnology* 85: 1869–1879.
- [58] Liu W., Chao Y., Yang X., Bao H., Qian S. (2004): Biodecolorization of azo, anthraquinonic and triphenylmethane dyes by white-rot fungi and a laccase-secreting engineered strain. – *Journal of Industrial Microbiology and Biotechnology* 31: 127–132.
- [59] Lorenzo M., Moldes D., Sanromán M.A. (2006): Effect of heavy metals on the production of several laccase isoenzymes by *Trametes versicolor* and on their ability to decolourise dyes. – *Chemosphere* 63: 912–917.
- [60] Mäkelä M.R., Hildén K.S., Hakala T.K., Hatakka A., Lundell T.K. (2006): Expression and molecular properties of a new laccase of the white rot fungus *Phlebia radiata* grown on wood. – *Current Genetics* 50: 323–333.
- [61] Matsushita N., Suzuki K. (2005): Identification of *Armillaria* species in Japan using PCR-RFLP analysis of rDNA intergenic spacer region and comparison of *Armillaria* species in the world. – *Journal for Forest Research* 10: 173–179.
- [62] Minter D.W., Rodríguez Hernández M., Mena Portales J. (2001): *Fungi of the Caribbean: an annotated checklist*. – PDMS Publishing, Isleworth.
- [63] Moreira M.T., Mielgo I., Feijoo G., Lema J.M. (2000): Evaluation of different fungal strains in the decolourisation of synthetic dyes. – *Biotechnology Letters* 22: 1499–1503.
- [64] Moturi B., Singara Charya M.A. (2009): Decolourisation of crystal violet and malachite green by fungi. – *Science World Journal* 4(4): 28–33.
- [65] Nagy L.G., Riley R., Tritt A., Adam C., Daum C., Floudas D., Sun H., Yadav J.S., Pangilinan J., Larsson H.K., Matsuura K., Barry K., LaButti K., Kuo R., Ohm R.A., Bhattacharya S.S., Shirouzu T., Yoshinaga Y., Martin F.M., Grigoriev I.V., Hibbett D.S. (2015): Comparative genomics of early-diverging mushroom-forming fungi provides insights into the origins of lignocellulose decay capabilities. – *Molecular Biology and Evolution* doi:10.1093/molbev/msv337.
- [66] Nakasone K.K., Burdsall JR. H.H. (1995): *Phlebia* species from eastern and southeastern United States. – *Mycotaxon* 54: 335–359.
- [67] Negrean G., Anastasiu P. (2004): *Hirneola auricula-judae* in Romania. – *Contribuții Botanice, Grădina Botanică "Alexandru Borza" Cluj-Napoca* 39: 21–39. (in Romanian)
- [68] Nerud F., Mišurcova Z. (1996): Distribution of ligninolytic enzymes in selected white-rot fungi. – *Folia Microbiologica* 41(3): 264–256.

- [69] Nikitina O.V., Shleev S.V., Gorshina E.S., Rusinova T.V., Serezhenkov V.A., Burbaev D.S., Belovolova L.V., Yaropolov A.I. (2005): Isolation and purification of enzymes from ligninolytic complex of the basidial fungus *Trametes pubescens* (Schumach.) Pilát and study of their properties. – *Biochemistry (Moscow)* 70(11): 1274–1279.
- [70] Novotný Č., Cajthaml T., Svobodová K., Šušla M., Šásek V. (2009): *Irpex lacteus*, a white-rot fungus with biotechnological potential. – *Folia Microbiologica* 54(5): 375–390.
- [71] Novotný Č., Troš N., Šušla M., Svobodova K., Mikeskova H., Valkova H., Malachova K., Pavko A. (2012): The use of the fungus *Dichomitus squalens* for degradation in rotating biological contactor conditions. – *Bioresource Technology* 114: 241–246.
- [72] Orth A.B., Royse D.J., Tien M. (1993): Ubiquity of lignin-degrading peroxidases among various wood-degrading fungi. – *Applied and Environmental Microbiology* 59(12): 4017–4023.
- [73] Otieno W., Pérez sierra A., Termorshuizen A. (2003): Characterization of *Armillaria* isolates from tea (*Cammelia sinensis*) in Kenya. – *Mycologia* 95(1): 160–175.
- [74] Palmieri G., Cennamo G., Sannia G. (2005): Remazol Brilliant Blue R decolourisation by the fungus *Pleurotus ostreatus* and its oxidative enzymatic system. – *Enzyme and Microbial Technology* 36: 17–24.
- [75] Pant D., Adholeya A. (2009): Concentration of fungal ligninolytic enzymes by ultrafiltration and their use in distillery effluent decolorization. – *World Journal of Microbiology and Biotechnology* 25: 1793–1800.
- [76] Pegler D.N., Læssøe T., Spooner B.M. (1995): British puffballs, earthstars and stinkhorns. An account of the British Gasteroid Fungi. – Royal Botanic Gardens, Kew.
- [77] Petersen R.H., Hughes K.W., Redhead S.A., Psurtseva N., Methven A.S. (1999): Mating systems in the Xerulaceae (Agaricales, Basidiomycotina): *Flammulina*. – *Mycoscience* 40: 411–426.
- [78] PKB (Plantwise Knowledge Bank, <http://www.plantwise.org/KnowledgeBank/Datasheet.aspx?dsid=7002>)
- [79] Pozdnyakova N.N., Nikiforova S.V., Makarov O.E., Turkovskaya O.V. (2010): Effect of polycyclic aromatic hydrocarbons on laccase production by white rot fungus *Pleurotus ostreatus* D1. – *Applied Biochemistry and Microbiology* 47(5): 543–548.
- [80] Qin G.-F., Zhao J., Korhonen K. (2007): A study on intersterility groups of *Armillaria* in China. – *Mycologia* 99(3): 430–441.
- [81] Ranadive K.R., Vaidya J.G., Jite P.K., Ranade V.D., Bhosale S.R., Rabba A.S., Hakimi M., Deshpande G.S., Rathod M.M., Forutan A., Kaur M., Naik-Vaidya C.D., Bapat G.S., Lamrood P. (2011): Checklist of Aphyllophorales from the Western Ghats of Maharashtra state, India. – *Mycosphere* 2(2): 91–114.
- [82] Rees B.J., Strid Å. (2001): Relationships between Australian and Northern hemisphere *Gymnopilus* species 1: new species and common misconceptions regarding earlier names. – *Australasian Mycologist* 20(1): 21–48.
- [83] Rigas F., Dritsa V. (2006): Decolorization of a polymeric dye by selected fungal strains in liquid cultures. – *Enzyme and Microbial Technology* 39: 120–124.
- [84] Robinson T., Chandran B., Nigam P. (2001): Studies on the decolourisation of an artificial textile-effluent by white-rot fungi in N-rich and N-limited media. – *Applied Microbiology and Biotechnology* 57: 810–813.
- [85] Robinson T., Nigam P.S. (2008): Remediation of textile dye waste water using a white-rot fungus *Bjerkandera adusta* through solid state fermentation (SSF). – *Applied Biochemistry and Biotechnology* 151: 618–628.
- [86] Ryu S.H., Lee A.Y., Kim M. (2008): Molecular characteristics of two laccase from basidiomycete fungus *Polyporus brumalis*. – *The Journal of Microbiology* 46(1): 62–69.
- [87] Ryvarden L., Gilbertson R.L. (1993, 1994): *European Polypores*, Vol I and Vol II. – Fungiflora, Oslo.
- [88] Sankaran K.V., Bridge P.D., Gokulapalan C. (2005): *Ganoderma* diseases of perennial crops in India-an overview. – *Mycopathologia* 159: 143–152.

- [89] Simone D., Annesi T. (2012): Occurrence of *Ganoderma adpersum* on *Pinus pinea*. – *Phytopathologia Mediterranea* 51(2): 374–382.
- [90] SIPMP (State Integrated Pest Management Program, University of California, <http://ipm.ucanr.edu/index.html>)
- [91] Smith A.H., Hessler L.R. (1968): The North American species of *Pholiota*. – Hafner Publishing Company, New York.
- [92] Steffen K.T., Schubert S., Tuomela M., Hatakka A., Hofrichter M. (2007): Enhancement of bioconversion of high-molecular mass polycyclic aromatic hydrocarbons in contaminated non-sterile soil by litter-decomposing fungi. – *Biodegradation* 18: 359–369.
- [93] Stoytchev I., Nerud F. (2000): Ligninolytic enzyme complex of *Armillaria* spp. – *Folia Microbiologica* 45(3): 248–250.
- [94] Šušla M., Novotný Č., Svobodová K. (2007): The implication of *Dichomitus squalens* laccase isoenzymes in dye decolorization by immobilised fungal culture. – *Bioresource Technology* 98: 2109–2115.
- [95] Szczepkowski A. (2007): Macromycetes in the dendrological Park of the Warsaw Agricultural University. – *Acta Mycologica* 42(2): 179–186.
- [96] Tinoco R., Verdin J., Vasquez-Duhalt R. (2007): Role of oxidizing mediators and tryptophan 172 in the decoloration of industrial dyes by the versatile peroxidase from *Bjerkandera adusta*. – *Journal of Molecular Catalysis B: Enzymatic* 46: 1–7.
- [97] Tomšovský M., Homolka L. (2003): Laccase and other ligninolytic enzyme activities of selected strains of *Trametes* spp. from different localities and substrates. – *Folia Microbiologica* 48(3): 413–416.
- [98] Ungureanu C., Balaş T., Favier L., Tănase C., Bahrim G. (2015): White-rot fungus implications in clofibric acid biodegradation. – *Romanian Biotechnological Letters* 20(3): 10388-10395.
- [99] Vaithanomsat P., Chedchant J., Kreetachat T., Kosugi A., Apiwatanapiwat W., Thanapase W., Chuntranuluck S., Mori Y. (2012): Improvement of lignin-degrading enzymes production from the white-rot fungus (*Lentinus strigosus*) and its application in synthetic dye removal. – *African Journal of Microbiology Research* 6(1): 137–148.
- [100] Valderrama B., Oliver P., Medrano-Soto A., Vazquez-Duhalt R. (2003): Evolutionary and structural diversity of fungal laccases. – *Antonie van Leeuwenhoek* 84: 289–299.
- [101] Valentiin L., Kluczek-Turpeinen B., Oivanen P., Hatakka A., Steffen K., Tuomela M. (2009): Evaluation of basidiomycetous fungi for pretreatment of contaminated soil. – *Journal of Chemical Technology and Biotechnology* 84: 851–858.