

ASSESSMENT OF THE EXTERNAL EXPOSURE DOSE TO HUMANS FROM ^{137}Cs AND ^{90}Sr IN THE COASTAL WATERS OF THE BALTIC SEA NEAR LITHUANIA

DAUNARAVIČIENĖ, A. * – JASAITIS, D.

*Department of Physics, Faculty of Fundamental Sciences,
Vilnius Gediminas Technical University
Sauletekio Ave 11, LT-10223 Vilnius, Lithuania
(phone: +370-5-274-4834)*

**Corresponding author
e-mail: Asta.Daunaraviciene@vgtu.lt.*

(Received 24th Nov 2016; accepted 16th Jan 2017)

Abstract. At present the Baltic Sea is considered to be the most contaminated with anthropogenic radionuclides in comparison to any other part of the World Ocean. Anthropogenic radionuclides (mainly ^{137}Cs and ^{90}Sr) found in the seawater are sources of the external exposure that contributes to the total radiation exposure to humans. The variations of activity concentrations of ^{137}Cs and ^{90}Sr in the water of the Baltic Sea near Lithuanian coast in 1985–2013 were analyzed. External exposure dose from these radionuclides to humans due to immersion in the seawater were calculated using the results of the measurements. An average external exposure dose from ^{137}Cs ranged between $6.28 \text{ nSv}\cdot\text{h}^{-1}$ to $1.5 \text{ nSv}\cdot\text{h}^{-1}$, from ^{90}Sr – between $1.73 \text{ nSv}\cdot\text{h}^{-1}$ to $0.53 \text{ nSv}\cdot\text{h}^{-1}$.

Keywords: *activity concentration, ionizing radiation, radionuclides, radioactivity, marine ecosystem*

Introduction

All living organisms, including humans, are being exposed to ionizing radiation at all times. Natural radiation comes from many sources including more than 60 naturally-occurring radioactive materials found in air, soil and water. Humans are also exposed to natural radiation from cosmic rays. According to the United Nations Scientific Committee on the Effects of Atomic Radiation report (UNSCEAR, 2010), the worldwide average radiation exposure from natural sources is approximately 2.4 mSv per year. However, the exposure to humans from ionizing radiation of natural sources is a continuing and inescapable feature of life on Earth.

Humans exposure to ionizing radiation also comes from anthropogenic sources. ^{137}Cs and ^{90}Sr are one of the most hazardous anthropogenic radionuclides due to their long physical and biological half-life (about 30 years) (IAEA, 2005). Anthropogenic radionuclides are penetrated in the marine environment mostly as a result of nuclear explosions, accidents at nuclear power plants and due to the operation of nuclear industry (HELCOM, 2009; Zalewska and Suplińska, 2013).

Review of Literature

For the first time ^{137}Cs and ^{90}Sr were penetrated into the Baltic Sea with the global fallout as a result of the tests of nuclear weapons in the atmosphere, particularly in the 1950s and 1960s (Livingston and Povinec, 2000; Ikäheimonen et al., 2009; Ilus, 2007). According to data published in Helsinki Commission proceedings (HELCOM, 2013), the total inputs of weapons test ^{137}Cs and ^{90}Sr into the Baltic Sea were 800 TBq and 500

TBq, respectively. During a long period of time this sea was contaminated by the North Sea waters where radioactive waste was discharged from various nuclear reprocessing plants (HELCOM, 2009; Zalewska and Lipska, 2006; Saniewski, 2013). The increase of the global fallout proceeded up to 1963, when a treaty concerning nuclear and thermonuclear weapon test stopped in the atmosphere, hydrosphere and cosmos were signed. This moment was the beginning of the decrease in radioactive contamination of the marine environment. The process was observed up to 1986. The average activity concentrations of ^{137}Cs and ^{90}Sr in the seawater until 1986 were $12 \text{ Bq}\cdot\text{m}^{-3}$ and $24 \text{ Bq}\cdot\text{m}^{-3}$, respectively (Styro et al., 1990).

The radiological situation in the Baltic Sea changed after the Chernobyl Nuclear Power Plant (ChNPP) accident, which became the main source of ^{137}Cs after 1986 (Ikäheimonen et al., 2009; Juranová et al., 2015). After the accident, an average activity concentration of ^{137}Cs in the surface waters of the Baltic Sea grew by more than an order of magnitude relative to the radioactive background ($12 \text{ Bq}\cdot\text{m}^{-3}$) formed after nuclear weapons tests in the atmosphere (Styro et al., 1990). According to data calculations (HELCOM, 2013), direct input of ^{137}Cs into the Baltic Sea from ChNPP accident was estimated to be 4700 TBq. The Chernobyl fallout was scattered very unevenly over the Baltic Sea area (Weiss, 2011). The Bothnian Sea and the Gulf of Finland were both regarded as the most contaminated regions of the Baltic (Zalewska and Lipska, 2006). In 1986, the respective average concentrations of ^{137}Cs in these regions were $480 \text{ Bq}\cdot\text{m}^{-3}$ and $500 \text{ Bq}\cdot\text{m}^{-3}$, whereas in the Baltic Proper the average activity concentration of this radionuclide was only $150 \text{ Bq}\cdot\text{m}^{-3}$ (Zalewska and Suplińska, 2013). Much less contaminated were the Bothnian Bay, the Belt Sea and the southern Baltic Proper where, in 1986, the average concentration was $84 \text{ Bq}\cdot\text{m}^{-3}$ (Zalewska and Suplińska, 2013). The activity concentration of ^{90}Sr following the accident of the ChNPP changed negligibly (Saniewski, 2013; Zaborska et al., 2014). The direct input of this radionuclide was estimated to be 80 TBq (Ikäheimonen et al., 2009). During a period of time, a leveling of the ^{137}Cs activity concentration in the surface waters took place; the end of this process may be ascribed to 1989, when its activity concentration appeared to be about $150 \text{ Bq}\cdot\text{m}^{-3}$ (Styra et al., 2008). This time was used as the beginning of self-purification process of the Baltic Sea from ^{137}Cs (Styra et al., 2008). During the period 1992–2010 concentrations of ^{137}Cs in the surface waters decreased in all parts of the Baltic Sea. The average activity concentration of ^{137}Cs in the Baltic Proper decreased to $40 \text{ Bq}\cdot\text{m}^{-3}$ in 2010. Concentrations in the Western Baltic and in the Gulf of Finland were lower at around $35 \text{ Bq}\cdot\text{m}^{-3}$ and $29 \text{ Bq}\cdot\text{m}^{-3}$, respectively. The highest concentrations were reported in the Archipelago and Aland Sea (equal to $44 \text{ Bq}\cdot\text{m}^{-3}$) (HELCOM, 2013). At present, concentrations of ^{137}Cs are relatively uniform in all regions of the Baltic and remain at approximately similar levels mainly because of the transport and mixing of water masses (Zalewska and Suplińska, 2013). ^{90}Sr concentration in the Baltic seawater varied in general from $5 \text{ Bq}\cdot\text{m}^{-3}$ to $15 \text{ Bq}\cdot\text{m}^{-3}$. The ^{90}Sr concentration decreases slowly with time and its behavior in seawater is different from ^{137}Cs (HELCOM, 2013).

The decrease of activity concentrations of these radionuclides in the coastal waters near Lithuania was observed too (Styro et al., 2012). However, radionuclides in the coastal waters are sources of external exposure that contribute to the total radiation exposure to humans.

The aim of this work is to analyze the average activity concentrations of ^{137}Cs and ^{90}Sr in the waters of the Baltic Sea near Lithuania and to assess the external exposure dose to humans due to these radionuclides.

Materials and Methods

The samples of the sea surface water for determining activity concentrations of ^{137}Cs and ^{90}Sr were taken in the Baltic Sea at the distance of 10 m from the seashore of Lithuania at 0.5 meter depth. Water samples were taken near Juodkrante (Fig. 1). The minimum volume of the analyzed water sample was 40–50 liters. The temperature, specific electric conductivity, water current direction, wind speed and direction were registered during the time of sampling.



Figure 1. Sample locations of the seawater

During the whole research period, the determination of activity concentrations of ^{137}Cs and ^{90}Sr in the coastal surface waters of the Baltic Sea were carried out using a single technique – radiochemical analysis and the same type of measuring equipment (Styro et al., 2011; Styro et al., 2012). This research technique consists of some main stages: the concentration of ^{137}Cs and ^{90}Sr together with a stable carriers, radiochemical cleaning and activity measuring.

The ^{137}Cs and ^{90}Sr were concentrated from the same seawater samples. After the radiochemical cleaning the yield of cesium was determined gravimetrically in the form of $\text{Cs}_3\text{Sb}_2\text{I}_9$. It varied within a range of 60–80 %. The activity of ^{137}Cs samples was registered by gamma spectrometer (CANBERRA) with HPGe detector (resolution 2 keV, efficiency 15 %). The determination error for ^{137}Cs amounted to 10 %.

^{90}Sr was measured by ^{90}Y emission using a low level background beta radiometer. A stable strontium yield was determined by the atomic absorption spectrometer and Y – gravimetrically in Y_2O_3 form. The yield values varied within a range of 60–80 %. The determination error for ^{90}Sr activity concentration amounted to 15 %.

The external exposure dose to humans from ^{137}Cs and ^{90}Sr was calculated with the following equation (Eckerman and Ryman, 1993; IAEA, 2001):

$$E_{\text{im}} = f DF_{\text{im}} C_{\text{v}}, \quad (\text{Eq. 1})$$

where E_{im} – external exposure dose to humans from radionuclide ($\text{Sv}\cdot\text{h}^{-1}$); f – exposure time to external radiation ($\text{h}\cdot\text{y}^{-1}$); DF_{im} – dose conversion coefficient for water immersion ($\text{Sv}\cdot\text{m}^3\cdot\text{Bq}^{-1}\cdot\text{y}^{-1}$), C_{v} – activity concentration of the radionuclide in the seawater ($\text{Bq}\cdot\text{m}^{-3}$).

Results and Discussion

The average values of ^{137}Cs and ^{90}Sr activity concentrations in the coastal waters of the Baltic Sea near Lithuania (Juodkrante) in 1985–2013 are illustrated in Fig. 2. An average activity concentration of ^{137}Cs was $18 \text{ Bq}\cdot\text{m}^{-3}$ and ^{90}Sr – $28 \text{ Bq}\cdot\text{m}^{-3}$ before the ChNPP accident.




Figure 2. Average values of ^{137}Cs and ^{90}Sr activity concentrations in the coastal waters of the Baltic Sea near Lithuania (Juodkrante) obtained in 1985–2013 (data from 1985–2003 after Nuclear Hydrophysics laboratory of Vilnius Gediminas Technical University)

After this accident the average activity concentration of ^{137}Cs considerably increased. The highest average values of ^{137}Cs in the coastal waters varied in the range from $95 \text{ Bq}\cdot\text{m}^{-3}$ to $117 \text{ Bq}\cdot\text{m}^{-3}$ were found from 1988 to 1995. This increase was attributed to the transport of more polluted water from the northern Baltic as well as to considerable riverine discharges (Styra et al., 2008). Since 1995 the situation stabilized and the activity concentration of ^{137}Cs started to decrease as a consequence of many processes, mainly radioactive decay, bioaccumulation and sedimentation (Styro et al., 2012). However, an unexpected increases of the average values of ^{137}Cs were also observed in 1999 and 2002. During 2003–2013, an average activity concentration in the coastal waters varied in the range from $63 \text{ Bq}\cdot\text{m}^{-3}$ to $30 \text{ Bq}\cdot\text{m}^{-3}$. However, in 2013 this average value exceeded almost two times the value found in 1985.

After ChNPP accident activity concentration of ^{90}Sr slightly increased. In 1986, the average concentration was $33 \text{ Bq}\cdot\text{m}^{-3}$. However, in 1987 it reached the value found before the accident (Fig. 2). During 1987–2002, an average activity concentration in the coastal

waters varied in the range from $25 \text{ Bq}\cdot\text{m}^{-3}$ to $18 \text{ Bq}\cdot\text{m}^{-3}$. In 2003, an average value of ^{90}Sr decreased to $13 \text{ Bq}\cdot\text{m}^{-3}$. Since then, the average activity concentration of ^{90}Sr in the seawater varied within a relatively narrow range from $13 \text{ Bq}\cdot\text{m}^{-3}$ to $10 \text{ Bq}\cdot\text{m}^{-3}$. The average activity concentration of ^{90}Sr in 2013 exceeded almost three times the value found in 1985.

According to data published in (Zalewska and Suplińska, 2013; Druteikienė et al., 2011; Lujanienė et al., 2010) and variations of average activity concentrations of ^{137}Cs and ^{90}Sr in the coastal waters near Lithuania, the Baltic is still the most contaminated seas in the world.

Certainly, variations of activity concentration of radionuclides in seawater have an impact on external exposure to humans. The summer season in the coastal Baltic Sea near Lithuania is from June to September. According to (Kahru et al., 2015), the number of days with the seawater warmer than 17°C have almost doubled (from 29 to 56 days) over the past 30 years. Approximately, humans are being exposed when swimming in the seawater from few to few dozen hours per year.

External exposure dose to humans calculated using the average activity concentrations of ^{137}Cs and ^{90}Sr in the seawater and equation (1) are presented in Figs. 3 and 4.




Figure 3. External exposure dose to humans from ^{137}Cs in the coastal waters of the Baltic Sea (Juodkrante) in 1985–2013

According to data obtained by the calculations, an average external exposure dose to humans from ^{137}Cs in the coastal waters in 1985–2013 ranged between $6.28 \text{ nSv}\cdot\text{h}^{-1}$ and $1.5 \text{ nSv}\cdot\text{h}^{-1}$ (Fig. 3), from ^{90}Sr – between $1.73 \text{ nSv}\cdot\text{h}^{-1}$ and $0.53 \text{ nSv}\cdot\text{h}^{-1}$ (Fig. 4). The highest exposure dose from ^{137}Cs was calculated in 1990, from ^{90}Sr – in 1986. After thirty years after ChNPP accident the exposure dose from ^{137}Cs decreased four times. The external exposure dose from ^{90}Sr due to immersion in the water decreased slower, almost three times.




Figure 4. External exposure dose to humans from ^{90}Sr in the coastal waters of the Baltic Sea (Juodkrante) in 1985–2013

The external exposure from ^{137}Cs and ^{90}Sr in the seawater to humans when swimming is unavoidable. However, the average external exposure dose was very small in comparison to natural sources and did not make any risk for humans health.

Conclusions

Activity concentrations of ^{137}Cs and ^{90}Sr in the Baltic Sea waters decrease in time due to their natural radioactive decay and the decrease of the environmental radioactive pollution. According to average values, the activity concentration of ^{137}Cs in the coastal waters near Lithuania has been decreasing since 1990. The average activity concentration of ^{137}Cs decreased from $117 \text{ Bq}\cdot\text{m}^{-3}$ found in 1990 to $30 \text{ Bq}\cdot\text{m}^{-3}$ in 2013. However, the average activity concentration of ^{137}Cs in 2013 exceeded almost two times the value found in 1985. During the same period the average activity concentration of ^{90}Sr decreased almost three times, i.e. to $10 \text{ Bq}\cdot\text{m}^{-3}$.

During the period from 1985 to 2013, an average external exposure dose to humans from ^{137}Cs in the coastal waters varied in the range from $6.28 \text{ nSv}\cdot\text{h}^{-1}$ to $1.5 \text{ nSv}\cdot\text{h}^{-1}$, from ^{90}Sr – from $1.73 \text{ nSv}\cdot\text{h}^{-1}$ to $0.53 \text{ nSv}\cdot\text{h}^{-1}$. The level of the external exposure was considerably lower than its limit ($1 \text{ mSv}\cdot\text{y}^{-1}$) and did not make any risk for humans health.

REFERENCES

- [1] Druteikienė, R., Morkūnienė, R., Lukšienė, B. (2011): Distribution of artificial radionuclides in the Baltic seaside environment. – Lithuanian Journal of Physics 51(1): 75-81.
- [2] Eckerman, K. F., Ryman, J. C. (1993): External exposure to radionuclides in air, water and soil. – US Environmental Protection Agency, Washington.

- [3] HELCOM, (2009): Radioactivity in the Baltic Sea, 1999–2006 HELCOM thematic assessment. – Baltic Sea environment proceedings No. 117, Helsinki.
- [4] HELCOM, (2013): Thematic assessment of long-term changes in radioactivity in the Baltic Sea, 2007–2010. – Baltic Sea environment proceedings No. 135, Helsinki.
- [5] IAEA, (2001): Generic models for use in assessing the impact of discharges of radioactive substances to the environment. – International Atomic Energy Agency, Vienna.
- [6] IAEA, (2005): Worldwide marine radioactivity studies (WOMARS). Radionuclide levels in oceans and seas. – International Atomic Energy Agency, Vienna.
- [7] Ikäheimonen, T. K., Outola, I., Vartti, V. P., Kolilainen, P. (2009): Radioactivity in the Baltic Sea: inventories and temporal trends of ^{137}Cs and ^{90}Sr in water and sediments. – *Journal of Radioanalytical and Nuclear Chemistry* 282(2): 419-425.
- [8] Ilus, E. (2007): The Chernobyl accident and the Baltic Sea. – *Boreal Environment Research* 12: 1-10.
- [9] Juranová, E., Hanslik, E., Marešová, D. (2015): Temporal development of radiocaesium and radiostrontium concentrations in the hydrosphere – methods and evaluation. – *Water, Air, & Soil Pollution* 226(10): 335.
- [10] Kahru, M., Elmgren, R., Savchuk O. P. (2016): Changing seasonality of the Baltic Sea. – *Biogeosciences Discussions* 12: 18855-18882.
- [11] Livingston, H. D., Povinec, P. P. (2000): Anthropogenic marine radioactivity. – *Ocean & Coastal Management* 43(8-9): 689-712.
- [12] Lujanienė, G., Beneš, P., Štamberg, K., Jokšas, K., Vopalka, D., Radžiūtė, E., Šilobritienė, B., Šapolaitė, J. (2010): Experimental study and modeling of ^{137}Cs sorption behaviour in the Baltic Sea and the Curonian Lagoon. – *Journal of Radioanalytical and Nuclear Chemistry* 286(2): 361-366.
- [13] Saniewski, M. (2013): Spatiotemporal variations of the ^{90}Sr in the southern part of the Baltic Sea over the period of 2005–2010. – *The Scientific World Journal*, 1013, ID 276098. dx.doi.org/10.1155/2013/276098.
- [14] Styra, D., Morkuniene, R., Daunaraviciene, A. (2008): Self-purification process of the Baltic Sea from ^{137}Cs radionuclide in 1986–2006. – In: Selected papers of the 7th international conference „Environmental engineering“ 1: 362-367.
- [15] Styro, D. B., Bumyalene, Zh. V., Kadzhene, G. I., Kleiza, J. V., Lukinskene, M. V., Pogrebnyak, E. V. (1990): Structure of the specific activity fields of man-made radionuclides extracted from the surface waters of the Baltic Sea in autumn of 1986 and 1987. – *Atomic Energy* 68: 14-18.
- [16] Styro, D., Morkuniene, R., Daunaraviciene, A. (2011): Radionuclide ^{90}Sr volume activity variations at the Baltic Sea coast near Juodkrante. – In: Selected papers of the 8th international conference „Environmental engineering“ 1: 357-361.
- [17] Styro, D., Morkūnienė, R., Daunaravičienė, A. (2012): On variations of volumetric activity of ^{90}Sr and ^{137}Cs in the Baltic Sea coastal waters near the shore of Lithuania in 2005–2009. – *Journal of Radioanalytical and Nuclear Chemistry* 293(3): 923-929.
- [18] UNSCEAR, (2010). Sources and effects of ionizing radiation. – New York: United Nations.
- [19] Weiss, D. (2011): Distribution pattern of artificial radionuclides in the Baltic Sea in the special event of the Chernobyl fallout. – *Isotopes in Environmental and Health Studies* 47(3): 254-264.
- [20] Zaborska, A., Winogradow, A., Pempkowiak, J. (2014): Caesium-137 distribution, inventories and accumulation history in the Baltic Sea sediments. – *Journal of Environmental Radioactivity* 127: 11-25.
- [21] Zalewska, T., Lipska, J. (2006): Contamination of the eastern Baltic Sea with ^{137}Cs and ^{90}Sr over the period 2000–2004. – *Journal of Environmental Radioactivity* 91(1-2): 1-14.
- [22] Zalewska, T., Suplińska, M. (2013): Anthropogenic radionuclides ^{137}Cs and ^{90}Sr in the southern Baltic Sea ecosystem. – *Oceanologia* 55(3): 485-517.