

Impacts of arbuscular mycorrhizal fungi on plant growth and yield of three pepper genotypes

Hong Duc NGUYEN – Au Trung VO – Katalin POSTA

Institute of Genetic, Microbiology and Biotechnology, Szent Istvan University, Hungary. Email: hongduc.real@gmail.com

Abstract: Field study was implemented at the experiment station of Szent Istvan University, Godollo, Hungary to investigate the effects of arbuscular mycorrhizal (AM) fungi on plant growth, yield of three pepper cultivars during the plant growth. The experiment was arranged in randomized complete block design with two factors of pepper cultivars and AM applications. Three pepper varieties were Karpia, Karpex, Kaptur while AM applications consisted of AM mixture with 6 different AM species and non-inoculated treatment. Greater shoot fresh weight in AM plants were observed while values of root fresh weight, shoot and root dry weight remained unchanged in inoculated plants, except the increases recorded in Karpex cultivar. AM application also enhanced significantly fruit yield in Karpia and Kaptur but not in Karpex cv. In addition, AM treated plants showed improved root colonization of AM fungi.

Keywords: Arbuscular mycorrhizal fungi, pepper, growth, fruit yield.

Introduction

Pepper (*Capsicum annuum* L.) is one of the main horticultural vegetables and cultivated worldwide due to important nutritional and economic values. In addition to pepper cultivars, pepper production and quality are diversified owing to various stress conditions that which often loses 70 % of yield forming a barrier in pepper cultivation (Gajanayake et al., 2011). The exploitation of symbiotic feature of AM fungi is one of the efficient approaches to improve crop tolerance to unfavored environment (Birhane et al., 2012). In fact, AM fungi are probably the most ubiquitous soil microbe that can colonize 80% of terrestrial plant species consisted of many important crops (Smith and Read, 2008). Many beneficial effects from mycorrhizal colonization including increased seedling survival, enhanced growth, fruit yield and quality, uniformity of horticultural crops, and earlier and increased flowering as well as induced resistance to abiotic and biotic stresses have been reported (Estrada-Luna et al., 2000; Estrada-Luna & Davies, 2003; Garmendia et al., 2004; Alejo-Iturvide et al., 2008; Mena-Violante et al., 2006; Kaya et al., 2009; Ruscitti et al., 2011; Ortas et al., 2011; Franco et al., 2013).

Although use of the AMF is widely investigated in many plants, little attention has been paid to use different cultivars as target plants for inoculation. Therefore, the aim of this study was to examine the potential of AM mixture for growth, fruit yield in different pepper genotypes under field conditions.

Materials and methods

Three sweet pepper (*Capsicum annuum* L.) hybrids, Karpia, Karpex and Kaptur were used for this study at experimental station of Szent István University, Gödöllő, Hungary. Under field, treatments including inoculation of AM mixture or no inoculation (control) and three cultivars were arranged in randomized complete block design with 30 replications each treatment. Pepper seedlings were grown in greenhouse in 7 weeks before planting in field. Mycorrhizal Inoculation with commercial product Symbivit® (mixture of *Glomus intraradices*, *G. mosseae*, *G. etunicatum*, *G. claroideum*, *G. microaggregatum*,

G. geosporum) (Symbiom Ltd., Lanskrout, Czech Republic; www.symbiom.cz), was utilized at 25 g of inoculum per pepper seedling at planting. **Assessment of mycorrhizal colonization** by gridline intersect method (Giovannetti & Mosse, 1980) after staining roots (Vierheilig et al.; 1998) with five plants per treatment. **Plant biomass and fruit yield:** The pepper harvesting was performed randomly by hand at the biological maturity stage and evaluated for each treatment. All data were evaluated by two-way factorial analysis of variance (ANOVA) and Tukey's Post hoc test at $P < 0.05$ by SAS 9.1 software.

Results and discussion

Pepper cultivar Karpex inoculated with AM mixture exhibited higher root fresh and dry weight than the counterpart while conversely, in Karpia and Kaptur root fresh and dry weight in AM plants were reduced and remained unchanged, respectively in relation to non-inoculated plants (Figure 1). Noticeably, all varieties pretreated with AM showed higher shoot fresh weight than their counterparts, however, the increase in dry shoot weight was only observed in Karpex cultivar applied by AM, not Karpia and Kaptur.




Figure 1. Effect of AM inoculation on root fresh weight (A - upper left), root dry weight (B - upper right), shoot fresh weight (C - lower left) and shoot dry weight (D - lower right) of three pepper cultivars, Karpia, Karpex, Kaptur. Each bar presents mean \pm standard deviation

AM inoculated plants in all pepper cultivars showed the greater AM colonization rate in roots than non-inoculated controls (Figure 2). Interestingly, fruit yield in cultivars inoculated with AM improved significantly than non-inoculated plants, nevertheless, Karpex cv. has the similar fruit yield to their control. The results of inclined biomass production, fruit yield are accordance to numerous reports showing that AM inoculation enhanced root and shoot dry weight, fruit yield in pepper plants (Abdel Latef & Chaoxing, 2014; Abdel Latef, 2013; Boonlue et al., 2012; Tanwar et al., 2013). The most common explanation is that AM symbiosis enhance absorption of soil mineral nutrients, water through the AM hyphae network (Smith & Read, 2008), induced resistance or tolerance

to various stresses (Garmendia et al., 2004; Alejo-Iturvide et al., 2008; Mena-Violante et al., 2006; Kaya et al., 2009; Ruscitti et al., 2011). However, no beneficial effect from AM inoculation on pepper plants were also found in other reports (Russo, 2006; Russo & Perkins-Veazie, 2010).




Figure 2. Arbuscular mycorrhizal colonization rate (A) and fruit yield (B) of AM inoculated and control plants of three pepper cultivars, Karpia, Karpex, Kaptur. AM, Arbuscular mycorrhizal fungi; Each bar presents mean \pm standard deviation. Different letters denote significant differences among treatments according to Tukey's post hoc test ($P < 0.05$) among treatments.

Conclusions

This study showed that application of AM mixture generally enhanced the growth, fruit yield of pepper plants but not in all cultivars tested under field conditions.

Acknowledgement

Authors thank to Stipendium Hungaricum fellowship for supporting this study and to Institute of Horticultural Technology at Szent Istvan University.

References

- Abdel Latef, A.A.H., Chaoning, H. (2014). Does inoculation with *Glomus mosseae* improve salt tolerance in pepper plants? *J. Plant Growth Regul.* **33**:3. 644-653. DOI: <http://doi.org/10.1007/s00344-014-9414-4>
- Abdel Latef, A.A. (2013). Growth and some physiological activities of pepper (*Capsicum annum* L.) in response to cadmium stress and mycorrhizal symbiosis. *J. Agr. Sci. Tech.* **15**: 1437-1448.
- Alejo-Iturvide, F., Márquez-Lucio, M.A., Morales-Ramírez, I., Vázquez-Garciduenas, M.S., Olalde-Portugal, V. (2008). Mycorrhizal protection of chili plants challenged by *Phytophthora capsici*. *Eur. J. Plant Pathol.* **120**: 13–20. DOI: <https://doi.org/10.1007/s10658-007-9188-7>
- Boonlue, S., Surapat, W., Pukahuta, C., Suwanarit, P., Suwanarit, A., Morinaga, T. (2012). Diversity and efficiency of arbuscular mycorrhizal fungi in soils from organic chili (*Capsicum frutescens*) farms. *Mycoscience.* **53**:1. 10–16. DOI: <https://doi.org/10.1007/s10267-011-0131-6>
- Birhane, E., Sterck, F.J., Fetene, M., Bongers, F., Kuyper, T.W. (2012). Arbuscular mycorrhizal fungi enhance photosynthesis, water use efficiency, and growth of frankincense seedlings under pulsed water availability conditions. *Oecologia.* **169**: 895–904. DOI: <https://doi.org/10.1007/s00442-012-2258-3>
- Estrada-Luna, A.A., Davies, F.T.Jr. (2003). Arbuscular mycorrhizal fungi influence water relations, gas exchange, abscisic acid and growth of micropropagated chile ancho pepper (*Capsicum annum*) plantlets during acclimatization and post-acclimatization. *J. Plant Physiol.* **160**: 1073–1083. DOI: <https://doi.org/10.1078/0176-1617-00989>
- Estrada-Luna, A.A., Davies, F.T.Jr., Egilla, J.N. (2000). Mycorrhizal enhancement of the physiology and growth of micropropagated chile ancho pepper (*Capsicum annum* L. cv. San Luis) plantlets during acclimatization and post-acclimatization. *Hortscience.* **35**: 426.

- Franco, A.D., Carrillo, M.A., Chairez, F.O., Cabrera, O.G. (2013). Plant nutrition and fruit quality of pepper associated with arbuscular mycorrhizal in greenhouse. *Rev. Mexicana Cienc. Agrícola*. **4**: 315–321.
- Gajanayake, B., Trader, B.W., Reddy, K.R., Harkess, R.L. (2011). Screening ornamental pepper cultivars for temperature tolerance using pollen and physiological parameters. *HortScience*. **46**: 878–884.
- Garmendia, I., Goicoechea, N., Aguirreolea, J. (2004). Effectiveness of three *Glomus* species in protecting pepper (*Capsicum annuum* L.) against verticillium wilt. *Biol. Control*. **31**: 296–305. DOI: <http://dx.doi.org/10.1016/j.biocontrol.2004.04.015>
- Giovannetti, M., Mosse, B. (1980). An evaluation of techniques for measuring vesicular-arbuscular mycorrhizal infection in roots. *New Phytol.* **84**: 489–500. DOI: <https://doi.org/10.1111/j.1469-8137.1980.tb04556.x>
- Kaya, C., Ashraf, M., Sonmez, O., Aydemir, S., Tuna, A.L., Cullu, M.A. (2009). The influence of arbuscular mycorrhizal colonisation on key growth parameters and fruit yield of pepper plants grown at high salinity. *Sci. Hort.* **121**: 1–6. DOI: <https://doi.org/10.1016/j.scienta.2009.01.001>
- Mena-Violante, H.G., Ocampo-Jiménez, O., Dendooven, L., Martínez-Soto, G., González-Castaneda, J., Davies, F.T.Jr., Olalde-Portugal, V. (2006). Arbuscular mycorrhizal fungi enhance fruit growth and quality of chile ancho (*Capsicum annuum* L. cv San Luis) plants exposed to drought. *Mycorrhiza*. **16**: 261–267. DOI: <https://doi.org/10.1007/s00572-006-0043-z>
- Ortas, I., Sari, N., Akpınar, Ç., Yetisir, H. (2011). Screening mycorrhiza species for plant growth, P and Zn uptake in pepper seedling grown under greenhouse conditions. *Sci. Hort.* **128**: 92–98. DOI: <https://doi.org/10.1016/j.scienta.2010.12.014>
- Ruscitti, M., Arango, M., Ronco, M., Beltrano, J. (2011). Inoculation with mycorrhizal fungi modifies proline metabolism and increases chromium tolerance in pepper plants (*Capsicum annuum* L.). *Braz. J. Plant Physiol.* **23**: 15–25. DOI: <https://doi.org/10.1590/s1677-04202011000100004>
- Russo, V.M., Perkins-Veazie, P. (2010). Yield and nutrient content of bell pepper pods from plants developed from seedlings inoculated, or not, with microorganisms. *HortScience*. **45**:3. 352–358.
- Russo, V.M. (2006). Biological amendment, fertilizer rate, and irrigation frequency for organic bell pepper transplant production. *HortScience*. **41**:6. 1402–1407.
- Smith, S.E., Read, D.J. (2008). *Mycorrhizal symbiosis*. 3rd edn. Academic Press
- Tanwar, A., Aggarwal, A., Kadian, N., Gupta, A. (2013). Arbuscular mycorrhizal inoculation and super phosphate application influence plant growth and yield of *Capsicum annuum*. *J. Soil Sci. Pl. Nutrit.* **13**:1. 55–66. DOI: <https://doi.org/10.4067/S0718-95162013005000006>
- Vierheilig, H., Coughlan, A.P., Wyss, U., Piché, Y., (1998). Ink and vinegar, a simple staining technique for arbuscular-mycorrhizal fungi. *Appl. Environm. Microbiol.* **64**:12. 5004–5007.