


Volume 5, Number 2 (2018)

Columella

Journal of Agricultural and Environmental Sciences



Szent István University Press
Gödöllő

COLUMELLA

Journal of Agricultural and Environmental Sciences

This peer reviewed journal of the Faculty of Agricultural and Environmental Sciences of the Szent István University, Gödöllő, Hungary publishes papers in English language.

Technical assistance is provided by the respective Scientific Committees of the Hungarian Academy of Sciences.

The journal is published in yearly volumes of two issues annually.

Editor-in-chief:
Béla URBÁNYI

Edited by:
Zoltán KENDE

Responsible publisher:
Mihály LAJOS

Editorial correspondence:
Faculty of Agricultural and Environmental Sciences of the Szent István University

Páter Károly utca 1.
H-2100 Gödöllő, Hungary
Phone: (+36 28) 522067
Fax: (+36 28) 410-804
web: www.columella.mkk.szie.hu
E-mail: columella@mkk.szie.hu

HU ISSN 2064-7816 (print)
HU ISSN 2064-9479 (online)
DOI: 10.18380/

Printed in Hungary, Gödöllő
Printed by Szent István Egyetemi Kiadó Nonprofit Kft. (Szent István University Press)
HU-2100 Gödöllő, Páter Károly utca 1.

Columella

Journal of Agricultural and Environmental Sciences

Volume 5, Number 2 (2018)

HU ISSN 2064-7816 (print)
HU ISSN 2064-9479 (online)
DOI: 10.18380/

Editor-in-chief
Béla URBÁNYI

Guest editor
Márton JOLÁNKAI

Edited by
Zoltán KENDE

© 2018 Szent István University
Faculty of Agricultural and Environmental Sciences, Gödöllő,
HUNGARY

Authors of current volume of Columella

Elizabeta ANĐELIĆ
Manda ANTUNOVIĆ
Károly BARTA
Csaba BOZÁN
Nguyen Hong DUC
Dario ILJKIĆ
Márton JOLÁNKAI
Gergő KOVÁCS
Ágnes KUN
Anett MÉSZÁROSNÉ PÓSS

Géza NAGY
Mária ONCSIK BÍRÓNÉ
Katalin POSTA
Péter RADÁCSI
Kálmán RAJKAI
Miro STOŠIĆ
Ferenc TÓTH
Franciska TÓTHÉ BOGDÁNYI
Ivana VARGA
Éva ZÁMBORI-NÉMETH
Angéla ZANKER

Reviewers of current volume of Columella

Franco MAGURNO
József CSAJBÓK
Katalin POSTA
Katalin Mária KASSAI
Endre DOBOS

Péter NAGY
Lajos BLASKÓ
Péter Gergő KOVÁCS
Tamás KISMÁNYOKY
Mirha Đikić

Scientific board of Columella

Manuel ACOSTA (CZ)
Adalbert BALOG (RO)
Zoltán BEDŐ (H)
János BERÉNYI (RS)
Sándor CSÁNYI (H)
Dénes DUDITS (H)
Werner EUGSTER (CH)
György FÜLEKY (H)
Lajos HELYES (H)
László HESZKY (H)
István HOLLÓ (H)
László HORNOK (H)
Márton JOLÁNKAI (H)
Tamás KISMÁNYOKY (H)
Vlado KOVAČEVIĆ (HR)
Jan KUČTIK (CZ)
Magdalena LACKO-BARTOŠOVÁ (SK)
Éva LEHOCZKY (H)
Miklós MÉZES (H)

Erika MICHÉLI (H)
László MIKLÓS (SK)
Péter István NAGY (H)
János NAGY (H)
Zoltán NAGY (H)
Miklos NEMÉNYI (H)
Alberto PARDOSSI (IT)
János PAUK (H)
Károly PENKSZA (H)
Péter PEPÓ (H)
Katalin POSTA (H)
Péter PÓTI (H)
Erich M. PÖTSCH (A)
Rezső SCHMIDT (H)
Peter SURAI (UK)
János TÓZSÉR (H)
Béla URBÁNYI (H)
Ottó VEISZ (H)
György VÁRALLYAY (H)

Table of Contents

Editorial

Márton JOLÁNKAI..... 4

Inoculation with *Septoglomus constrictum* improves tolerance to heat shock in tomato plants

Nguyen Hong DUC – Katalin POSTA..... 7

***In vitro* activity of the cinnamon essential oil against the plant pathogen *Septoria melissae* desm.**

Gergő KOVÁCS – Éva ZÁMBORI-NÉMETH – Péter RADÁCSI – Géza NAGY..... 15

Preliminary results of a pot experiment with the combined effects of a terrestrial isopod species (*Porcellionides pruinosus*, Brandt 1833) and organic mulching on tomato

Anett Mészárosné PÓSS – Angéla ZANKER – Franciska TÓTHNÉ BOGDÁNYI – Ferenc TÓTH..... 21

Yield components of winter oilseed rape regard to plant population

Elizabeta ANĐELIĆ – Manda ANTUNOVIĆ – Miro STOŠIĆ – Dario ILJKIĆ – Ivana VARGA..... 33

Calculation nitrogen and sodium budget from lysimeter-grown short-rotation willow coppice experiment

Ágnes KUN – Csaba BOZÁN – Mária ONCSIK BÍRÓNÉ – Károly BARTA..... 43

OBITUARY


György FÜLEKY (1945-2018) prominent agrochemist passed away

Kálmán RAJKAI..... 58

Editorial

How I edited an agricultural paper once

The title was adopted from the marvellous short story of Mark Twain, published in 1870 by The Galaxy: an illustrated monthly magazine edited in New York. The humorous story summarizes the adventures of a journalist who is appointed by an agricultural paper temporarily. The regular editor of the paper was going off for a holiday, and someone accepted the terms he offered, and took his place. He tried to handle the journal in accordance with the general patterns of newspapers. Namely – it does not matter what you write, as long as it is attractive to the readers. The more astonishing the information is the better is the reputation of the periodical. In this belief our man wrote various crazy articles about turnips that should never be pulled from the soil but harvested from trees, or how to hatch the eggs of the guano bird, and why not produce pumpkins instead of gooseberries etc. The readers of the paper soon started to revolt, the poor regular editor had to return from his leave, and our man was fired. At the end of the story there is a real “Mark-Twainian” conclusion. Specific papers in general and agricultural papers in particular should avoid employing journalists having no sufficient information on the matter they write about. This should be left for political parties since they welcome anybody with the philosophy “that the less a man knows the bigger noise he makes, and the higher the salary he commands”.



Front page of an original GALAXY magazine from the Dave Thomson collection.

Márton Jolánkai

So, it isn't an easy job neither to produce nor to edit a scientific paper. Columella - Journal of Agricultural and Environmental Sciences has reached a milestone. We are five years old now, and the present issue is just the 10th in our history. Of course it is not (or not *yet*) an anniversary. We don't have to be proud and also we cannot be satisfied with the results of this brief period. However it would be worth to see some of the records of the paper. The foundation of the journal was in 2014. We are completing the fifth volume by now. Ten regular issues and one conference proceedings book have been edited, so far containing 74 full papers and 50 conference papers. These have been selected from among 145 articles. The rest of them were discarded during the reviewing processes. We believe in the wise opinion of a onetime professor of ours according to whom "the value of a paper is determined by the power of the reviewer". During the past five years altogether 105 international reviewers contributed to the success of our work. The magnitude of papers published is 944 pages. The editorial process required some 1500 working hours. All papers are open access published and so they are available on the internet having DOI numbers and the journal is ISSN registered with both hard copies and electronic issues. The journal is indexed and stored by a wide range of repositories and it is a scientific journal approved by the Hungarian Academy of Sciences.

Finally we have to return to the initial question again; what is the mission of a scientific paper nowadays? I believe it is the support of science in general, and the dissemination of scientific results in a controlled, broad and open access way to the public. Our journal, Columella provides a forum for scientific publications in the field of agricultural and environmental sciences. The journal of one of the most ambitious agricultural faculties of Hungary has a task to disseminate novel research results, in favour of creating a better world.

The editors would say thanks to the authors of the present issue, and also welcome the readers who may read, use and broadcast the scientific information compiled. We also do hope that they may become future authors as well.

Márton Jolánkai

References

- Berkove L. (2004): The best short stories of Mark Twain. Penguin Books. Random House. Canada.
- Schmidt B. (2018): Mark Twain Quotations, Newspaper Collections, & Related Resources.
<http://www.twainquotes.com/Galaxy/gindex.html>
- Twain M. (1997): The L 1,000,000 bank-note. Three stories. A mysterious visit. How I edited an agricultural paper / Az egymillió fontos bankjegy. Három elbeszélés. Titokzatos látogatás. Hogyan szerkesztettem mezőgazdasági lapot; vál., jegyz. Heltai Pál, ford. Szász Imre; Corvina, Bp. (Kétnyelvű olvasmányok)

Inoculation with *Septoglomus constrictum* improves tolerance to heat shock in tomato plants

Nguyen Hong DUC – Katalin POSTA

Institute of Genetics, Microbiology and Biotechnology, Department of Microbiology and Environmental Toxicology, Szent István University, Páter Károly Street 1, 2100 Gödöllő, Hungary.
E-mail: hongduc.real@gmail.com

Abstract: Arbuscular mycorrhizal fungi (AMF) are symbiotic soil fungi colonizing roots of about 80% of vascular plants. This symbiosis enhances the growth and survival of numerous plant species including vegetables; moreover, it offers some other benefits for the plants. This work aimed to study the impact of AMF on tomato plant tolerance to heat shock. Tomato (*Solanum lycopersicum* L.) plants inoculated or not with *Septoglomus constrictum* were placed in a commercial potting media at 26/22°C with 16/8h photoperiod for six weeks, then exposed to normal (26°C for 6h) or high temperature (42°C for 6h). Arbuscular mycorrhizal (AM) colonization rate, level of lipid peroxidation (malondialdehyde – MDA), hydrogen peroxide (H₂O₂) accumulation and antioxidative enzymes in roots and leaves were measured after the stress application. AM colonization rate of mycorrhizal plants was 73% under non-stress conditions and 68% under heat shock conditions while no mycorrhizal colonization found in non-AM treatments. MDA and H₂O₂ content substantially increased in leaves of all plants after exposure to the heat shock. Leaf and root peroxidase (POD), leaf catalase (CAT) and root superoxide dismutase (SOD) activities of mycorrhizal plants were enhanced compared to those in non-AM ones while the activity of leaf SOD and root CAT in mycorrhizal plants remained unchanged. Furthermore, there were significant decreases in MDA and H₂O₂ content in leaves of inoculated plants compared with non-AM ones under heat shock conditions. Our results indicate that AM inoculation can increase stress tolerance against heat shock by enhancing the activity of antioxidative enzymes. Further research is required to understand the mechanisms that contribute to heat tolerance.

Abbreviations: AM, arbuscular mycorrhizal; AMF, arbuscular mycorrhizal fungi; CAT, catalase; FW, fresh weight; MDA, malondialdehyde; POD, peroxidase; ROS, reactive oxygen species; SOD, superoxide dismutase.

Keywords: arbuscular mycorrhizal fungi, *Septoglomus constrictum*, heat shock stress, stress tolerance, tomato plants

Received 12 March 2018, Revised 19 November 2018, Accepted 14 December 2018

Introduction

Due to the effects of global warming, heat stress has become the major challenge for crop production on earth. Heat stress causes anatomical, morphological, physiological, biochemical, and genetic responses *in planta* (Camejo et al. 2005; Chen et al. 2012; Min et al. 2014), diminishing crop yield and quality. When plants are subjected to very high temperatures, severe injuries in the cell, even death, may take place within minutes. These could be ascribed to a devastating collapse of the cellular system (Schoffl et al. 1999). Nevertheless, such cell damages and death may take place merely after plants are exposed to moderately high temperatures in the long term (Wahid et al. 2007). As direct consequences of heat stress, proteins are denatured and aggregated while the fluidity of membrane lipids is elevated in plants. Indirectly

or slowly, high-temperature stress causes enzyme inactivation in mitochondria and chloroplast, protein synthesis prohibition, degradation of proteins and membrane integrity loss (Howarth 2005). These damages eventually lead to starvation, plant growth reduction, decreased ion flux, generation of toxic compounds as well as reactive oxygen species (ROS) (Howarth 2005).

ROS consist of peroxides, superoxide, hydroxyl radical, and singlet oxygen. ROS overproduction may result in cell death as a consequence of oxidative stress, such as peroxidation of membrane lipid, causing oxidative damage to nucleic acids (Tanou et al. 2009). Peroxidation of membrane lipid is detected by measuring malondialdehyde (MDA) which is a widely used marker of oxidative lipid injury caused by environmental stresses (Kong et al. 2016). To reduce oxidative damage under temperature

stresses, plants have evolved different antioxidative strategies to detoxify harmful ROS components by non-enzymatic and enzymatic antioxidant defence systems where peroxidase (POD), catalase (CAT), superoxide dismutase (SOD) are main enzymatic ROS scavengers (Wu et al., 2014). SOD is the first defense against ROS (Alscher et al. 2002) due to its ability to catalyze the dismutation of $O_2^{\cdot-}$ to H_2O_2 (Wu et al. 2014). Subsequently, CAT and other scavenging enzymes detoxicate H_2O_2 to H_2O and O_2 (Apel and Hirt 2004). POD can also generate and detoxify H_2O_2 in the first and next phase, respectively (Siegel 2003).

Studies on plants under temperature stresses are extensively carried out to develop strategies to deal with the adverse effects of heat stress on crop productivity through breeding heat-tolerant varieties, suitable crop shifts and cultivation practices. Most of them are costly and time-consuming whereas noticeably, using beneficial microbes has been proved as a potential solution to improve plant tolerance to various abiotic and biotic stresses. Arbuscular mycorrhizal fungi (AMF), one of the common soil microbes, can form the symbiotic association with roots of 80% of terrestrial plant species. The application of AMF enhances not only nutrient and water uptake but plant tolerance to abiotic stress (Birhane et al. 2012).

Cabral and coworkers (2016) showed that inoculation of an AM mixture including *Rhizophagus irregularis* BEG140, *Rhizophagus irregularis*, *Funneliformis mosseae* BEG95, *Funneliformis geosporum*, *Claroideoglossum claroideum* in wheat plants mitigated adverse effects of temperature stress at 35°C (day) and 25°C (night) for seven days. In maize, inoculation of *Claroideoglossum etunicatum* reduced relative membrane permeability and MDA concentration in roots and leaves of plants while it increased soluble sugar and proline content in roots but lowered leaf proline content, relative to non-AM plants as exposure to 35°C and 40°C for one week (Zhu et al. 2010). Improved net photosynthetic rate (P_n), stomatal conductance (g_s), transpiration rate (E), the maximum quantum efficiency of photosystem II (Fv/Fm) together with higher chlorophyll contents in leaves of mycorrhizal maize plants under such heat stress conditions were observed (Zhu et al. 2011). Enhanced

biomass of mycorrhizal plants under heat stress conditions was also detected (Zhu et al. 2010; Maya and Matsubara 2013; Matsubara et al. 2014; Maya et al. 2014). AM application triggered the higher activity of SOD, ascorbate peroxidase in whole plants (roots, tubers, and leaves) (Matsubara et al. 2014; Maya and Matsubara 2013) and resulted in heightened leaf ascorbic acid and polyphenol in cyclamen plants after exposure to heat stress (Maya and Matsubara 2013). Similarly, POD, SOD, CAT activities were elevated in roots and leaves in mycorrhizal plants under temperature stresses (Zhu et al. 2010). It is worth mentioning that the level and duration of stress (acute versus chronic) significantly affect plant responses (Tattersall et al. 2007; Pinheiro and Chaves 2011). Previous studies on mycorrhizal plants focused on the chronic heat stresses which usually lasted for one week. Therefore, mechanisms underlying the effect of AM inoculation on the ROS metabolism and antioxidative enzymes of host plants under heat shock representing acute stress remain unknown.

Tomato (*Solanum lycopersicum* L.) is a main vegetable crop in the world, widely cultivated optimally in agricultural areas with temperatures between 20°C and 30°C. Tomato is a primary dietary component in different countries because it contains a rich source of vitamins, antioxidant compounds, minerals, sugars, providing significant nutritional value for the human. Nevertheless, tomato productivity is substantially decreased by abiotic stresses (Schwarz et al. 2010).

The objective of this study was, hence, to contribute to the understanding of the effect of AM inoculation on ROS metabolism and the antioxidative activity in tomato plants under heat shock (42°C for 6h). *Septoglossum constrictum*, distributed around the world (Opik et al. 2010), was chosen as the fungal inoculant in our study. A degree of lipid peroxidation (estimated by MDA) and H_2O_2 accumulation, together with POD, SOD, CAT activity in leaves and roots of mycorrhizal tomato plants and non-AM tomato plants under heat shock were examined.

Material and methods

Tomato (*Solanum lycopersicum*) seeds, cultivar MoneyMaker, were soaked in 2.5% sodium hypochlorite for 20 minutes, then washed with

distilled water five times and placed on wet papers in Petri dishes for germination for three days at room temperature. Germinated seeds were put in plastic pots (0.5-lit volume) with 0.5 kg of the sterile mixture of sand and soil (4:1, v/v). The loamy soil with pH 7.1, 1.61% organic matter, N 15.6 mg kg⁻¹, available P 36 mg kg⁻¹, available K 60 mg kg⁻¹ (Duc et al., 2017) was used.

The experiment consisted of 12 plants without AM inoculation and 12 plants inoculated by *Septoglomus constrictum* (formerly *Glomus constrictum* Trappe.). The AM inoculum was cultured in the sterile sand with *Zea mays* as host plants for four months. Thirty grams of the AM inoculum (27 spores g⁻¹) were utilised for each pot in AM treatment while plants in non-AM treatment were added the same amount of autoclaved inoculum and 3 ml aliquot of a filtrate (< 20 µm) of the AM inoculum in order to provide a general microbial population free AMF propagules. Pots were put randomly in a growth chamber (EKOCHIL 1500), and the pot positions were changed weekly. Growing conditions, 26/20°C with 16/8 hours photoperiod, light intensity 600 µmol m⁻² s⁻¹ and 60% humidity were applied. Pots were watered twice and fertilized with Long Ashton nutrient solution (Hewitt, 1966) with low phosphorus level (3.2 µM Na₂HPO₄·12H₂O) once a week. After six weeks of plant growth, heat shock was carried out by transferring six non-AM plants and six AM plants to 42°C for 6h (Zhou et al. 2014) whereas six plants without AMF and six mycorrhizal plants were kept under non-stress conditions. Then all leaf and root samples were collected for further analysis.

Assessment of arbuscular mycorrhizal colonization

Root samples were washed by tap water and cleaned before staining according to Vierheilig et al. (1998). AM colonization was examined by visual inspection of fungal structures under a stereomicroscope at x 100 magnification. AM colonization rate was determined by the gridline intersect method (Giovannetti and Mosse 1980).

Measurement of hydrogen peroxide accumulation and oxidative damage to lipids

The H₂O₂ content was determined by the method

of Alexieva et al. (2001). Shortly, leaf samples (500 mg) were ground with a cold 0.1% (w/v) trichloroacetic acid (TCA) (5 ml), subsequently, centrifuged at 12,000×g at 4°C for 15 min. The reaction mixture consisted of 100 mM potassium phosphate buffer (pH 7.0, 0.5 ml), 1 M KI (1 ml) and the leaf extract supernatant (0.5 ml). The reaction occurred in the dark for 1h. Its absorbance was recorded at 390 nm.

The leaf lipid peroxidation was estimated by the method of Heath and Packer (1969). In detail, leaf samples (200 mg) were ground in 0.1% TCA (5 ml), then centrifuged at 10,000×g for 5 min. A 1 ml of leaf supernatant was mixed with 20% TCA (4 ml) containing 0.5% 2-thiobarbituric acid (TBA). Then the mixture was heated at 95°C for 15 min and immediately cooled. Absorbances of the mixture at 532 nm and 600 nm were recorded for MDA estimation. The content of MDA was estimated using an extinction coefficient of 155 mM⁻¹ cm⁻¹.

Measurement of antioxidant enzymatic activities

Leaf and root samples (500 mg) were frozen in liquid nitrogen and ground with 50 mM Tris-HCl buffer pH 7.8 (3 ml) containing 1 mM Na₂EDTA and 7.5% (w/v) polyvinylpyrrolidone K25. Then, crude extracts were centrifuged at 10,000 x g for 20 minutes at 4°C. The supernatants were collected to examine enzyme activities by U-2900 UV-VIS spectrophotometer (Hitachi). Soluble protein contents were estimated according to the method of Bradford (1976).

The activity of peroxidase (POD, EC 1.11.1.7) was tested according to Rathmell and Sequeira (1974). A 2.2 ml reaction mixture included 10 µl plant extract, 12mM H₂O₂ (100 µl), 50 mM Guaiacol (100 µl) and 0.1 M sodium phosphate buffer (pH 6.0). Changes in the absorption at 436 nm for 5 minutes were recorded. The activity of POD was presented by the changes in absorbance per mg protein per minute.

The activity of superoxide dismutase (SOD, EC 1.15.1.1) was examined by the method of Beyer and Fridovich (1987). A 2 ml 50 mM phosphate buffer (pH 7.8) containing 55 µM NBT, 2 mM EDTA, 9.9 mM L-methionine and 0.025% Triton X-100 and leaf extract (20 µl), 1 mM riboflavin (20 µl) was used for the reaction

mixture. Absorbance changes of the reaction at 560 nm were recorded.

The activity of catalase (CAT, EC 1.11.1.6) was assayed by the method described by Aebi and Lester (1984). The reaction mixture (3 ml) consisted of 10 mM H₂O₂, 50 mM potassium phosphate buffer (pH 7.0) and the enzyme extract. Absorbance decreases of the reaction at 240 nm were recorded. CAT activity was presented as absorbance changes per mg protein per minute.

Statistical analysis

All data were tested by two-way factorial analysis of variance (ANOVA) using SAS 9.1 (SAS Institute, Cary, North Carolina). The means were compared at the 5% level by Duncan posthoc test.

Results

No dead plants were observed in our experiment after heat shock treatment. AM colonization rate of plants colonized by *S. constrictum* was 73% under non-stress conditions and 68% under heat shock conditions whereas no AM colonization was observed in plants without mycorrhiza (Figure 1A). Also, heat shock (42°C in 6 hours) did not change the AM colonization significantly

although a slight decrease occurred. Under non-stress conditions, both AM and non-AM plants had similar MDA values, however; when plants were subjected to heat shock, MDA levels considerably increased in non-AM plants (by 42%) but not in AM plants (Figure 1B).

In addition, AM plants also showed a significant lower MDA (17% lower) than non-AM plants under heat shock conditions. Similarly, there was no significant difference in H₂O₂ level in AM and non-AM plants under non-stress conditions (Figure 1C). Heat shock increased H₂O₂ level by six-fold in non-AM plants and by over three-fold in AM plants. H₂O₂ accumulation reduced by 40% in AM plants, as compared to the corresponding in non-AM plants under heat shock.

As shown in Figure 2A, POD activity in leaves of mycorrhizal plants was not significantly different from that of non-AM plants under non-stress conditions. Nonetheless, exposure to heat shock substantially increased (61-76%) leaf POD level in plants although AM symbiosis induced a considerable higher (9%) POD activity than plants without AM inoculation. Notably, heat shock substantially heightened (19-26%) SOD activity of leaves in plants but there was




Figure 1. Mycorrhizal colonization rate (A), malondialdehyde (MDA) (B) and H₂O₂ (C) accumulation in leaves of non-AM plants (AM-) and plants colonized by *Septoglomus constrictum* (AM+) exposed to non-stress, heat-shock conditions. Bars present mean ± standard deviation (n = 4). Different letters present significant differences among treatments according to Duncan posthoc test (P < 0.05).

no significant difference in leaf SOD activity between AM and non-AM plants (Figure 2B). Likewise, leaf CAT activity of colonized plants was considerably improved (increased by 18%), in comparison to that of uncolonized plants under heat shock despite no significant differences in this enzyme between both AM and non-AM plants under non-stress conditions (Figure 2C).

In roots, POD activity had the same pattern as in leaves (Figure 2D). Root POD activity in AM and non-AM plants under non-stress conditions did not differ considerably, nevertheless; under heat shock conditions AM plants expressed an improved POD activity (increased by 22%), relative to non-AM plants. In contrast to leaf SOD activity, this enzyme was enhanced significantly

(increased by 87%) in AM plants, as compared to non-AM plants under heat stress conditions although no substantial differences between AM and non-AM plants under normal conditions were recorded (Figure 2E). Regarding root CAT activity, no considerable differences between AM and non-AM plants were found under non-stress as well as heat shock conditions although the level of CAT was elevated by 75-102% in plants subjected to the heat stress (Figure 2F).

Discussion

AMF can enhance the host tolerance to temperature stresses in maize (Zhu et al. 2010; 2011), in cyclamen (Maya and Matsubara 2013), in citrus (Wu, 2011), in tomato (Abdel Latif and




Figure 2. Peroxidase (POD) (A), superoxide dismutase (SOD) (B), catalase (CAT) (C) activity in leaves and POD (D), SOD (E) and CAT (F) in roots of non-AM plants and plants colonized by *Septoglomus constrictum* exposed to non-stress, heat-shock conditions. Bars present mean \pm standard deviation ($n = 4$). Different letters present significant differences among treatments according to Duncan posthoc test ($P < 0.05$).

Chaoxing 2011), however; little information on how AM symbiosis responds to heat shock was explored. In this study, the effect of AM inoculation with *Septoglomus constrictum*, an uncommon AM isolate in previous studies under heat stress, on plant tolerance to heat shock in tomato plants was investigated.

Temperature stresses can negatively impact on the growth and development of AM symbiosis (Zhu et al. 2010; 2011). Earlier studies showed that the development of AMF was inhibited by low temperatures (Zhu et al. 2010) whereas high temperatures influenced detrimentally the activity of AMF (Martin and Stutz 2004) and negative or neutral AM colonization (Compant et al. 2010). In the present study, heat shock had no significant effect on mycorrhizal colonization of tomato plants, which might be due to the length of heat stress applied. This result is in line with the observations in previous studies under heat stress in cyclamen (Maya and Matsubara, 2013), in maize (Zhu et al. 2010).

Extreme temperatures cause unbalanced cellular homeostasis, resulting in overproduction of ROS, membrane lipid peroxidation, and damage plant cells. In this study, we observed that there was an elevated MDA and H₂O₂ accumulation in leaves of tomato plants exposed to heat shock, nevertheless, substantially lower MDA and H₂O₂ contents were found in colonized plants than in plants without AM inoculation, suggesting that the presence of the AMF could alleviate the oxidative stress and peroxidation. The present results concur with observations in tomato plants (Abdel Latef and Chaoxing 2011). The authors illustrated that inoculation of *Funneliformis mosseae* decreased MDA in plants as compared to non-AM plants under cold stress (8°C) for 1 week. Zhu et al. (2010) also showed that maize plants colonized by *Claroideoglossum etunicatum* reduced substantially MDA in roots and leaves when plants were subjected to various temperature stresses (5, 15, 35, 40°C) for one week.

Plant tolerance to high-temperature stress has been linked with an increase in activities of antioxidant

enzymes (Sairam et al. 2000). Under heat shock, AM plants showed an enhanced POD, CAT activity in leaves, relative to non-AM plants, which is consistent with lower MDA and H₂O₂ contents observed in leaves of AM plants in our study. In roots, POD and SOD activity in mycorrhizal plants were significantly higher than those in uncolonized plants. These findings suggest that AMF treatment improved the effectiveness of antioxidant systems to protect the host plant against oxidation damage under heat shock conditions. Our results are in accordance with earlier mycorrhiza studies in tomato plants under cold stress (Abdel Latef and Chaoxing 2011), in maize under high-temperature stresses (Zhu et al., 2010). However, leaf SOD and root CAT were not improved in AM plants in our study, which was inconsistent with observations of Abdel Latef and Chaoxing (2011), Zhu et al. (2010) and Maya and Matsubara (2013). The findings may indicate that different AMF isolate could induce differently antioxidant systems in the host plant under heat stress conditions. Similarly, Cekic et al. (2012) reported that pepper plants colonized by *Rhizophagus irregularis* enhanced substantially CAT activity in leaves whereas the leaf SOD level remained unchanged in both AM and non-AM plants when plants exposed to 1 mM NaCl stress.

In conclusion, our results point out that inoculation of AMF, *Septoglomus constrictum*, could enhance tomato plant tolerance against heat shock by decreasing oxidative stress (reduced H₂O₂ and MDA content) and increasing activities of main ROS scavengers such as leaf and root POD, root SOD and leaf CAT. Nonetheless, further studies are necessary to elucidate mechanisms by which AMF influence antioxidant production, proline, photosynthesis, respiration, and water status in plants to better understand their benefits under heat stress for agricultural application.

Acknowledgement

Authors thank Stipendium Hungaricum fellowship for supporting this study.

References

- Abdel Latef, A.A.H., Chaoxing, H. (2011): Effect of arbuscular mycorrhizal fungi on growth, mineral nutrition, antioxidant enzymes activity and fruit yield of tomato grown under salinity stress. *Scientia Horticulturae*. **127**: 3. 228–233. DOI: <http://doi.org/10.1016/j.scienta.2010.09.020>

- Aebi, H., Lester, P. (1984): Catalase in vitro. *Meth Enzymol.* **105**: 121–126.
- Alexieva, V., Sergiev, I., Mapelli, S., Karanov, E. (2001): The effect of drought and ultraviolet radiation on growth and stress markers in pea and wheat. *Plant, Cell & Environment.* **24**: 1337-134. DOI: <https://doi.org/10.1046/j.1365-3040.2001.00778.x>
- Alscher, R.G., Erturk, N., Heath, L.S. (2002): Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. *Journal of Experimental Botany.* **53**: 372. 1331–1341. <http://doi.org/10.1093/jexbot/53.372.1331>
- Apel, K., Hirt, H. (2004): Reactive oxygen species: Metabolism, oxidative Stress, and signal transduction. *Annual Review of Plant Biology.* **55**: 1. 373–399. DOI: <https://doi.org/10.1146/annurev.arplant.55.031903.141701>
- Beyer, W.F., Fridovich, I. (1987): Assaying for superoxide dismutase activity: some large consequences of minor changes in conditions. *Anal Biochem.* **161**: 2. 559–566. DOI: [https://doi.org/10.1016/0003-2697\(87\)90489-1](https://doi.org/10.1016/0003-2697(87)90489-1)
- Birhane, E., Sterck, F.J., Fetene, M., Bongers, F., Kuyper, T.W. (2012): Arbuscular mycorrhizal fungi enhance photosynthesis, water use efficiency, and growth of frankincense seedlings under pulsed water availability conditions. *Oecologia.* **169**: 4. 895–904. DOI: <https://doi.org/10.1007/s00442-012-2258-3>
- Bradford, M.M. (1976): A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**: 1-2. 248-254. DOI: [https://doi.org/10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3)
- Cabral, C., Ravnskov, S., Tringovska, I., Wollenweber, B. (2016): Arbuscular mycorrhizal fungi modify nutrient allocation and composition in wheat (*Triticum aestivum* L.) subjected to heat-stress. *Plant and Soil.* **408**: 1-2. 385–399. DOI: <http://doi.org/10.1007/s11104-016-2942-x>
- Camejo, D., Rodríguez, P., Morales, M.A., Dell'Amico, J.M., Torrecillas, A., Alarcón, J.J. (2005): High temperature effects on photosynthetic activity of two tomato cultivars with different heat susceptibility. *Journal of Plant Physiology.* **162**: 3. 281–289. DOI: <http://doi.org/10.1016/j.jplph.2004.07.014>
- Cekic, F.O., Unyayar, S., Ortas, I. (2012): Effects of arbuscular mycorrhizal inoculation on biochemical parameters in *Capsicum annuum* grown under long term salt stress. *Turkish Journal of Botany.* **36**: 1. 63–72. DOI: <http://doi.org/10.3906/bot-1008-32>
- Chen, L., Ren, Y., Zhang, Y., Xu, J., Sun, F., Zhang, Z., Wang, Y. (2012): Genome-wide identification and expression analysis of heat-responsive and novel microRNAs in *Populus tomentosa*. *Gene.* **504**: 2. 160–165. DOI: <https://doi.org/10.1016/j.gene.2012.05.034>
- Compant, S., van der Heijden, M.G.A., Sessitsch, A. (2010): Climate change effects on beneficial plant-microorganism interactions. *FEMS Microbial Ecol* **73**: 2. 197–214. DOI: <https://doi.org/10.1111/j.1574-6941.2010.00900.x>
- Duc, N. H., Mayer, Z., Pék, Z., Helyes, L., Posta, K. (2017): Combined inoculation of arbuscular mycorrhizal fungi, *Pseudomonas fluorescens* and *Trichoderma* spp. for enhancing defense enzymes and yield of three pepper cultivars. *Applied Ecology and Environmental Research.* **15**: 3. 1815-1829. DOI: https://doi.org/10.15666/aeer/1503_18151829
- Giovanetti, M., Mosse, B. (1980): An evaluation of techniques for measuring vesicular-arbuscular mycorrhizal infection in roots. *New Phytology.* **84**: 3. 489-500. DOI: <https://doi.org/10.1111/j.1469-8137.1980.tb04556.x>
- Heath, R.L., Packer, L. (1969): Photoperoxidation in isolated chloroplast: I. Kinetics and stoichiometry of fatty acid peroxidation. *Arch. Biochem Biophys.* **125**: 1. 189-198. DOI: [https://doi.org/10.1016/0003-9861\(68\)90654-1](https://doi.org/10.1016/0003-9861(68)90654-1)
- Hewitt, E.J. (1966): Sand and water culture methods used in the study of plant nutrition. 2nd edn. London: Commonwealth Agricultural Bureau.
- Howarth, C.J. (2005): Genetic improvements of tolerance to high temperature. In: Ashraf, M., Harris, P.J.C. (Eds.). *Abiotic Stresses: Plant resistance through breeding and molecular approaches.* Howarth Press Inc., New York.
- Kong, W., Liu, F., Zhang, C., Zhang, J., Feng, H. (2016): Non-destructive determination of Malondialdehyde (MDA) distribution in oilseed rape leaves by laboratory scale NIR hyperspectral imaging. *Sci. Rep.* **6**: 35393. DOI: <https://doi.org/10.1038/srep35393>
- Martin, C.A., Stutz, J.C. (2004): Interactive effects of temperature and arbuscular mycorrhizal fungi on growth, P uptake and root respiration of *Capsicum annuum* L. *Mycorrhiza.* **14**: 4. 241–244. DOI: <https://doi.org/10.1007/s00572-003-0261-6>
- Matsubara, Y., Ishioka, C., Maya, M.A., Liu, J., Takami, Y. (2014): Bioregulation potential of arbuscular mycorrhizal fungi on heat stress and anthracnose tolerance in cyclamen. *Acta Horticulturae.* **1037**: 813–818. DOI: <https://doi.org/10.17660/actahortic.2014.1037.108>

- Maya, M.A., Matsubara, Y. (2013): Influence of arbuscular mycorrhiza on the growth and antioxidative activity in cyclamen under heat stress. *Mycorrhiza*. **23**: 5. 381-390. DOI: <https://doi.org/10.1007/s00572-013-0477-z>
- Maya, M.A., Ito, M., Matsubara, Y. (2014): Tolerance To Heat Stress and Anthracnose in Mycorrhizal Cyclamen. *Acta Hort.*, **1025**: 143–148. DOI: <http://doi.org/10.17660/ActaHortic.2014.1025.21>
- Min, L., Li, Y., Hu, Q., Zhu, L., Gao, W., Wu, Y., Ding, Y., Liu, S., Yang, X., Zhang, X. (2014): Sugar and auxin signalling pathways respond to high-temperature stress during anther development as revealed by transcript profiling analysis in cotton. *Plant Physiol.* **164**: 3. 1293–1308. DOI: <https://doi.org/10.1104/pp.113.232314>
- Opik, M., Vanatoa, A., Vanatoal, E., Moora, M., Davison, J., Kalwij, J. M., Reier, U., Zobel, M. (2010): The on-line database MaarjAM reveals global and ecosystemic distribution patterns in arbuscular mycorrhizal fungi (Glomeromycota). *New Phytol.* **188**: 223–241. DOI: <https://doi.org/10.1111/j.1469-8137.2010.03334.x>
- Pinheiro, C., Chaves, M.M. (2011): Photosynthesis and drought: can we make metabolic connections from available data? *J Exp Bot.* **62**: 3. 869-882. DOI: <https://doi.org/10.1093/jxb/erq340>
- Rathmell, W.G., Sequeira, L. (1974): Soluble peroxidase in fluid from the intercellular spaces of tobacco leaves. *Plant Physiology*. **53**: 2. 317-318. DOI: <https://doi.org/10.1104/pp.53.2.317>
- Sairam, R.K., Srivastava, G.C., Saxena, D.C. (2000): Increased antioxidant activity under elevated temperature: a mechanism of heat stress tolerance in wheat genotypes. *Biol Plant.* **43**: 2. 245–251. DOI: <https://doi.org/10.1023/A:1002756311146>
- Schoffl, F., Prandl, R., Reindl, A., (1999): Molecular responses to heat stress. In: Shinozaki, K., Yamaguchi-Shinozaki, K. (Eds.). *Molecular Responses to Cold, Drought, Heat and Salt Stress in Higher Plants*. R.G. Landes Co., Austin, Texas, pp. 81–98.
- Schwarz, D., Roupshael, Y., Colla, G., Venema, J.H. (2010): Grafting as a tool to improve tolerance of vegetables to abiotic stresses: thermal stress, water stress and organic pollutants. *Sci. Hortic.* **127**: 2. 162–171. DOI: <https://doi.org/10.1016/j.scienta.2010.09.016>
- Siegel, B.Z. (2003): Plant peroxidases – an organic perspective. - *Plant Growth Regulation*. **12**: 3. 303-312. DOI: <https://doi.org/10.1007/bf00027212>
- Tanou, G., Molassiotis, A., Diamantidis, G. (2009): Induction of reactive oxygen species and necrotic death-like destruction in strawberry leaves by salinity. *Environ Exp Bot.* **65**: 2-3. 270–281. DOI: <https://doi.org/10.1016/j.envexpbot.2008.09.005>
- Tattersall, E.A., Grimplet, J., Deluc, L., Wheatley, M.D., Vincent, D., Osborne, C., Ergul, A., Lomen, E., Blank, R.R., Schlauch, K.A., Cushman, J.C., Cramer, G.R. (2007): Transcript abundance profiles reveal larger and more complex responses of grapevine to chilling compared to osmotic and salinity stress. *Funct Integr Genomics*. **7**: 4. 317-333. DOI: <https://dx.doi.org/10.1007/s10142-007-0051-x>
- Vierheilig, H., Coughlan, A. P., Wyss, U., Piché, Y. (1998): Ink and vinegar, a simple staining technique for arbuscular-mycorrhizal fungi. *Applied and Environmental Microbiology*. **64**: 12. 5004-5007.
- Wahid, A., Gelani, S., Ashraf, M., Foolad, M.R. (2007): Heat tolerance in plants: An overview. *Environmental and Experimental Botany*. **61**: 3. 199-223. DOI: <https://doi.org/10.1016/j.envexpbot.2007.05.011>
- Wu, Q.S. (2011): Mycorrhizal efficacy of trifoliolate orange seedlings on alleviating temperature stress. *Plant Soil Environ.* **57**: 10. 459-464.
- Wu, Q.S., Zou, Y.N., Abd_Allah, E.F. (2014): Mycorrhizal Association and ROS in Plants. In: P. Ahmad (Ed): *Oxidative Damage to Plants*. Elsevier Inc. pp 453- 475. DOI: <http://dx.doi.org/10.1016/B978-0-12-799963-0.00015-0>
- Zhou, J., Wang, J., Li, X., Xia, X-J., Zhou, Y-H., Shi, K., Chen, Z., Yu, J-Q. (2014): H₂O₂ mediates the crosstalk of brassinosteroid and abscisic acid in tomato responses to heat and oxidative stresses. *Journal of Experimental Botany*. **65**: 15. 4371–4383. DOI: <https://doi.org/10.1093/jxb/eru217>
- Zhu, X.C., Song, F.B., Xu, H.W. (2010): Influence of arbuscular mycorrhiza on lipid peroxidation and antioxidant enzyme activity of maize plants under temperature stress. *Mycorrhiza*. **20**: 5. 325-332. DOI: <https://doi.org/10.1007/s00572-009-0285-7>
- Zhu, X.C., Song, F.B., Liu, S.Q., Liu, T.D. (2011): Effects of arbuscular mycorrhizal fungi on photosynthesis and water status of maize under high temperature stress. *Plant Soil*. **346**: 1-2. 189-199. DOI: <https://doi.org/10.1007/s11104-011-0809-8>

***In vitro* activity of the cinnamon essential oil against the plant pathogen *Septoria melissae* desm.**

Gergő KOVÁCS^{1,2} – Éva ZÁMBORI-NÉMETH¹ – Péter RADÁCSI¹ – Géza NAGY³

1: Department of Medicinal and Aromatic Plants, Szent István University, H-1118 Budapest, Villányi út 29-43., Hungary. E-mail: kovacs.gergo.phd@gmail.com

2: Department of Plant Pathology, Szent István University, H-1118 Budapest, Villányi út 29-43., Hungary

3: Directorate of Plant Protection, Soil Conservation and Agri-environment, National Food Chain Safety Office, H-1118 Budapest, Budaörsi út 141-145., Hungary

Abstract: *Septoria* leafspot caused by the mitosporic fungus *Septoria melissae* Desm. is the most significant disease of lemon balm (*Melissa officinalis* L.). The fungus frequently appears in the plantations and causes serious yield losses or decreases the quality of the drug. At present plant protection of lemon balm is inadequate due to the lack of authorised plant protection products. The essential oil of the cinnamon bark (CEO) has a strong antimicrobial activity proved by several *in vitro* experiments. Therefore the goal of our work was to test antifungal effect of the CEO against the pathogen *Septoria melissae* Desm.

In vitro tests were carried out with three concentrations (0.3%, 0.1%, 0.03%) of the CEO against a Hungarian and a Polish isolate of the fungus. Inhibition of the germination of the conidia as well as the mycelial radial growth was investigated. Supplementary tests were carried out with colonies of the fungi transferred to growth media without CEO after an incubation period of 14 and 35 days on media containing CEO.

Our results showed that all the applied concentrations of CEO have very strong (98.07-100%) inhibitory effect on the mycelial growth of both isolates. Germination of conidia was also blocked on each medium containing CEO. However, the effect of the CEO at 0.03% concentration was reversible. Mycelium began to grow again on media without CEO after transfer.

Based on the results, further investigation of CEO as a potential plant protection product in lemon balm crops is recommended.

Keywords: *Septoria* leafspot, lemon balm, growth inhibition, environmental friendly, medicinal plants

Received 13 April 2018, Revised 21 September 2018, Accepted 10 October 2018

Introduction

Septoria leafspot caused by the mitosporic fungus *Septoria melissae* Desm. is the most important disease of lemon balm (*Melissa officinalis* L.) (Nagy et Horváth, 2010; Jadczyk et Pizoń, 2017; Wielgusz et Seidler-Łożykowska, 2017). This pathogen frequently appears in lemon balm crops and causes serious yield losses expressed by the severe leaf fall as a consequence of the lack of proper pest management. Furthermore, even a moderate infection may highly influence the quality of the drug by decreasing the essential oil (EO) content and modifying the rate of the main compounds (Aulerio et al., 1995; Kowalska et al., 2014). At present, the plant protection of lemon balm is inadequate, due to the strict regulation of the maximum residue levels allowed in herbal products (Kowalska et al., 2014; Bernáth et Zámboři-Németh, 2015). In Hungary, only a few plant protection products are authorised in

this crop which cannot provide reliable control of the pathogens (Ocskó et al. 2017).

In the light of the facts mentioned above, the development of new and environmental friendly methods for the protection of lemon balm is essential. EOs are potential substances for this purpose. The antimicrobial activity of these compounds has been confirmed by several *in vitro* and *in vivo* investigations in the past decades. EO of the cinnamon bark (CEO) is one of the most highly investigated EOs in the field of food science. The results of an *in vitro* experiment of Ju et al. (2018) demonstrated the efficacy of the CEO against *Penicillium* and *Aspergillus* species. The results of López et al. (2007) indicate, that *trans*-cinnamaldehyde, the main compound of this oil might be responsible for its antimicrobial activity. Another *in vitro* experiment of Feyaerts et al. (2018) carried out with 175 EOs confirmed the previous

observation. The authors' results showed, that the EOs which contain high amount of aldehydes, had the highest inhibitory effect in vapour phase.

Field trials were carried out by Hochbaum et Nagy (2013) with the combination of the EOs of *Thymus vulgaris* and *Cinnamomum zeylanicum*. Spraying a mixture of the EOs significantly decreased the disease incidence of *Monilinia* blossom blight (*Monilinia laxa*) of apricot and leafcurl (*Taphrina deformans*) of peach in the applied 0.05% and 0.1% concentrations, respectively. The wide spectrum of antimicrobial activity of CEO was also demonstrated by a small plot trial of Kovács et al. (2013). The application of this EO gave the best result against the *Fusarium* head blight (*Fusarium graminearum*, *F. culmorum*) of winter wheat on artificially inoculated plants.

In medicinal plant production only few data are available (e.g. Kovács et Nagy (2014)) about the use of EOs as plant protection product. Therefore, the goal of our recent study was to test the efficacy of the EO of *C. zeylanicum* bark (*Aetheroleum cinnamomi zeylanici corticis*) against the most important pathogen of lemon balm: *S. melissae* Desm.

Materials and methods

Characterisation of the applied essential oil

The investigated CEO was purchased from a commercial company (Aromax Zrt., Hungary). The chemical composition was determined

Table 1. Chemical composition of the applied cinnamon essential oil according to the GC-MS analysis

| Compound | RT | LRI | % |
|----------------------------------|-------|----------|------|
| α -Pinene | 5.56 | 938.000 | 0.6 |
| Benzaldehyde (artificial almond) | 6.28 | 967.000 | 0.2 |
| β -Pinene | 6.64 | 980.856 | 0.1 |
| p-Cimol (para) | 8.09 | 1025.856 | 1.4 |
| Limonene | 8.19 | 1028.529 | 1.5 |
| 1,8-Cineole | 8.38 | 1033.503 | 2.3 |
| Linalool | 10.76 | 1097.246 | 2.8 |
| α -Terpineol | 14.55 | 1189.123 | 1.2 |
| cis-Cinnamaldehyde | 15.73 | 1217.000 | 0.1 |
| trans-Cinnamaldehyde | 17.72 | 1264.000 | 78.8 |
| trans-Anetol | 18.51 | 1283.175 | 2.8 |
| Eugenol | 21.44 | 1360.822 | 1.3 |
| α -Copaene | 22.03 | 1376.986 | 0.04 |
| β -Caryophyllene | 23.68 | 1419.951 | 2.5 |
| trans-Cinnamyl acetate | 24.80 | 1448.000 | 3.6 |
| α -Humulene | 25.07 | 1454.187 | 0.5 |
| orto-Methoxy Cinnamaldehyde | 28.24 | 1536.000 | 0.1 |
| Benzyl benzoate | 36.83 | 1773.296 | 0.1 |
| Total | | | 99.9 |

by gas chromatography mass spectrometric method, using an Agilent Technologies 6890N instrument equipped with HP-5MS capillary column (30 m \times 0.25 mm i.d. \times 0.25 μ m) and an Agilent Technologies MS 5975 inert mass selective detector. The temperature program was the following: initial temperature 60°C, then by a rate of 3°C/min up to 240°C; the final temperature was kept for 5 min. Carrier gas was helium (1 mL min⁻¹), injector and detector temperatures were 250°C. Split ratio: 30:1. Injected quantity: 0.2 μ l (solvent: n-hexane). The percentage composition of the EO was computed from the GC peak areas. Ionization energy was 70 eV. The MS were recorded in full scan mode that revealed the total ion current (TIC) chromatograms (mass range m/z 50–550 uma). The components were identified by linear retention indices that were calculated using the generalized equation of Van Den Dool and Dec. Kratz (1963), and by mass spectra by using NIST MS Search 2.0 library and Adams mass spectra library (Adams, 2007).

According to the GC-MS analysis, the applied EO contains *trans*-cinnamaldehyde in 78.8% (Table 1.). Other components of the oil in higher amounts were the followings: *trans*-cinnamyl-acetate (3.6%), *trans*-anetol (2.8%), linalool (2.8%) and eugenol (1.3%).

Origin and maintenance of fungal isolates

Two isolates of *S. melissae* Desm. were used in our experiment. Target pathogens were

isolated from infected leaves of lemon balm on malt extract agar (MEA). The leaves were collected at Budapest-Soroksár (47°24'08.7"N 19°09'03.9"E) in Hungary (HBS) and at Warsaw-Wilanów (52°09'36.9"N 21°06'08.2"E) in Poland (PWW). The fungal isolates have been maintained on malt extract agar (MEA) at 24°C without light.

Experimental methods to test the antimicrobial properties

Inhibitory effect of CEO on both the *germination of conidia* and the *mycelial radial growth* were tested by the following method.

For the testing of the effect of CEO on the *germination of conidia*, suspension of conidia was prepared by suspending conidial exudates developed on the surface of monosporic cultures of the isolate PWW in distilled water. Concentration of the suspension was determined by haemocytometer. The amount of conidia in the suspension was approximately 1.6×10^6 conidia/ml. Preparation of media carried out according to the followings. CEO was evenly diluted in MEA at the 0.3%; 0.1% and 0.03% concentrations in Petri dishes. Silwet Star wetting agent in 0.02% was also applied in each treatment to improve homogeneity. Conidia growing on medium containing no CEO served as control. Besides, MEA mixed with the wetting agent were also tested. 30 µl of conidial suspension was spread evenly on the surface of the media in two replications. The number of germinated conidia was counted 24 hours after inoculation on ten randomly chosen $10^6 \mu\text{m}^2$ area of the media. Conidium was considered germinated if measured length of the germ tube was longer than the length of the conidium. Ratio of the germinated and non germinated conidia was expressed in percentage.

For the testing of the effect of CEO on the *mycelial radial growth* preparation of media and the treatments were carried out in the same way as described above. 31 days old monosporic cultures of the slow growing fungus *S. melissae* Desm. were used to obtain adequate amount of inoculum. Mycelial fragments from the margin of the cultures of both origin, HBS and PWW, were placed onto previously prepared media

aseptically. The experiment was set up in 10 replicates. The radial growth of mycelium was determined by measuring the diameter of the colonies 0 and 14 days after inoculation. The ratio of inhibition was calculated by the following formula: $PI = ([C_A - C_S] - [T_A - T_S]) / (C_A - C_S) \times 100$, where C_S is the area of the colonies on intact medium at the time of inoculation, C_A is the area of the colonies on intact medium on day 14 after inoculation, T_S is the area of colonies on the treated medium at the day of inoculation and T_A is the size of the culture on the treated medium on day 14 after inoculation. Growth inhibition was expressed in percentage comparing the growth rate of CEO treated colonies with the control ones.

The *survival rate of the treated cultures* was investigated with HBS isolates. After the maintenance of 14 and 35 days of the cultures on medium containing 0.3%, 0.1% and 0.03% CEO, mycelial fragments were placed onto media containing no CEO. The mycelial radial growth of these cultures was measured on the 3rd, 8th and 14th days after transfer. The mycelial growth was calculated by the following formula: $MReg = S_A - S_S$, where S_S is the area of the colonies on intact medium at the time of transfer and S_A is the area of the colonies on intact medium at the time of the assessment. The results of the calculations were expressed in mm^2 .

Statistical analysis

Data were analysed by the IBM SPSS Statistics 22 software. Univariate ANOVA (ANOVA) was used to evaluate the significant differences among the inhibitory effect of the investigated CEO in different dilutions. Multivariate ANOVA (MANOVA) was used to detect the significant differences between the growth rates of the transferred cultures during survival rate test. The normality of the residuals was tested according to the Saphiro-Wilk and Kolgomorov-Smirnov tests. If the normality could not be justified by the mentioned analyses, it was verified by the skewness and the kurtosis. Homogeneity of variances was tested by the Levene's method. If the homogeneity assumption was not violated, Tukey *post hoc* test was used to group the genotypes. Otherwise the separation was made by the Games-Howell test.

Table 2. The rate of inhibition of mycelial radial growth of the two *Septoria melissae* Desm. isolates by cinnamon essential oil (CEO) (Legends: The abc letters refer to the significantly different groups according to the ANOVA test)

| Treatment | Isolate | |
|-------------|----------|-----------|
| | HBS | PWW |
| CEO 0.3% | 97.68% a | 100.00% a |
| CEO 0.1% | 98.07% a | 100.00% a |
| CEO 0.03% | 99.65% a | 100.00% a |
| Silwet Star | 15.38% b | 1.01% b |

Table 3. The effect of cinnamon essential oil (CEO) treatments on the germination of conidia of the PWW isolate

| Treatment | Conidia | |
|-------------|------------|----------------|
| | germinated | not germinated |
| CEO 0.3% | 0% | 100% |
| CEO 0.1% | 0% | 100% |
| CEO 0.03% | 0% | 100% |
| Silwet Star | 100% | 0% |
| Control | 100% | 0% |

Results and discussion

Inhibitory effect on the germination of conidia and on the mycelial radial growth

Conidia of the PWW isolate did not germinate on any of the treated medium containing CEO (Table 3.; Figure 1.). The Silwet Star wetting agent did not influence the germination of the conidia.

The CEO achieved a powerful inhibitory effect on the *mycelial growth* of both isolates of *S. melissae* Desm. (Table 2.) as well. All applied

concentrations showed approximately 100% growth inhibition. Differences among the concentrations were not significant according to statistical analysis. The applied wetting agent showed a very low growth inhibition (1.01-15.38%) of the mycelia of both isolates.

Information about the inhibitory effect of the CEO against *Septoria* species was not found in the international literature. The effect of other EOs (thyme, fennel, rosemary, etc.) on the related fungus *Septoria tritici* (syn.: *Zymoseptoria tritici*) was investigated by Matusinsky et al. (2015). The EO of thyme inhibited mycelial growth by 100%. The strong antifungal activity of CEO was demonstrated by several authors, although against organisms other than *Septoria* spp. In the *in vitro* experiment of Kovács et al. (2013) CEO showed 100% inhibition of the mycelial growth of *Fusarium graminearum*. Another trial carried out with the pure compound *trans*-cinnamaldehyde (the main component of CEO) revealed a strong inhibitory effect on the mycelial growth of both *Aspergillus flavus* (100% inhibition) and *Penicillium islandicum* (91% inhibition) (López et al., 2007).

Survival rate of the treated cultures

Although a good inhibitory activity of CEO was detected in each applied concentrations, survival rate trials showed that the fungal mycelium was not completely destroyed in all treatments. The transferred mycelial fragments from colonies grown for 14 days on MEA containing 0.03% CEO began to grow on MEA without CEO. The measured growth intensity of these colonies was




Figure 1. Inhibition of the germination of conidia by cinnamon essential oil (CEO) (Legends: Surface of the media in the following treatments: A – control; B – Silwet Star 0.02%; C – CEO 0.03%; D – CEO 0.1%; E – CEO 0.3%. Scale 100 μ m)




Figure 2. Mycelial radial growth of HBS colonies on MEA transferred after 14 days incubation on CEO containing MEA (Legends: The ABC letters means the significantly different groups in one time according to the MANOVA test. Normal letters belongs to the CEO 0.03% and italic letters to the control. The stars sign the CEO 0.3% and CEO 0.1% treatments where mycelial growth was not observed).

significantly higher than that of the control ones (Figure 2.). After a longer incubation period (up to 35 days) on MEA containing 0.03% CEO only a few colonies, which were transferred to MEA without CEO started to grow with very different intensity. This part of the experiment was not evaluated statistically. Colonies transferred from MEA containing higher CEO concentrations (0.1% and 0.3%) were not able to grow on MEA without CEO.

Conclusion

Our experiment showed that the investigated CEO had a powerful *in vitro* inhibitory effect on conidial germination and the mycelial growth of both *S. melissae* Desm. isolates. In accordance with the results of the *in vitro* experiment of Kovács et al. (2013) carried out with the wheat pathogen *Fusarium graminearum*, CEO had approximately 100% growth inhibition even in the lowest (0.03%) concentration. However, in our experiment this CEO concentration caused

only a reversible inhibition and was not lethal to the mycelium of *S. melissae* Desm. Treated isolates managed to grow on MEA with no CEO after transfer.

According to Gutiérrez et al. (2010) EOs break the integrity of the cytoplasmic membrane, which leads to the collapse of fungal cell. CEO might have the same effect on the conidial cell or on the membrane of the germ tube of *S. melissae* Desm.

Based on our results CEO applied in 0.1% and 0.3% concentrations can be a potentially efficient plant protection product used for the control of *S. melissae* Desm. However, the practical application needs further investigations including *in vivo* trials.

Acknowledgements

Géza Nagy's research was supported by the European Union and the State of Hungary, co-financed by the European Social Fund in the framework of TÁMOP 4.2.4. A/1-11-1-2012-0001 'National Excellence Program'

References

- Adams, R.P. (2007): Identification of essential oil components by gas chromatography/mass spectrometry, 4th Edition. Carol Stream, Ill: Allured Pub Corp.
- Aulerio, A. Z., Zambonelli, A., Bianchi, A., Albasini, A. (1995): Micromorphological and chemical investigations into the effects of fungal diseases on *Melissa officinalis* L., *Mentha pieprita* L. and *Salvia officinalis* L. Journal of Phytopathology. **143** (3): 179–183.
- Bernáth J., Zámoriné Németh É. (2015): Gyógynövény kultúrák magyarországi növényvédelmének időszzerű kérdései, (Up to date aspects of the plant protection in the Hungarian medicinal plant production) Növényvédelem, **51** (1): 25–36.

- Feyaerts, A. F., Mathé, L., Luyten, W., De Graeve, S., Van Dyck, K., Broekx, L., Van Dijck, P. (2018): Essential oils and their components are a class of antifungals with potent vapour-phase-mediated anti-*Candida* activity. *Scientific Reports*. DOI:10.1038/s41598-018-22395-6
- Gutiérrez, G., Batlle, R., Sánchez, C., Nerín, C. (2010): New approach to study the mechanism of antimicrobial protection of an active packaging. *Foodborne Pathogens and Disease*. **7** (9): 1063–1069.
- Hochbaum T., Nagy G. (2013): Egy illóolaj kombináció alkalmazásának lehetősége kajszi- és őszibarack kórokozói, valamint kártevő molyfajai ellen, (Efficacy of essential oils against the major pathogens and fruit moths of apricot and peach). *Növényvédelem*. **49** (1): 8–15.
- Jadczak, P., Pizoń, K. (2017): Identification of taxa of microscopic fungi occurring on selected herbal plants and possible methods of their elimination. *World Scientific News*. **69**: 1-17.
- Ju, J., Xu, X., Xie, Y., Guo, Y., Cheng, Y., Qian, H., Yao W. (2018): Inhibitory effects of cinnamon and clove essential oils on mold growth on baked foods. *Food Chemistry* **240**: 850–855.
- Kovács B., Horváth A., Sárosi S., Nagy G. (2013): A mentafélék és a fahéj kivonatának és illóolajának alkalmazási lehetősége az őszi búza fuzáriumos kalászfertőzése ellen, (Application of extracts and essential oils of mints and cinnamon against fusarium head blight of winter wheat). *Növényvédelem*. **49**: (9) 403–409.
- Kovács F., Nagy G. (2014). Mentaolajok hatása a *Ramularia menthicola* kórokozó és gazdanövénye kapcsolatban. *Kertgazdaság* **46**: (2) 78-84.
- Kowalska, J., Remlein-Starosta, D., Seidler-Łożykowska, K., Jan Bocianowski, J. (2014): Can *Trichoderma asperellum* [t1] stimulate growth of lemon balm (*Melissa officinalis* L.) in different systems of cultivation? *Acta Scientiarum Polonorum, Hortorum Cultus* **13** (1): 91–102.
- López, P., Sánchez, C., Batlle, R., Nerín, C. (2007): Vapor-phase activities of cinnamon, thyme, and oregano essential oils and key constituents against foodborne microorganisms. *Journal of Agricultural and Food Chemistry*. **55**: 4348–4356.
- Matusinsky, P., Zouhar, M., Pavela, R., Novy, P. (2015): Antifungal effect of five essential oils against important pathogenic fungi of cereals. *Industrial Crops and Products*. **67**: 208–215.
- Nagy G., Horváth A. (2010): Gyógynövények szeptóriás levélfoltosságai Magyarországon. (*Septoria* leaf spot diseases of medicinal plant species in Hungary) *Növényvédelem*. **46** (4): 145–153.
- Ocskó Z., Erdős Gy., Molnár J. (2017): Növényvédő szerek, termésvetelő anyagok 2017. vol. I. Agrixen Kiadó. Budapest. p. 789.
- Van Den Dool, H., and Kratz, P. D. (1963): A generalization of the retention index system including linear temperature programmed gas—liquid partition chromatography. *Journal of Chromatography*. **11**: 463-471.
- Wielgusz, K., Seidler-Łożykowska, K. (2017): Fungi colonizing and damaging different parts of some medicinal plants. *Herba Polonica*. **63** (2): 18–26.

Preliminary results of a pot experiment with the combined effects of a terrestrial isopod species (*Porcellionides pruinosus*, Brandt 1833) and organic mulching on tomato

Anett Mészárosné PÓSS – Angéla ZANKER – Franciska TÓTHNÉ BOGDÁNYI – Ferenc TÓTH

Szent István University, Faculty of Agricultural and Environmental Sciences, Plant Protection Institute, 2100 Gödöllő, Páter Károly utca 1., Hungary; E-mail: toth.ferenc@mkk.szie.hu

Abstract: Organic mulching practice has a beneficial effect on soil life. Increased soil activity may in turn lead to increased soil fertility. To study the effect of terrestrial isopods and organic mulch on potted tomatoes we measured the growth, vitality and yield of plants. Half of the pots had the surface of the soil covered with a mulch of equal quantity of walnut (*Juglans regia*), elm (*Ulmus minor*) and maple (*Acer platanoides*) leaves; and half of them had the soil uncovered. 20 *Porcellionides pruinosus* (Brandt, 1833) individuals were introduced to every second pot. There were four treatments in 10 replicates: (1) I+M+ (isopods, mulching); (2) I+M- (isopods, no mulching); (3) I-M+ (no isopods, mulching) and (4) I-M- (no isopods, no mulching). To determine the microbial activity of the soil, fluorescein diacetate hydrolysis activity was measured. The fluorescein diacetate concentration was higher in the “combined” treatment than in the “isopods only” treatment, and in the “mulch only” treatment. Regardless of treatment, overall microbial activity figures were lower after the experiment than their respective starting values. The “combined” treatment significantly increased the generative growth of tomatoes. The number of flowers was significantly higher, and thus significantly more tomato was harvested when compared to the “control” treatment. Our preliminary results indicate the beneficial effects of *P. pruinosus*, because its presence had an advantageous influence on tomato yield.

Keywords: Oniscidea, soil fertility, woodlice, soil cover

Received 4 September 2018, Revised 25 September 2018, Accepted 30 October 2018

Introduction

Higher habitat and ecological diversity in a home-scale horticulture decreases the pressure from pests (Philpott 2013). Tomato production has an increasing trend both in the total area of production and the amount of harvest worldwide (FAOSTAT data, 2018). Tomato is typical to most home vegetables patches as well, with most of the gardeners cultivating more than one variety.

Consumers believe that landraces are nutritionally more valuable (Casals et al. 2011). While this argument has not yet been proven in all cases, genetic variability differs among various landraces and depends on selection pressure (Passam et al. 2007).

For our study we selected a Hungarian landrace (Dány) with the preconception in mind that landraces are better adopted to local conditions. We wanted to find a method that has the potential to increase the success of home-scale tomato production, to increase the amount of yield and the vitality of soil at the same time. In our study we were looking for providing production conditions that are relatively less experienced with yet are low on the budget and energy demand.

The benefits of using natural mulches are manifold. Suppressing weeds, preserving soil moisture and therefore, stabilizing soil structure are just some of them. One has to mention that besides contributing to the production of healthy, marketable crops, good mulching practice provides suitable conditions for beneficial organisms, making the use of mulch a factor of key importance within the complexity of crop protection (Mancinelli et al. 2015, Diver et al. 1999). Mancinelli et al. (2015) found that organic mulching, irrigation and fertilization decreased CO₂-emission and increased C-storage of the soil. There are many soil-dwelling living organisms involved in nutrient cycles to maintain soil fertility. Terrestrial isopods hide under dead plant matter, using it as shelter and food source at the same time (Stachursky 1968). Woodlice are also known for manipulating the soil: to consume decaying organic matter and release it back as faeces to the soil again. Within their intestinal tract, decaying materials are broken down to small, organic molecules, which are now available for the soil bacteria and fungi for the processes of immobilization, mineralisation and humification. This is how isopods have an indirect influence on the microbial activity of

the soil (Zimmer and Topp 1999; Lavelle et al. 2006; Vilisics et al. 2012). *Porcellionides pruinosus* was found the most abundant species in compost heaps (Farkas and Vilisics 2013).

The positive effect of mulching has been well documented in recent times (Campiglia et al., 2011, Shirgure, 2012, Kumar, 2014), and the beneficial effects of faecal pellets of woodlice and millipedes on the growth of test plants were proven in pot experiments (Gere 1956; Farkas et al. 2017).

Therefore we expected mulching to increase the microbial activity of the soil, and tomato yield; and by adding (not faecal pellets but living individuals of) woodlice to the system, to further increase the beneficial effects of mulch. We also predicted that the lack of mulch would make woodlice leave the pots, and that in those pots there would be no significant change in either the microbial activity of the soil or in the yield.

Methods

Selection of plants and test animals

For our purposes, the regional cultivar Dány was used. This determinate, mid to late season variety is mainly for fresh consumption. It has a moderate growth vigour, and a strong foliage rejuvenation (ÖMKi 2015). The growing period for this variety was observed to start late March to early April with sowing, and to end mid to late September with the final harvests. Tomato seeds were sown in Florimo® general potting soil

(*Sphagnum* moss peat, flat peat, organic humus, composted cattle manure, clay, fertilizer mixture, with a pH of 6.4 ± 0.5). The plants were grown in greenhouse until the age of six weeks, when the seedlings were transplanted one by one into a pot. There were 40 pots with the same soil, one seedling per pot. Thereafter Isopods were placed into the pots with or without leaf litter.

Our test animal was *Porcellionides pruinosus*. Besides its simple availability and synanthropic lifestyle, this isopod became our test species because it responds well to various environmental circumstances and is easy to rear among artificial conditions.

Porcellionides pruinosus individuals were collected from the Regional Waste Management Center of Púsztaámor by handsorting. The animals were bred and kept at the Institute of Plant Protection of Szent István University. For identification we used the taxonomic key of Farkas and Vilisics (2013).

Experimental design

The study started on 24 May 2016 in the experimental area of the Plant Protection Institute of the Szent István University in Hungary (SZIE NVI) in a partially covered glasshouse and it was terminated on 23 September 2016, after the growing period. Before distributing the same amount of potting soil („Original soil”) into every pot, the enzymatic activity of the soil was measured with *fluorescein diacetate*




Figure 1. Pot experiment with four different treatments: studying the effect of the addition of mulch and isopods, using a tomato landrace variety as test plant

(FDA) hydrolysis assays tests. There were four treatments in 10 replicates as follows: (1) I+M+ (isopods, mulching); (2) I+M- (isopods, no mulching); (3) I-M+ (no isopods, mulching) and (4) I-M- (no isopods, no mulching) (Fig 1.).

Placement of *Porcellionides pruinosus* individuals and mulch

In each I+ pot, 20 specimen of *P. pruinosus* adults (with an average body weight of 228.5 mg) were placed. We chose to not differentiate between genders as this type of selection would have hurt the individuals, but gravid females were excluded. The number of individuals, adults and juveniles were counted at the completion of the experiment. Mulch material was collected from the surrounding areas: the mulch mixture contained walnut (*Juglans regia*), elm (*Ulmus minor*) and maple (*Acer platanoides*) leaf litter of local trees in a ratio of 1:1:1. Every mulched pot was covered with an average of 39.6 g of leaf matter. The soil of the pots was sieved at the end of the experiment, and the number of adult and juvenile isopods was recorded.

Soil microbial activity

To measure and monitor the microbial activity of soil, we followed the method of SCHNÜRER and ROSSWALL (1982). Each pot was sampled before the experiment and at the end of the experiment again by taking 2 g of soil. Samples were incubated for two hours at 24°C with 10 ml sodium-phosphate buffer (60mM, pH= 7.6) and 100 µl of FDA (2 mg/l). The rotary shaker was set at 300 RPM. After two hours the reaction was halted by the addition of acetone (10 ml). The solution was sieved, and the amount of hydrolysed FDA was measured. Absorbance values were obtained by using a Haca DR/2000 spectrophotometer at 490 nm. Calibration curves were drawn using solutions of known fluorescein concentration, and thus, the amount of fluorescein obtained from our samples during the two-hour period was calculated. Each test was replicated ten times. The values were calculated for one gram of dry soil as mg of fluorescein.

Parameters of tomato

Both the tomatoes and their respective pots were sampled every fortnight during the growing

season. The number of flower buds, flowers and leaves were recorded. Indicators of maturity: the number of fruits, their weight were also measured. The number of flower buds, flowers and fruits, and the total weight fruits per plant were regularly monitored. Root weight was measured at the end of the growing season.

Statistical analyses

Levene's test for homogeneity of variance, ANOVA with Tukey's posthoc test, Kruskal-Wallis test with Mann-Whitney pairwise comparisons and Welch test (PAST® program: Hammer et al 2001) were used for statistical analyses.

Results

Soil microbial activity

The fluorescein concentration values (µg/ml) of treatments revealed that the addition of mulch material resulted in higher microbial activity (Fig. 2) ($F=14.35$, $df=21.77$, $p=7.015 \times 10^{-6}$).

Regardless of the presence of isopods, there was a significant difference between treatments with or without mulch, that is, between treatments with




Figure 2. Total soil microbial activity by fluorescein concentrations (µg/ml) in a tomato pot experiment with mulch and the isopod *Porcellionides pruinosus* [I=Isopod, M=Mulch; the same letters above bars indicate the lack of significant ($p < 0.05$) difference; error bar: \pm SE; ANOVA, Tukey's posthoc test]




Figure 3. The number of *Porcellionides pruinosus* individuals at the end of a tomato pot experiment where isopods and mulch were both added to the soil [I=Isopod, M=Mulch; the same letters above bars indicate the lack of significant ($p < 0.05$) difference; error bar: \pm SE; Kruskal-Wallis test, Mann-Whitney pairwise comparisons]

I-M- and I-M+; and between treatments I+M- and I+M+. The fluorescein values measured for the two no-mulch treatments (I-M- and I+M-) were also significantly lower than the initial values (Original soil) measured prior to the pot experiment. Statistical analysis confirmed that the original microbial activity of the soil was not significantly different from the values obtained in treatments with mulch (I-M+ and I+M+) but was significantly higher than those without mulch (I-M-) and (I+M-) (Annex 1).

effect on number of *P. pruinosus* individuals (Fig.3., Annex 2) ($H=17.42$; $p=3.40 \times 10^{-6}$). The number of juveniles were also recorded, but due to low recapture no reliable conclusions were drawn from these data.

Parameters of tomato

Number of flower buds, flowers and leaves

No significant difference was found in the maximum number of flower buds and leaves between the treatments (Annexes 3-4) There was a significant difference however, in the maximal number of flowers ($F=5.246$, $df=19.83$, $p=0.008$; CI95% and average I-M- 0.682 and 3.9, I-M+ 0.787 and 5.5, I+M- 0.889 and 4.5,

Porcellionides pruinosus and mulch

The number of *Porcellionides pruinosus* individuals at the end of the experiment was lower than initially, but mulching had a beneficial




Figure 4. The maximum number of tomato flowers in a pot experiment studying the effect of *Porcellionides pruinosus* and mulch [I=Isopod, M=Mulch; the same letters above bars indicate the lack of significant ($p < 0.05$) difference; error bar: \pm SE; ANOVA, Tukey's post hoc test]




Figure 5. Tomato yield according to the presence or lack of a woodlice species, *Porcellionides pruinosus* and mulch in a tomato pot experiment [I=Isopod, M=Mulch; the same letters above bars indicate the lack of significant ($p < 0.05$) difference; error bar: \pm SE; ANOVA, Tukey’s post hoc test]

I+M+ 0.989 and 6.1; Tukey’s pairwise I+M+ 0.005) (Fig. 4, Annex 5).

When only isopods were added (I+M-), the decreasing trend in the number of flower buds and flowers was slower than in any of the other treatments.

The number of fruits, their weight and root weight

None of the treatments had a significant effect on the number of fruits per plant ($p = 0.079$).

Neither the addition of isopods, nor the addition of mulch alone had a significant effect on fruit weight, but isopods and mulch together had a recognizable effect on tomato yield ($F = 5.78$, $df = 19.76$, $p = 0.005$) (Annex 6).

The effect of adding isopods to pots on tomato yield is recognizable ($p = 0.013$).

On the other hand, no significant difference was recorded when only mulch was added (M+). According to these results, the difference between the yield of tomato plants treated and those not treated with isopods was statistically not proven. None of the treatments alone had a significant effect on the number of fruits harvested, but there was a significant dissimilarity between the effect of the “combined” (I+M+) and the “control” (I-M-) treatment on the weight of the yield (Fig. 5).

Measuring the dry and fresh matter content of the roots revealed that none of the treatments had a significant influence on the growth of the root system ($p = 0.486$). Yield (fruit weight per plant) kept increasing continuously as the season wore on. The increase was the highest with the “mulch only” (I-M+) treatment; and its values were followed by the values of the “combined” treatment (I+M+) (Fig. 6).




Figure 6. Cumulative tomato yield growth during the experiment with four treatments (I=Isopod, M=Mulch) in a tomato pot experiment

Discussion

Adding mulch alone did not have a significant effect on tomato yield, but a combination of mulch and isopods did. While Tuf and Tufová (2005) claimed that soil macroinvertebrates, due to their role in the ecosystem, increase soil fertility, we only recorded a significant effect of macroinvertebrates when combined with mulch.

Since the type and origin of mulch materials may have an impact on production indicators (Zribi et al. 2015), based on our previous studies (Fehér et al, 2017), we now decided to use organic mulch and in particular, the locally available leaf litter: an even mixture of walnut (*Juglans regia*), elm (*Ulmus minor*) and maple (*Acer platanoides*). Similarly to our experiment, Förster et al. (2006) also mulched with a various mixture of forest leaves, but in their set up faecal pellets of millipedes and woodlice were also added to crops, while we preferred to use the organisms themselves, which were found useful by Tantachasatid et al. (2017) in their experiment. Our prediction, based on the results of Förster et al. (2006) was proven: soil microbial activity declined in the absence of mulch in our treatments. Without a proper organic cover, soil was more prone to desiccation, its microbial activity decreased, and it also failed to provide isopods with habitat and shelter.

Within this limited timeframe of our present experiment mulch material did not have the time to enhance the nutrient content of the soil. We believe mulch provided a habitat and a shelter for our isopods as was found by Diekötter et al (2010). Similarly to a natural environment, isopods were not confined: they had the ability to leave the pots, and we assume that they hardly ever found their way back to the same, or to any pot. Where there were none at the beginning, none was found at the termination of the experiment either. No new isopods settled to pots, and migration was low, probably because it was hindered by the mulch leaves. In the end the number of juveniles was measured but it is not a reliable data because the sex ratio was not determined before the experiment.

Individuals of *Porcellionides pruinosus* contributed to leaves being decomposed and

buried into the soil, although the contribution was definitely smaller than that of the dung beetle, also considered an ecosystem engineer (Johnson et al. 2016). This raises the question: would the impact of isopods on soil and plant have been higher had the isopods been introduced in considerably larger numbers. If so, is there an optimum number of individuals to be introduced. Is this optimum amount different for each isopod species, for soil types, and in general, what influence ecological conditions have on this data? To answer these questions, further studies are needed.

Conclusions

Our preliminary results indicate the beneficial effects of the combined application of leaf litter mulch and *Porcellionides pruinosus* on tomato yield.

The observation that isopods were found only in pots where they were introduced at the beginning of the experiment may suggest that these animals do not leave their micro-environment when the conditions were favourable for their survival. One may thus suspect that the useful influence of isopods is only present when the conditions favoured by these arthropods are created within the course of crop production protocol, that is, when mulching is part of the management.

We propose to test this innovative system in larger scale experiments. The biomass quantity of woodlice in our experiment was not remarkable enough and more experiments with higher isopod density are suggested.

We consider treating production areas with mulch and isopods a preliminary step in elaborating a new element of technology in crop production. We contemplate the possibilities of expanding our studies to a larger scale, to arable lands. At present, *P. pruinosus* is already a marketable item: it is reared mainly as prey (e.g. <http://www.ebay.co.uk/itm/40-Isopods-Porcellionides-pruinosus-Dart-frog-Newts-and-Salamander-food-culture-/171520772125>, <https://www.insecte.org/forum/viewtopic.php?t=134763>, <https://argiope.se/ovogram/>). We suggest a series of detailed studies to take other steps towards the commercial sale of isopods to enhance soil fertility.

Acknowledgements

The authors would like to express their appreciation to Dr. Elizabeth Hornung and Dr. Diána Csonka for their comments and guidance during our research. The authors would like to thank Dr. Sándor Farkas for the revision of the text and Zita Lakiné Sasvári for her valuable help in determining the fluorescein concentration of our soil samples. We were glad to work with Krisztina Boziné Pullai (ÖMKi, PhD student, SZIU) and Renáta Petrikovszki (PhD student, SZIU). This research was partly supported by the Higher Education Institutional Excellence Program (1783-3/2018/FEKUTSTRAT) awarded by the Ministry of Human Capacities within the framework of plant breeding and plant protection researches of Szent István University, and by the ÚNKP-17-3 and ÚNKP-18-3 New National Excellence Program of the Ministry of Human Capacities

References

- Birkás M (2010): Long-term experiments aimed at improving tillage practices. *Acta Agronomica Hungarica*, 58 (Suppl.): 75-81. DOI: 10.1556/AAgr.58.2010.Suppl.1.11
- Campiglia E, Mancinelli R, Radicetti E (2011): Influence of no-tillage and organic mulching on tomato (*Solanum lycopersicum* L.) production and nitrogen use in the mediterranean environment of central Italy. *Scientia Horticulturae* 130, 588-598. DOI:10.1016/j.scienta.2011.08.012.
- Casals J, Pascual L, Canizares J, Cebolla-Cornejo J, Casanas F, Nuez F (2011): The risks of success in quality vegetable markets: Possible genetic erosion in Marmande tomatoes (*Solanum lycopersicum* L.) and consumer dissatisfaction. *Scientia Horticulturae*, 130, 78-84 DOI: 10.1016/j.scienta.2011.06.013
- Diekötter T, Wamser S, Wolters V, Birkhofer K (2010): Landscape and management effects on structure and function of soil arthropod communities in winter wheat. *Agriculture, Ecosystems & Environment* 137 (1-2), 108-112. ISSN 0167-8809, DOI: 10.1016/j.agee.2010.01.008
- Diver S, Kuepper G, Born H (1995) Organic tomato production. National Center for Appropriate Technology (ATTRA) Publication #CT073/149 NCAT IP439.
- FAOSTAT (www.fao.org/faostat/en/#home) [Retrieved 15-06-2018]
- Farkas S, Otártics MZs, Mátyás R (2017): Possible ecosystem services of terrestrial isopods. Abstract Book of the 10th International Symposium on Terrestrial Isopod Biology p. 7-8. ISBN: 978-963-87343-9-6
- Farkas S, Vilisics F (2013): Magyarország szárazföldi ászkarák faunájának határozója (Isopoda: Oniscidea), Kaposvár. *Natura Somogyiensis* 23, 89-124.
- Fehér A, Mészárosné Póss A, Turóczy Gy, Tóth F (2017) Különböző szerves talajtakaró anyagok hatása a burgonya ép gumókihozatalára, valamint károsító- és nem károsító-eredetű minőségromlására. *Növényvédelem* 53 (9): 399-404.
- Südiné F.A, Mészárosné Póss A, Tóth F, Turóczy Gy (2017): Termésnövelés és károsítók elleni védekezés burgonyában szerves talajtakaró anyagokkal. *Biokultúra* 28 (5): 14-15.
- Förster B, Muroya K, Garcia M (2006): Plant growth and microbial activity in a tropical soil amended with faecal pellets from millipedes and woodlice. *Pedobiologia* 50: 281-290. DOI:10.1016/j.pedobi.2006.03.001
- Gere G (1956): Erdei avarfogyasztó Diplopoda és Isopoda fajok humifikációs szerepének vizsgálata növénynevelési módszerrel / The role of diplopods and isopods in humification studied with plant growing method. *Állattani Közlemények* 45: 284-293.
- Gregory S (2009): Woodlice and waterlice (Isopoda: Oniscidea & Asellota) in Britain and Ireland. Field Studies Council/Centre for Ecology & Hydrology, ISBN: 10-0955767288, ISBN-13: 978-0955767289
- Hammer Ø, Harper DAT, Ryan PD (2001): PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4 (1): 4-9.
- Johnson SN, Lopaticki G, Barnett K, Facey SL (2016): An insect ecosystem engineer alleviates drought stress in plants without increasing plant susceptibility to an above-ground herbivore. *Functional Ecology* 30 (6): 894-902. DOI: 10.1111/1365-2435.12582
- Kumar R, Sood S, Sharma S, Kasana RC, Pathania VL, Singh B, Singh RD (2014): Effect of plant spacing and organic mulch on growth, yield and quality of natural sweetener plant *Stevia* and soil fertility in western Himalayas *International Journal of Plant Production* 8(3):311-334 www.researchgate.net/publication/262263478_Effect_of_plant_spacing_and_organic_mulch_on_growth_yield_and_quality_of_natural_sweetener_plant_Stevia_and_soil_fertility_in_western_Himalayas

- Lavelle P, Decaëns T, Aubert M, Barot S, Blouin M, Bureau F, Margerie P, Mora P, Rossi JP (2006): Soil invertebrates and ecosystem services. *European Journal of Soil Biology* 42: S3-S15 DOI:/10.1016/j.ejsobi.2006.10.002
- Loksa I (1988) Ászkarák In: Jermy T, Balázs K (Eds) A növényvédelmi állattan kézikönyve I. Akadémiai Kiadó, Budapest, 177–182.
- Mancinelli R, Marinari S, Brunetti P, Radicetti E, Campiglia E (2015): Organic mulching, irrigation and fertilization affect soil CO₂ emission and C storage in tomato crop in the Mediterranean environment. *Soil & Tillage Research* 152: 39-51 DOI: 10.1016/j.still.2015.04.001
- Ömki Tomato (2015) <http://www.biokutatas.hu/images/stories/kiadvanyok/paradicsom-mb-net.pdf>
- Passam HC, Karapanos IC, Bebeli PJ, Savvas D (2007): A review of recent research on tomato nutrition, breeding and post-harvest technology with reference to fruit quality. *The European Journal of Plant Science and Biotechnology*. 1 (1): 1-21.
- Philpott SM. (2013): Biodiversity and Pest Control Services. In: Levin S.A. (ed.) *Encyclopedia of Biodiversity Second Edition*, Waltham, MA: Academic Press. Volume 1, pp. 373-385.
- Schnürer J, Rosswall T (1982): Fluorescein diacetate hydrolysis as a measure of total microbial activity in soil and litter. *Applied and Environmental Microbiology*, 43(6): 1256-1261. DOI:10.1016/0038-0717(85)90036-7
- Shirgure, PS (2012): Sustainable acid lime fruit production and soil moisture conservation with different mulches. *Agricultural Engineering Today* 36 (3): 21-26. https://www.researchgate.net/publication/295504007_Sustainable_Acid_Lime_Fruit_Production_and_Soil_Moisture_Conservation_with_Different_Mulches
- Stachursky A (1968): Emigration and mortality rates and the food-shelter conditions of *Lygidium hypnorum* L. (Isopoda). *Ekologia Polska, Seria A*, 16: 445-459.
- Sutton SL, Hassall M, Willows R, Davis RC, Grundy A, Sunderland KD (1984): Life histories of terrestrial isopods: a study of intra- and interspecific variation. – *Symp. Zool. Soc. Lond.* 53: 269-294.
- Tantachasatid, P, Boyer J, Thanisawanyankura S, Séguy L, Sajjaphan K (2017): Soil macrofauna communities under plant cover in a no-till system in Thailand. *Agriculture and Natural Resources*, 51 (1): 1-6, ISSN 2452-316X, DOI: 10.1016/j.anres.2016.08.004.
- Tuf I H, Tufová J (2005): Communities of terrestrial isopods (Crustacea: Isopoda: Oniscidea) in epigeon of oak-hornbeam forests of SW Slovakia – *Ekológia* 24: 114-123.
- Vilisics F, Szekeres S, Hornung E (2012): Size dependent differences in litter consumption of isopods: preliminary results. *ZooKeys* 176, 247-259 DOI:10.3897/zookeys.176.2470
- Zimmer M, Topp W (1999): Relationship between woodlice and microbial density and activity in the field. *Biol Fertil Soils* 30: 117. DOI:10.1007/s003740050597
- Zribi W, Aragüés R, Medina E, Faci JM (2015): Efficiency of inorganic and organic mulching materials for soil evaporation control. *Soil & Tillage Research* 148: 40-45. DOI:10.1016/j.still.2014.12.003

Annexes

Annex1: Soil microbial activity

| | I-M- | I-M+ | I+M- | I+M+ | Original soil |
|---|-------------|-----------|-------------|-----------|---------------|
| Average | 0.8736 | 1.1477 | 0.9068 | 1.1473 | 1.1486 |
| Deviation | 0.075020294 | 0.1602 | 0.13805297 | 0.1943 | 0.115313293 |
| SE | 0.0237235 | 0.0506 | 0.04365618 | 0.0614 | 0.036465265 |
| CI 95% | 0.046497205 | 0.0993 | 0.08556454 | 0.1204 | 0.071470606 |
| One-way ANOVA | | | | | |
| | Sum of sqrs | df | Mean square | F | p(same) |
| Between groups: | 0.802225 | 4 | 0.200556 | 9.891 | 7.892E-06 |
| Within groups: | 0.912425 | 45 | 0.0202761 | | |
| Total: | 1.71465 | 49 | | | |
| omega^2: | 0.4156 | | | | |
| Levene's test for homogeneity of variance. based on means: p(same) = 0.7645 | | | | | |
| Based on medians: p(same) = 0.8287 | | | | | |
| Welch F test in the case of unequal variances: F=14.35, df=21.77, p=7.015E-06 | | | | | |
| Tukey's pairwise comparisons: Q below diagonal. p(same) above diagonal | | | | | |
| | I-M- | I-M+ | I+M- | I+M+ | Original soil |
| I-M- | | 0.0009404 | 0.9848 | 0.0009562 | 0.0009059 |
| I-M+ | 6.087 | | 0.004092 | 1 | 1 |
| I+M- | 0.7373 | 5.35 | | 0.004165 | 0.003932 |
| I+M+ | 6.078 | 0.008883 | 5.341 | | 1 |
| Original soil | 6.107 | 0.01999 | 5.37 | 0.02887 | |

Annex2: *Porcellionides pruinosus* and mulch

| | | | | |
|---|-----------|-----------|----------|-----------|
| Kruskal-Wallis test for equal medians | | | | |
| H (chi2): | 17.42 | | | |
| Hc (tie corrected): | 28.14 | | | |
| p (same): | 3.40E-06 | | | |
| There is a significant difference between sample medians | | | | |
| Mann-Whitney pairwise comparisons. raw p-values. uncorrected significance | | | | |
| | I-M- | I-M+ | I+M- | I+M+ |
| I-M- | | 1 | 0.1681 | 0.0002312 |
| I-M+ | 1 | | 0.1681 | 0.0002312 |
| I+M- | 0.1681 | 0.1681 | | 0.001151 |
| I+M+ | 0.0002312 | 0.0002312 | 0.001151 | |

Annex3: Maximum number of flower buds

| | Average | CI 95% |
|------|---------|-------------|
| I-M- | 7.2 | 1.998778921 |
| I-M+ | 6.3 | 2.08756356 |
| I+M- | 7.7 | 1.802267141 |
| I+M+ | 8.6 | 1.466702744 |

One-factor variance analysis
Summary

| Groups | Quantity | Sum | Average | Variance |
|--------|----------|-----|---------|-------------|
| I-M- | 10 | 72 | 7.2 | 10.4 |
| I-M+ | 10 | 63 | 6.3 | 11.34444444 |
| I+M- | 10 | 77 | 7.7 | 8.45555556 |
| I+M+ | 10 | 86 | 8.6 | 5.6 |

Variance analysis

| Factors | SS | df | MS | F | p-value | F crit. |
|----------------|-------|----|------------|-------------|-----------|----------|
| Between groups | 27.7 | 3 | 9.23333333 | 1.031657356 | 0.3901503 | 2.866266 |
| In groups | 322.2 | 36 | 8.95 | | | |
| Sum | 349.9 | 39 | | | | |

Annex4: Maximal number of leaves

| | Average | CI 95% |
|------|---------|-------------|
| I-M- | 12.1 | 1.110646258 |
| I-M+ | 11.8 | 0.703648605 |
| I+M- | 12.5 | 0.602332503 |
| I+M+ | 12.8 | 1.124016514 |

One-factor variance analysis
Summary

| Groups | Quantity | Sum | Average | Variance |
|--------|----------|-----|---------|------------|
| I-M- | 10 | 121 | 12.1 | 3.21111111 |
| I-M+ | 10 | 118 | 11.8 | 1.28888889 |
| I+M- | 10 | 125 | 12.5 | 0.94444444 |
| I+M+ | 10 | 128 | 12.8 | 3.28888889 |

Variance analysis

| Factors | SS | df | MS | F | p-value | F crit. |
|----------------|----|------|----|------------|-------------|------------|
| Between groups | | 5.8 | 3 | 1.93333333 | 0.885496183 | 0.45783582 |
| In groups | | 78.6 | 36 | 2.18333333 | | |
| Sum | | 84.4 | 39 | | | |

One-way ANOVA

| | Sum of sqrs | df | Mean square | F | p(same) |
|-----------------|-------------|----|-------------|--------|---------|
| Between groups: | 5.8 | 3 | 1.93333 | 0.8855 | 0.4578 |
| Within groups: | 78.6 | 36 | 2.18333 | | |
| Total: | 84.4 | 39 | | | |

omega²: -0.008662
Levene's test for homogeneity of variance. based on means: 0.355
p(same) =
Based on medians: p(same) = 0.5658

Welch F test in the case of unequal variances: F=1.004. df=19.41. p=0.4124

Tukey's pairwise comparisons: Q below diagonal. p(same) above diagonal

| | I-M- | I-M+ | I+M- | I+M+ |
|------|--------|-------|-------|------|
| I-M- | | | | |
| I-M+ | 0.642 | | | |
| I+M- | 0.8561 | 1.498 | | |
| I+M+ | 1.498 | 2.14 | 0.642 | |

Annex5: Maximum number of flowers

| | I-M- | I-M+ | I+M- | I+M+ | |
|--|-------------|-------------|-------------|-------------|----------------|
| Average | | 3.9 | 5.5 | 4.5 | 6.1 |
| Deviation | 1.100504935 | 1.269295518 | 1.433720878 | 1.595131482 | |
| SE | 0.348010217 | 0.401386486 | 0.45338235 | 0.504424865 | |
| CI 95% | 0.682087492 | 0.786703056 | 0.888613078 | 0.988654568 | |
| One-way ANOVA | | | | | |
| | Sum of sqrs | df | Mean square | F | p(same) |
| Between groups: | 29.2 | | 3 | 9.73333 | 5.246 0.004167 |
| Within groups: | 66.8 | | 36 | 1.85556 | |
| Total: | 96 | | 39 | | |
| omega^2: | 0.2415 | | | | |
| Levene's test for homogeneity of variance. | | | | | |
| based on means: p(same) = | 0.6626 | | | | |
| Based on medians: p(same) = | 0.7355 | | | | |
| | F=5.246. | | | | |
| Welch F test in the case of unequal variances:df=19.83. | | | | | |
| | p=0.007889 | | | | |
| Tukey's pairwise comparisons: Q below diagonal. p(same) above diagonal | | | | | |
| | I-M- | I-M+ | I+M- | I+M+ | |
| I-M- | | | 0.0583 | 0.7589 | 0.004985 |
| I-M+ | 3.714 | | | 0.3691 | 0.7589 |
| I+M- | 1.393 | 2.321 | | | 0.0583 |
| I+M+ | 5.107 | 1.393 | 3.714 | | |

Annex6: Tomato Yield

| | I-M- | I-M+ | I+M- | I+M+ | |
|---|-------------|-------------|-------------|-------------|---------------|
| Average | | 149.69 | 164.88 | 209.9 | 228.94 |
| Deviation | 44.67739796 | 75.84869149 | 46.58826748 | 44.82772456 | |
| SE | 14.12823375 | 23.98546226 | 15.05652313 | 14.17577119 | |
| CI 95% | 27.69082931 | 47.01064219 | 29.51024308 | 27.78400099 | |
| One-way ANOVA | | | | | |
| | Sum of sqrs | df | Mean square | F | p(same) |
| Between groups: | 41573.9 | | 3 | 13858 | 4.647 0.00759 |
| Within groups: | 107362 | | 36 | 2982.27 | |
| Total: | 148936 | | 39 | | |
| omega^2: | 0.2148 | | | | |
| Levene's test for homogeneity of variance. | | | | | |
| based on means: p(same) = | 0.006713 | | | | |
| Based on medians: p(same) = | 0.009981 | | | | |
| Welch F test in the case of unequal variances: F=5.78. df=19.76. p=0.005236 | | | | | |
| Tukey's pairwise comparisons: Q below diagonal. p(same) above diagonal | | | | | |
| | I-M- | I-M+ | I+M- | I+M+ | |
| I-M- | | | 0.9244 | 0.08299 | 0.01304 |
| I-M+ | 0.8796 | | | 0.2704 | 0.05875 |
| I+M- | 3.487 | 2.607 | | | 0.8632 |
| I+M+ | 4.589 | 3.709 | 1.103 | | |

Yield components of winter oilseed rape regard to plant population

Elizabeta ANĐELIĆ – Manda ANTUNOVIĆ – Miro STOŠIĆ – Dario ILJKIĆ – Ivana VARGA

Faculty of Agrobiotechnical Sciences Osijek, Josip Juraj Strossmayer University of Osijek, Vladimira Preloga 1, 31000 Osijek; E-mail: ivana.varga@pfos.hr

Abstract: The aim of this study was to analyze the yield components of winter oilseed rape depending on the plant population in 2017/2018 growing season. Three plant populations were considered in the study: 20, 40 and 60 plants m^{-2} . At harvest several yield components were determined: plant height (cm), plant mass (g), height of the first fertile branch (cm), number of fertile lateral branches, number of pods per plant, length of the pod (cm), number of seeds per pod, mass of seeds per pod (g), number of seeds per plant, 1000 grain mass (g) and yield of seeds per plant (g). At the plant population of 40 plants m^{-2} the plants were the highest (153.4 cm), with the highest plant mass (295.3 g) and the number of lateral fertile branches (5.6 branches per plant). Furthermore, the plants from 40 plants m^{-2} had the highest number of pods per plant (716 pieces), the longest pods (6.5 cm) with the largest number of seeds per pod (21.0) and the number of seeds per plant (15 036 seeds). The highest and significant correlation coefficient was found between number of pods per plant and plant mass ($r=0.890^{***}$) and than between the number of lateral branches and number of pods per plant ($r=0.850^{***}$). Linear regression showed that for every centimeter increment of pod length the number of seeds increase for 4.8 seeds at 20 and 40 plants m^{-2} and for 5.5 seeds per pod at 60 plants m^{-2} . Furthermore, regression analysis showed that for every centimeter pod length increment the seeds mass per pod increase on average of all plant population for 0.02 g.

Keywords: winter oilseed rape, 2017/2018, plant population, yield components

Received 18 October 2018, Revised 13 December 2018, Accepted 17 December 2018

Introduction

According to FAOStat data (2018) in the world production of oilseed rape from 2012 to 2016 was on about 35 mil. ha with seed yield of 2.0 t ha^{-1} , whereas in the Europe there was about 25% of world sown area, and it counts on average 8.6 mil. ha with seed of yield 2.9 t ha^{-1} (Table 1). The greater producers in the EU are France, Germany and United Kingdom, while in the world great producers are Canada, China, India and Australia. In Republic of Croatia the average

Table 1. Production of oilseed rape in the world from 2012 to 2016 (FAOStat, 2018)

| | Harvested area (ha) | Seed yield (t ha^{-1}) |
|--|---------------------|---------------------------|
| The greater producers in the Europe | | |
| <i>Europe total</i> | 8 619 138 | 2.9 |
| France | 1 520 668 | 3.3 |
| Germany | 1 355 440 | 3.9 |
| Russian Federation | 993 076 | 1.1 |
| Poland | 873 228 | 2.9 |
| Ukraine | 705 758 | 2.5 |
| Great Britain | 675 303 | 3.4 |
| The greater producers in the world | | |
| <i>World total</i> | 35 175 387 | 2.0 |
| Canada | 8 322 800 | 2.1 |
| China | 7 537 637 | 1.9 |
| India | 6 085 748 | 1.2 |
| Australia | 2 726 891 | 1.3 |

production area of winter oilseed rape was 21 948 ha and seed yield 2.8 t ha^{-1} (Statistical Yearbook of the Republic of Croatia 2017). The importance of rape has thus increased in recent years and today it is cultivated on every continent. Nowadays the cultivation of oilseed rape have a great importance due to its usage as vegetable oil in human diet, but also for the biodiesel production as a renewable source, as a catch crop for green manuring and as a forage crop: nutrition in the form of rape cake and meal (Pepó, 2013; Carré and Pouzet, 2014; Lääniste et al. 2016; Nath et al. 2016; Zając et al., 2016; Novoselec et al. 2018).

The number of plants per unit area is one of the most important yield components in plant production (Ma et al., 2014; Masarirambi et al., 2012; Pepó and Murányi, 2014; Varga et al., 2015; Li et al., 2017; Varga et al., 2017; Vinze, 2017). According to Balodis and Gaile (2016) plant population in relation to the sowing rate particularly affects the number of pods per plant and seed number per pod. The optimum density or plant population results in mature plants that are sufficiently crowded to efficiently use resources such as water, nutrients, and sunlight, yet not so crowded that some plants die or are unproductive. Thus, the distribution of plants

per unit area is of great importance for yield stability. There are several factors that have influence on the realized plant population like soil properties, seed quality, field germination, sowing time, plant morphology, diseases, pests, seedbed preparation etc. (Sidlauskas et al., 2003; Kristek et al., 2015; Balodis and Gaile 2016; Kovacevic et al., 2017; Zebec et al., 2017).

The optimum time for sowing winter oilseed rape in Croatia is from 25th August to 10th September. It is therefore sown at 12.5 cm or 25 cm distance between the rows and at the depth from 1.5 cm to 2.5 cm. The recommended seed rate for winter sown oilseed rape in the Croatia is 2.5 – 5 kg ha⁻¹. In Europe oilseed rape is usually sown as winter crop at sown rate around 70 seeds m⁻² (Roques and Berry, 2016). It can be also sown as a spring crop, when it is sown in higher density as 150 seeds m⁻² (Lääniste et al. 2016) or 100 – 110 seeds m⁻² (Leach et al., 1999) for hybrids.

The optimum plant population for oilseed rape hybrids is 30 – 50 plants m⁻² at harvest (Pospišil, 2013.). On the one hand, in lower plant populations oilseed rape form greater lateral branches, which can somewhat compensate the lack of plants. On the other hand, larger number of plants per m⁻² can cause decrease in the stem diameter, thus plants are more sensible to lie down. Leach et al. (1999) indicate that the yield of winter oilseed rape increases if 50-60 plants m⁻² is achieved. Authors suggested that the productivity of the plant at lower plant population is compensated by increasing the leaf area, multiple lateral fertile branches and with more pods per plant. On the contrary, authors stated that in large plant population, no significant increase in yields was found because of the greater possibility of disease in oilseed rape crops. Mendham et al. (1981) shown that plant populations of 20–30 plants m⁻² produce yields comparable to crops with 70–80 plants m⁻², in some instances a crop with only 9–10 plants m⁻² produced acceptable yields if the plants were healthy and evenly distributed. Diepenbrock (2000) reported that plant population had the greatest effect on yield components of individual plants.

There is a lack of research available on winter oilseed rape yield components in the conditions of Croatia, especially regard to plant population. It is therefore, essential to understand how individual plants interact with each other and

the environment and to possibly come up with the ideal crop density levels to optimize yields. The aim of this study was to determine the winter oilseed rape yield components regard to different plant populations in 2017/2018 vegetation.

Material and methods

Field trial was set up in Eastern Croatia (location Beravci) on Family farm «Ivica Anđelić». Winter oilseed rape was sown on 5th September 2017 (hybrid Hybrirock, KWS). One month after sowing (5th November 2017) when plants had about 5-6 leaves, the plant population correction was made to get three different densities: 20, 40 and 60 plants m⁻². Each plant population was established in plot size 10 m x 3 m. There were total 171 kg ha⁻¹ N, 67 kg ha⁻¹ P i 87 kg ha⁻¹ K added with fertilization (Table 2). All protection against weeds and pests was made properly.

During the vegetation, the number of leaves per plant were determined in 7 dates (from 5th November 2017 to 5th April 2018). In 2017 the number of leaves represent the average number of leaves per plant counted on 30 individual plants, whereas in 2018 the number of leaves per plant represent the average number of 45 individual plants. Thus, the number of leaves were determined on 270 individual plants.

Besides leaves counting, during vegetation the realized plant population was determined three times for every treatment in 4 replications. The harvest was on 11th June 2018 and therefore from the every plant population (20, 40 and 60 plants m⁻²) the 15 individual plants were harvested manually. Therefore, the yield components were

Table 2. Fertilization of winter oilseed rape for 2017/2018 vegetation year

| Fertilizer | Amount (kg ha ⁻¹) | N | P ₂ O ₅ | K ₂ O |
|--|-------------------------------|-----------|-------------------------------|------------------|
| 5 th July 2017 (with soil tillage) | | | | |
| NPK 0-20-30 | 200 | - | 40 | 60 |
| UREA | 100 | 46 | - | - |
| 5 th September 2017 (with sowing) | | | | |
| NPK 15-15-15 | 180 | 27 | 27 | 27 |
| 1 st March and 12 th April 2018 (top dressing) | | | | |
| KAN | 180 | 49 | - | - |
| KAN | 180 | 49 | - | - |
| Total | 171 | 67 | 87 | |




Figure 1. The mean air temperature (°C) in 2017/2018 winter oilseed rape vegetation as compared to the long term mean (LTM = 1961-1990) for Meteorological station Slavonski Brod (Croatian Meteorological and Hydrological Service, 2018)




Figure 2. The total monthly precipitation (mm) in 2017/2018 winter oilseed rape vegetation as compared to the long term mean (LTM = 1961-1990) for Meteorological station Slavonski Brod (Croatian Meteorological and Hydrological Service, 2018)

determined on 45 individual plants. The plants were marked and afterwards the plant height (cm), the plant height to first fertile branch (cm), total number of pods per plant were measured. From every plant 100 individual pods were separated in the paper bag, and than on every pod several measurements were made: the pods length (cm), the number of seeds per pod and mass of seeds per pod. Total of 4500 individual pods were analyzed. Afterwards, the number of seeds per plant, the 1000 grain mass and the yield of seeds per plant (g per plant) were determined.

During vegetation period, from August 2017 to June 2018 (Figure 1) the average air temperature was 1.9°C higher as compared to the long-term mean (LTM, 1961-1990) (Croatian Meteorological and Hydrological Service, 2018). The winter period was not so cold, so the plants were not destroyed. Even though, in January the mean air temperature was for 4.5°C which was higher for 5.7°C comparing to LTM. During vegetation, there were total

824.1 mm precipitation (Figure 2), which was for 156.8 mm higher as compared to the LTM (667.3 mm). In spring 2018 there was 173.8% higher rainfall in March, and than the lack of rainfall in April (17.7 mm), but towards to winter oilseed rape maturation, in May and June there were higher rainfall and warm air temperatures, thus the harvest of winter oil seed rape and other winter crops in Croatia started about 20 days earlier than usual.

The differences between the mean values were calculated in SAS 9.4. with one way ANOVA as plant population as the main factor.

Results

In order to determine the phase of growth for winter period, the number of leaves per plant were determined (Figure 3). Thus, the rosette was in the optimum phase for winter period. Also after winter period there was not extreme decrease of planed plant population (Table 3). The mean plant height was 138.8 cm (Table 4)




Figure 3. The average number of leaves per oilseed rape plant in 2017/2018 vegetation

Table 3. Realized winter oilseed rape plant population

| Planned plant population | Date | | | | | |
|---------------------------|---------------------------------|-------------|---------------------------------|-------------|---------------------------------|-------------|
| | 5 th November 2017 | | 5 th March 2018 | | 11 th June 2018 | |
| | Realized plants m ⁻² | % of planed | Realized plants m ⁻² | % of planed | Realized plants m ⁻² | % of planed |
| 20 plants m ⁻² | 19 | 95 | 18 | 90 | 18 | 90 |
| 40 plants m ⁻² | 39 | 98 | 39 | 98 | 39 | 98 |
| 60 plants m ⁻² | 59 | 98 | 58 | 97 | 57 | 95 |

Table 4. The plant height (cm), plant mass (g), height to first fertile branches (cm), the number of lateral branches of winter oilseed rape at different plant population

| | Plant height (cm) | Plant mass (g) | Hight to first fertile branches (cm) | Number of lateral branches |
|---------------------------|--------------------|--------------------|--------------------------------------|----------------------------|
| 20 plants m ⁻² | 123.7 ^c | 112.0 ^b | 28.4 ^a | 3.5 ^b |
| 40 plants m ⁻² | 153.4 ^a | 295.3 ^a | 21.0 ^b | 5.6 ^a |
| 60 plants m ⁻² | 139.1 ^b | 167.9 ^b | 19.1 ^b | 4.4 ^b |
| Average | 138.8 | 191.8 | 22.9 | 4.5 |

The differences between the means within the column are marked with different letter (abc) at $p < 0.05$

Table 5. The number of pods per plant, the pods length (cm), number of seeds per pod and mass of seed per pod (g) of winter oilseed rape at different plant population

| | Number of pods per plant | Pods length (cm) | Number of seeds per pod | Mass of seed per pod (g) |
|---------------------------|--------------------------|-------------------|-------------------------|--------------------------|
| 20 plants m ⁻² | 339 ^b | 6.2 ^b | 20.4 ^a | 0.09 ^a |
| 40 plants m ⁻² | 716 ^a | 6.5 ^a | 21.0 ^a | 0.10 ^a |
| 60 plants m ⁻² | 462 ^b | 6.4 ^{ab} | 20.3 ^a | 0.10 ^a |
| Average | 506 | 6.4 | 20.6 | 0.10 |

The differences between the means within the column are marked with different letter (abc) at $p < 0.05$

Table 6. The number of seeds per plant, 1000 grain mass (g) and yield of seeds per plant (g) of winter oilseed rape at different plant population

| | Number of seeds per plant | 1000 grain mass (g) | Seed yield per plant (g) |
|---------------------------|---------------------------|---------------------|--------------------------|
| 20 plants m ⁻² | 6 916 ^b | 4.71 ^a | 35.0 ^b |
| 40 plants m ⁻² | 15 036 ^a | 4.65 ^a | 85.2 ^a |
| 60 plants m ⁻² | 9 379 ^b | 4.85 ^a | 45.0 ^b |
| Average | 10 423 | 4.74 | 55.1 |

The differences between the means within the column are marked with different letter (abc) at $p < 0.05$

and it varied from 123.7 cm (20 plants m⁻²) to 153.4 cm (40 plants m⁻²). The highest plant mass (295.3 g) had the plants at 40 plants m⁻², and there were no statistically differences determined for plant mass between 20 and 60 plants m⁻². The plants formed 4.5 lateral branches and the mean height of first fertile branch was 22.9 cm.

At plant population of 40 plants m⁻² plants formed the higher number of pods per plant (Table 5), but also, at 40 plants m⁻² the number of seeds per plant and seed yield per plant (Table 6) were the highest, which was significant ($p < 0.05$) in comparison with number of pods per plant, number of seeds per plant and seed yield per plant at 20 and 60 plants m⁻² where the differences

was not significant. The average length of pods was 6.4 cm with 20.6 seeds which weighted on average 0.10 g (Table 5). The 1000 grains mass was not significant different between plant population.

In order to determine the relationship between the yield components the Pearson's correlation coefficient was calculated (Table 7). Significant and positive correlation for plant population and plant height was found ($r = 0.385^{**}$) and significant but negative correlation was between plant population and the height to first fertile branch ($r = -0.372^*$). Other yield components (the number of lateral branches, the number of pods per plant, plant mass, pod length, number of seeds per pod, mass of seeds per pod and

Table 7. The Pearson's correlation coefficient between yield components of winter oilseed rape (N = 45)

| | PP | PH | BH | NB | NP | PM | PL | NSP | MS | PS |
|-----|----------|----------|-----------|----------|----------|----------|----------|----------|----------|----|
| PP | - | | | | | | | | | |
| PH | 0.385** | - | | | | | | | | |
| BH | -0.372* | -0.300* | - | | | | | | | |
| NB | 0.229ns | 0.625*** | -0.465*** | - | | | | | | |
| NP | 0.156ns | 0.662*** | -0.481*** | 0.850*** | - | | | | | |
| PM | 0.179ns | 0.666*** | -0.467*** | 0.782*** | 0.890*** | - | | | | |
| PL | 0.246ns | 0.500*** | -0.378** | 0.561*** | 0.508*** | 0.415** | - | | | |
| NSP | 0.001ns | 0.247ns | -0.251ns | 0.296* | 0.335* | 0.199ns | 0.755*** | - | | |
| MS | -0.075ns | 0.368* | -0.369* | 0.542*** | 0.619*** | 0.631*** | 0.448** | 0.481*** | - | |
| PS | -0.073ns | 0.373* | -0.368* | 0.545*** | 0.619*** | 0.631*** | 0.454** | 0.488*** | 0.757*** | - |

p < 0.001 ***; p < 0.01 **; p < 0.05 *; ns – not significant)

PP – plant population; PH – plant height; BH – height to first fertile branch; NB – number of lateral branches; NP – number of pods per plant; PM – plant mass; PL – pod length; NSP – number of seeds per pod; MS – mass of seeds per pod; SY – seed yield per plant

seed yield per plant) did not have significant relationship with winter oilseed rape plant population. For other winter oilseed rape yield components, the highest and significant correlation coefficient was found between number of pods per plant and plant mass ($r = 0.890^{***}$) and than between the number of lateral branches and number of pods per plant ($r = 0.850^{***}$).

Linear regression was also calculated for pod length and number of seeds per pod and pod length and mass of seeds per pod (Figure 4). According to linear regression analysis the equations showed quite similar relationship between the pod length and number of seeds per pod at every plant population. Thus, it was found that for every centimeter increment of pod length the number of seeds increase for about 4.8 seeds at 20 and 40 plants m^{-2} and for about 5.5 seeds per pod at 60 plants m^{-2} . In regression analysis of the mass of seeds per pod (g) and pod length (cm) it was found that for every centimeter pod length increment the mass of seeds per pod increase for 0.02 or 0.03 g.

Discussion

In the period from August 2017 to June 2018, temperatures were higher by 1.9°C than the long term mean (9.6°C) and during this period the precipitation amount was 156.8 mm higher than the long term mean (667.3 mm). The weather conditions were suitable for the development of

oilseed rape. According to the data obtained in this study, the planned plant population was not drastically reduced to harvest. It is important to note that in the winter period from December 2017 to March 2018 was not extremely cold and did not affect the plant's population reduction. Zając et al. (2013) state that as a result of very cold winter, oilseed rape plants developed a larger number of lateral branches to compensate for reduced plant population. In this study, high temperatures in April and May led to earlier harvest of oilseed rape, which was about 20 days before the optimal harvest dates (3rd decade of the June).

The plant population is a very important factor winter oilseed rape production. Pospíšil et al. (2014) and Diepenbrock (2000) stated that the plant population had the greatest influence on the seed yield and the yield component of winter oilseed rape. Zhang et al. (2012) found that the number of pods per plant and the number of seeds per pod are the most varied yield component of winter of oilseed rape.

The basic yield components of oilseed rape seed are the number of plants per unit area (m^2), the number of pods per plant, the number of seeds per pod and the mass of 1000 seeds. Even though Momoh and Zhou (2001) and Clarke and Simpson (1978) found that in small plant population oilseed rape will increase the number of lateral branches, in this study on




Figure 4. The linear regression of pod length (cm) and number of seeds per pod and pod length (cm) and mass of seeds per pod (g) of winter oilseed rape at three different plant population

the contrary, the smallest plant population (20 plants m⁻²) did not result with highest number of lateral branches (3.5 lateral branches). In our study some relationships between several yield components (parameters of an individual plant productivity) were observed. The number of pods per plant depends on the number of developed pods and aborted pods, which vary depending on different pollination conditions. Abiotic stress like high temperatures during flowering can also affect the number of pods per plant. Angadi et al. (2000) reported that

high temperatures at flowering affected yield formation more than high temperature at pod development and that in such conditions *Brassica* sp. could have more pods per plant, but with less seeds due to seed abortion was also more common. According to Pospíšil (2013) the number of pods per plant in our agroecological conditions could vary from 100 to 600. In our experiment the highest number of pods per plant was 716 in plant population of 40 plants m⁻² and at 20 and 60 plants m⁻² it was decreased by 52.7% and 35.5%, respectively. Balodis and

Gaile (2016) stated that the differences in the oilseed rape yield component “pods per plant” showed plant ability to compensate seed yield in cases when plant density was lower, such as due to decreased sowing rate and poor wintering. So, in our experiment this was not confirmed. Based on three year experiment in Croatia, Pospišil et al. (2014) found that the number of pods per plant varied from 140 (cultivar Ricco, 2009/10 season) to 528 (hybrid Turan, 2011/12 season).

Correlations are important for the breeder in order to associate all the possible valuable features in the newly created genotypes. It was interesting to find that plant height was positive and extremely significant correlated with the number of lateral fertile branches, number of seeds per plant and pod length (Table 7). Zhang and Zhou (2006) reported that number of seed per pod and 1000 seed mass were positively correlated with seed yield per plant. Besides correlations, the regression analysis can explain prediction and connection between the yield components. Zajac et al. (2011) found a high coefficient of determination $R^2 = 0.945$ between the seed mass per oilseed rape pod and the total mass of the pod, but also authors found that the length of the pod was very poorly correlated with seed mass. This is quite similar to our results, which showed a small coefficient of determination ($R^2 = 0.32$ to $R^2 = 0.35$) between pods length and seeds mass per pod.

Zajac et al. (2011) reported that mean number of seeds per pod is around 19 and that it should be considered that location of particular pod on a plant was also important; more seeds per pod were found in the middle part of plant in comparison with upper and lower parts. Authors stated that the seed number per pod can increase up to 27 seeds on the main stem, in relation to pod location on the plant. Even though it would be interesting, in our study, the location of pods on a plant was not observed, but there were no significant difference for average number (20.6) of seeds per pod regard to plant population (Table 5). The opposite findings were in Li et al. (2017) study, where with increasing winter oilseed rape plant population, decreased number of seeds per pod. Pospišil et al. (2014) emphasize the genotype differences in number of seeds per pod and between 11 hybrids and 5 cultivars, the cultivar Ricco had the highest number of seeds per pod in two seasons, 30,00 in 2010/11 and 33,41 in 2011/12.

Our study did not show the significant influence of plant population on 1000 grain mass. This is similar to Li et al. (2017) found that with increasing winter oilseed rape plant population there were no significant influence on 1000 grain mass. In study with different nitrogen fertilizer rate, Spitek and Pospišil (2017) did not find significant difference in 1000 grain mass (on average 5.65 g).

Our trial results showed the highest significant ($p < 0.05$) seed yield per plant of 85.2 g at 40 plants m^{-2} in comparison with other plant populations (Table 6). Similar results were obtained in the study of Nasiri et al. (2017). Authors include 6 different oilseed rape genotypes (Ahmadi, Okapi, Opera, L72, Karajl and SW102) and found that the seed yield at 60 and 80 plants m^{-2} in all varieties was significantly reduced compared to the 40 plants m^{-2} . In the study of different plant population (30, 40, 50 and 60 seeds m^{-2}) by Ratajczak et al. (2017) indicate that the lowest seed yield was obtained at 30 m^{-2} seeds (38.9 dt ha^{-1}), while there was no statistically justified differences between yield of seeds of other sowing density and the yield ranged from 41.0 dt ha^{-1} (40 seeds m^{-2}) to 41.4 dt ha^{-1} (60 seeds m^{-2}). Zhang et al. (2012) also found yield decrement with increasing plant population and they stated that the highest seed yield per plant was at 36 plants m^{-2} .

Conclusion

In this study field trial was set up to analyze the yield components of winter oilseed rape depending on the plant population (20, 40 and 60 plants m^{-2}) in 2017/2018 year. It is obvious in our experiment that, generally the best results were found at 40 plants m^{-2} . Thus, there were the highest plants developed (153.4 cm per plant) with the largest plant mass (295.3 g per plant) and furthermore, plants formed the most fertile side branches (5.6 branches per plant) and the largest number of pods per plant (716 pieces). Also, plants had the longest pods length (6.5 cm) and therefore the largest number of seeds per pod (average 21.0). The highest seed yield per plant of 85.2 g was found at 40 plants m^{-2} , which was 58.9% and 47.2% higher compared to 20 and 60 plants m^{-2} , respectively. Even though this results showed only one oilseed rape season, our results coincides with the recommendations that the winter oilseed rape hybrids should be sown within 30 to 50 plants m^{-2} .

References

- Angadi, S.V., Cutforth, H.W., Miller, P.R., McConkey, B.G., Entz, M.H., Brandt, S.A., Volkmar, K.M. (2000): Response of three *Brassica* species to high temperature stress during reproductive growth. *Canadian Journal of Plant Science*, **80**: 4. 693-701. <https://doi.org/10.4141/P99-152>
- Balodis, O., Gaile, Z. (2016): Sowing date and rate effect on winter oilseed rape (*Brassica napus* L.) yield components' formation. *Proceedings of the Latvian Academy of Sciences. Section B*, **70**: 6 (705). 384–392. doi: 10.1515/prolas-2016-0058
- Carré, P., Pouzet, A. (2014): Rapeseed – Tremendous potential for added value generation? *Oilseeds and fats, Crops and Lipids*, **21**: 1. D102: 1-12. doi:10.1038/srep12574
- Clarke, J.M., Simpson, G.M. (1978): Influence of irrigation and seeding rates on yield and yield components of *Brassica napus* cv. Tower. *Can. J. Plant Sci.* **58**: 731-737. <https://doi.org/10.4141/cjps78-108>
- Croatian Meteorological and Hydrological Service, 2018 (<http://meteo.hr/index.php>, accessed on 11th October 2018).
- Diepenbrock, W. (2000): Yield analysis of winter oilseed rape (*Brassica napus* L.): A review. *Field Crops Research*, **67**: 35-49. doi: 10.1016/S0378-4290(00)00082-4
- FAOStat, Food and Agriculture Organization of the United Nations (2017) <http://www.fao.org/faostat/en/#data/QC> (accessed on 10th May 2018).
- Kovacevic, V., Antunovic, M., Varga, I., Iljkic, D., Jovic, J. (2017): Response of soybean and barley to Fertdolomite application on acid soil. *Columella – Journal of Agricultural and Environmental Sciences*. **4**: 2. 7-15. DOI: 10.18380/SZIE.COLUM.2017.4.2.7
- Kristek, A., Kristek, S., Varga, I., Drmić, Z. (2015): Results of sugar beet production depending on the hybrids selection and the number of fungicide application (*in Croatian*). *Poljoprivreda/Agriculture*, **21**: 2. 15-22. DOI: <http://dx.doi.org/10.18047/poljo.21.2.3>
- Lääniste, P., Eremeev, V., Mäeorg, E., Jõudu, J. (2016): Effect of sowing date on oil, protein and glucosinolate concentration of winter oilseed rape (*Brassica napus* L.). *Agronomy Research* **14**(S2). 1384–1395.
- Leach, J. E., Stevenson, H. J., Rainbow, A. J., Mullen, L. A. (1999): Effects of high plant populations on the growth and yield of winter oilseed rape (*Brassica napus*). *The Journal of Agricultural Science*, **132**: 2. 173-180.
- Li, M., Shahbaz Naeem, M., Ali, S., Zhang, L., Liu, L., Ma, N., Zhang C. (2017): Leaf Senescence, Root Morphology, and Seed Yield of Winter Oilseed Rape (*Brassica napus* L.) at Varying Plant Densities. *BioMed Research International*, 2017, Article ID 8581072. <https://doi.org/10.1155/2017/8581072>
- Ma, N., Yuan, J., Li, M., Li, J., Zhang, L., Liu, L., Naeem, M.S., Zhang, C. (2014): Ideotype population exploration: growth, photosynthesis, and yield components at different planting densities in winter oilseed rape (*Brassica napus* L.). *PloS one*, **9**: 12: e114232. doi: 10.1371/journal.pone.0114232.
- Masarirambi, M.T., Mandisodza, F.C., Mashingaidze, A.B., Bhebhe, E. (2012): Influence of plant population and seed tuber size on growth and yield components of potato (*Solanum tuberosum*). *Int. J. Agric. Biol.*, **14**: 545–549.
- Mendham, N., Shipway, P., Scott, R. (1981): The effects of seed size, autumn nitrogen and plant population density on the response to delayed sowing in winter oil-seed rape (*Brassica napus*). *The Journal of Agricultural Science*, **96**: 2. 417-428. <https://doi.org/10.1017/S0021859600066181>
- Momoh, E.J.J., Zhou, W. (2001): Growth and yield responses to plant density and stage of transplanting in winter oilseed rape (*Brassica napus* L.). *Journal of Agronomy and Crop Science*, **186**: 4. 253-259. <https://doi.org/10.1046/j.1439-037x.2001.00476.x>
- Nasiri, A., Samdaliri, M., Rad, A. S., Shahsavari, N., Kale, A. M., Jabbari, H. (2017): Effect of plant density on yield and physiological characteristics of six canola cultivars. *Journal of Scientific Agriculture*, **1**. 249-253. DOI: <https://doi.org/10.25081/jsa.2017.v1.819>
- Nath, U.K., Kim, H.-T., Khatun, K., Park, J.-I., Kang, K.-K., Nou, I-S. (2016): Modification of Fatty Acid Profiles of Rapeseed (*Brassica napus* L.) Oil for Using as Food, Industrial Feed-Stock and Biodiesel. *Plant Breed. Biotech*, **4**: 2. 123-134. DOI: <https://doi.org/10.9787/PBB.2016.4.2.123>
- Novoselec, J., Klir, Ž., Domaćinović, M., Lončarić, Z., Antunović, Z. (2018): Biofortification of feedstuffs with microelements in animal nutrition. *Poljoprivreda/Agriculture*, **24**: 1. 25-34. DOI: 10.18047/poljo.24.1.4

- Pepó, P. (2013): Effects of nutrient supply and sowing time on the yield and pathological traits of winter oilseed rape. *Acta Agronomica Hungarica*, **61**: 3. 195–205.
- Pepó, P., Murányi, E. (2014): Plant density impact on grain yield of maize (*Zea mays* L.) hybrids on chernozem soil of the Eastern Hungary. *Columella - Journal of Agricultural and Environmental Sciences*, **1**: 2. 95-100. DOI: 10.1556/AAgr.61.2013.3.3
- Pospišil, M. (2013): Ratarstvo II. dio – industrijsko bilje. *Zrinski. Čakovec*.
- Pospišil, M., Brčić, M., Pospišil, A., Butorac, J. (2014): Yield and yield components of investigated rapeseed hybrids and cultivars (*in Croatian*). *Poljoprivreda/Agriculture*, **20**: 1. 3-9.
- Ratajczak, K., Sulewska, H., Szymańska, G. (2017): New winter oilseed rape varieties – seed quality and morphological traits depending on sowing date and rate. *Plant Production Science*, **20**: 3. 262-272. <https://doi.org/10.1080/1343943X.2017.1304809>
- Roques, S., Berry, P. (2016): The yield response of oilseed rape to plant population density. *The Journal of Agricultural Science*, **154**: 2. 305-320. <https://doi.org/10.1017/S0021859614001373>
- Sidlauskas, G., Bernotas, S. (2003): Some factors affecting seed yield of spring oilseed rape (*Brassica napus* L.). *Agronomy Research*, **1**: 2. 229-243.
- Spitek, J., Pospišil, M. (2017): Influence of nitrogen and sulfur rate in topdressing upon the yield and yield components of rapeseed (*in Croatian*). *Glasnik Zaštite Bilja*, **40**: 4. 76-81.
- Statistical Yearbook of the Republic of Croatia 2017.
- Varga, I., Kristek, A., Antunović, M. (2015): Growth analysis of sugar beet in different sowing density during vegetation. *Poljoprivreda/Agriculture*, **21**: 1: 28-34. <https://doi.org/10.18047/poljo.21.1.5>
- Varga, I., Antunović, M., Iljkić, D. (2017): Vliv rozdílného hnojení dusíkem na růst kořene cukrové řepy (*in Czech*). *Listy cukrov. řepář.*, **133**: 4. 138-141.
- Vinze, É. (2017): The effect of sowing date and plant density on yield elements of different winter oilseed rape (*Brassica napus* var. *napus* f. *biennis* L.) genotypes. *Columella – Journal of Agricultural and Environmental Sciences*, **4**: 1. 21-25. DOI: 10.18380/SZIE.COLUM.2017.4.1.suppl
- Zajac, T., Oleksy, A., Synowiec, A., Klimek-Kopyra, A. (2011). Comparison of morphological traits, productivity and canopy architecture of winter oilseed rape (*Brassica napus* L.) and white mustard (*Sinapis alba* L.). *Journal of Applied Botany and Food Quality*, **84**. 183-191.
- Zajac, T., Kulig, B., Oleksy, A., Stokłosa, A., Styrc, N., Pyziak, K. (2013): Development and yield of morphologically different groups of winter oilseed rape canopy I. Productivity and morphology of plants. *Acta Sci. Pol., Agricultura*, **12**: 1. 45-56.
- Zajac, T., Klimek-Kopyra, A., Oleksy, A., Lorenc-Kozik, A., Ratajczak, K. (2016): Analysis of yield and plant traits of oilseed rape (*Brassica napus* L.) cultivated in temperate region in light of the possibilities of sowing in arid areas. DOI: 10.5586/aa.1696
- Zebec, V., Semialjac, Z., Marković, M., Tadić, V., Radić, D., Rastija, D. (2017): Influence of physical and chemical properties of different soil types on optimal soil moisture for tillage (*in Croatian*). *Poljoprivreda/Agriculture*, **23**: 2. 10-18. DOI: 10.18047/poljo.23.2.2
- Zhang, G., Zou, W. (2006): Genetic analyses of agronomic and seed quality traits of synthetic oilseed *Brassica napus* produced from interspecific hybridization of *B. campestris* and *B. oleracea*. *J Genet.*, **85**: 1. 45-51.
- Zhang, S., Liao, X., Zhang, C., Xu, H. (2012): Influences of plant density on the seed yield and oil content of winter oilseed rape (*Brassica napus* L.). *Industrial Crops and Products*, **40**: 27-32.

Calculation nitrogen and sodium budget from lysimeter-grown short-rotation willow coppice experiment

Ágnes KUN¹ – Csaba BOZÁN¹ – Mária ONCSIK BÍRÓNÉ¹ – Károly BARTA²

1: National Agricultural Research and Innovation Centre, Research Institute of Irrigation and Water Management (NARIC ÖVKI), Szarvas, Hungary 5540 Szarvas Anna-liget utca 35.,
E-mail: kun.agnes@ovki.naik.hu

2: University of Szeged, Department of Physical Geography and Geoinformatics, Szeged, Hungary

Abstract: Alternative water resources utilization should take priority over the conventional irrigation water resources (surface and subsurface waters) in the future in Hungary as well, similarly to the global trends because of the climate change impacts. However because of the environmental risks (e.g. nitrate leaching and soil salinization) the reuse of the wastewater require sustainable practices hence the farmers and researchers are responsible for development soil management practices and irrigation principles. Aim of our study was to determine the impact of a wastewater originated from an African intensive catfish farm on the nitrogen budget of the soil-water-plant system in order to evaluate the nitrogen substitution effect and the risk of the nitrate leaching. On the other hand, the aim of the study was to calculate also sodium budget to assess the risk of the sodicity regard to the high sodium concentration of the wastewater. The experiment was conducted at the National Agricultural Research and Innovation Centre (NAIK), Research Institute of Irrigation and Water Management (OVKI) in Szarvas, Hungary. The experiment was set up in the NAIK ÖVKI Lysimeter Station in 2014 with energy willow. During the study (between 03.07.2015 and 21.04.2017) irrigation water quality, rainwater quality, willow N and Na uptake by stems and N and Na leaching was measured. Nitrogen and sodium budget were calculated for two years (2015, 2016) from these data. According to our results the wastewater had high nitrogen content what was able to increase the nitrogen amount in the examined budget however without supplementary fertilizer it could not able to balance the budget only just at W60 treatment (irrigation with wastewater: N concentration 22,7 mg/l). The wastewater had high environmental risk as soil sodicity according to results of the sodium budget.

Keywords: nitrate leaching, salinization, wastewater, rainwater quality

Received 19 July 2018, Revised 12 December 2018, Accepted 28 December 2018

Introduction

Sustainable soil and water management and use of alternative water resources for agricultural production are one of the key elements of the fight against the consequences of the continuous increase in global population and effects of climate change (Singh 2015). The new water resources play determining role because of the water scarcity in addition to water and energy saving irrigation methods (Francés et al. 2017). The reuse of treated wastewater for irrigation is a suitable practice to close the water cycle in the agro-industrial sector (Vergine et al. 2017).

The climate change affects our country due to the basin character, the unfavourable distribution of rainfall is the consequence of the change of the regional climate patterns. Other example is the decreasing surface water quantity at summer time thereby there is a growing number of water bodies with water scarcity and regions with water stress. This water scarcity is occurred at summer time when the irrigation water demand is the highest (Hungarian River Basin

Management Plan 2015). Alternative water resources utilization should take priority over the conventional irrigation water resources (surface and subsurface waters) in the future in Hungary as well, similarly to the global trends because of the global warming and water scarcity.

However, inappropriate management of irrigation with wastewater can cause substantial risks to public health and the surrounding environment (Elgallal et al. 2017). Understanding the processes of nitrate leaching and introduction of deeper-rooted trees or other crops can minimise the ground water contamination (Khajanchi Lal et al. 2015). On the other hand, wastewater usually has a higher concentration of total dissolved solids and major ions and a higher electrical conductivity than fresh water especially in regions with hot climates or it can originate from many sources such as detergents and washing material, the chemicals used during the treatment process and other sources (Elgallal et al. 2017). Removing salts from wastewater for irrigation purposes is prohibitively expensive.

Therefore, there is a need for specific measures and management strategies to prevent and control the effects of salinity and sodicity during irrigation with wastewater (Elgallal et al. 2017).

The energy plantation may provide prosperous opportunity for the wastewater reuse in the future through irrigation (Vermees 2017). Energy willow and poplar have many favourable advantages compared to other plants: long growing season, high evapotranspiration rate, ability to take up nutrients and toxic elements with minimum leaching potential (Isebrands and Richards 2014). According to Ligetvári (2014) the use of sewage can make safer production of energy crops or forests or even not directly consumed agricultural products.

Arable crops can be characterised by low species and breed number of crops with minimum crop rotation (Grónás et al. 2006). To improve the biodiversity and to break the monotony of the landscape, installation of energy plantation would be one feasible prosperous solution. Diversification of arable crops would be beneficial to decrease the risk of the market and the cost of production (Grónás et al. 2006).

Aim of our lysimeter experiment was to determine the impact of a wastewater originated from an African intensive catfish farm on the nitrogen budget of the soil-water-plant system in order to evaluate the *nitrogen substitution effect* and *the risk of the nitrate leaching*. On the other hand, the aim of the study was to calculate also sodium budget to assess the *risk of the sodicity* regard to the high sodium concentration of the wastewater.

Materials and methods

Experiment site

The experiments were conducted at the National Agricultural Research and Innovation Centre (NAIK), Research Institute of Irrigation and Water Management (OVKI) in Szarvas, Hungary. The experiment was set up in the NAIK ÖVKI Lysimeter Station in 2014 in 64 pieces of 1 m³ vessels with energy willow. The year 2015 was the second year of the plantation when the irrigation started. The willow clone (no. 77, 82) were selected by the NAIK Forest Research Institute of Püspökladány Experimental Station.

Each treatment occurred in 8 replication/ lysimeter. The 1 m³ lysimeter vessels were filled with disturbed meadow topsoil. The soil was characterised by high clay content (~70-80%), low humus (~2%), lime content less than 0.5%, total soluble salt content less than 0.08%, and pH_{KCl} values between 5.88-6.97 (Kun et al. 2018).

Irrigation treatments

Three different irrigation waters were applied in the experiment. First one was originated from the Oxbow Lake of Körös River (K15, K30, K60 – numbers mean one-time irrigation doses in mm), while the second one was the wastewater (W15, W30, W60) from an intensive African catfish farm in Szarvas. The third irrigation water type (HG60) used for irrigation (only 60 mm doses) was treated wastewater, which was diluted (1:3) with Körös water than gypsum was added to it according to the following equation:

$$x = S z_e \times E$$

where the x means the gypsum quantity (mg/l or g/cm³), $S z_e$ = residual sodium carbonate index (RSC), E = equivalent weight of gypsum (86,1). There were one non-irrigated treatment (Control). The irrigation water quality was describe according to the 90/2008. (VII. 18.) Hungarian decree and FAO guideline as well (Kun 2018).

Sampling method and analyses

The irrigation water sampling occurred in the irrigation periods at every time between 15th April and 30th September according to Hungarian standard (MI-10-172/9-1990). Inorganic nitrogen (ammonium, nitrate, nitrite, total inorganic-N) and sodium was determined according to the Hungarian standards (MSZ EN ISO 11732:2005, MSZ EN ISO 13395:1999, MSZ EN ISO 13395:1999, MSZ 12750-20:1972, MSZ 1484-3:2006, respectively). In 2015 3 replication represents the Körös River and the wastewater quality, in 2016 the number of the replication was 7. In case of diluted and improved wastewater treatment (HG60) the irrigation quality was analysed based on 4 replications. The rainwater was collected at 4 rainfall events (14.06.2016, 20.09.2016, 19.04.2017, 29.06.2017) meaning 4 replications and the same parameters were

Table 1. The irrigation and the leachate water amounts (mm) in the experiment

| Treatments | Irrigation water amounts (mm) | | Leachate water amounts (mm) | |
|------------------------------------|-------------------------------|------|-----------------------------|------------------------|
| | 2015 | 2016 | 1 st period | 2 nd period |
| Körös 15 mm (K15) | 195 | 90 | 383 | 23 |
| Körös 30 mm (K30) | 390 | 180 | 409 | 106 |
| Körös 60 mm (K60) | 780 | 360 | 621 | 532 |
| Wastewater 15 mm (W15) | 195 | 90 | 463 | 358 |
| Wastewater 30 mm (W30) | 390 | 180 | 470 | 145 |
| Wastewater 60 mm (W60) | 780 | 360 | 717 | 313 |
| Diluted wastewater + gypsum (HG60) | 720 | 360 | 674 | 676 |
| Control (non-irrigated) | 0 | 0 | 655 | 130 |

Remark: 1st period: between 15.06.2015 and 17.06.2016 2nd period: between 17.06.2016 and 21.04.2017

analysed as from the irrigation waters. The lysimeter is suitable to collect the leachate. The leachate water amounts were measured 33 times between 03.07.2015 and 21.04.2017 for 22 months in 62 lysimeter vessels. (In H15 and HG60 treatments there was 1-1 not-working lysimeter.) The leachate sampling occurred in winter time in both years (16.01.2016, 28.02.2016, 06.02.2017, 08.02.2017) and the analysed parameters were nitrate and sodium according to the methods above. The results represent 2 replications per treatments per the analysed periods.

The willow stem sampling occurred in 2015 at time of the willow harvest (02.12.2015). Harvested stems from each treatment (in 8 replication) were collected and analysed for dry mass, total Kjeldahl-nitrogen and sodium concentration according to Hungarian standards (MSZ-08-1783-1:1983, MSZ EN ISO 5983-2:2009, MSZ-08-1783-5:1983).

Nitrogen and sodium budget calculation and statistical analyses

Based on the leachate analyses results nitrogen and sodium budget was calculated for two periods. First period (2015) was between 15.06.2015 and 17.06.2016 and the second period (2016) was between 17.06.2016 and 21.04.2017. For the calculation the input quantities were: the nitrogen fertilizers, the nitrogen content of the irrigation waters and the rain and the output quantities were the nitrogen amount that was accumulated in the willow stems and the nitrogen losses by leaching. The applied fertiliser amount was 40 kg N/ha in the first period and there were no applied fertiliser in 2016. Input inorganic-N of the irrigation water

were calculated by multiplying irrigation water amounts (Table 1). The input inorganic-N of the rainwater calculated by multiplying the precipitation volumes. The applied irrigation water amounts were different in 2015 and 2016 (Table 1), because the weather was different in the experiment years. In the first year the precipitation was 116.8 mm during the irrigation period but in the second year it was 220.6 mm. In 2015 the total precipitation was 400.6 mm and in 2016 it was 632.8 mm. Nitrate leaching loads were calculated by multiplying drainage volumes for each period (Table 1). The output nitrogen by the willow stem was calculated from the dry biomass data in 2015 and 2016 (not published data) and the Kjeldahl-nitrogen of the willow stems in 2015. The subtraction of the input and output N means the nitrogen budget (ΔN values). According to Arronson and Bergström (2001) the estimated ΔN values express the combined effect of mineralization, immobilization, build-up of the soil pool of nitrogen, and gaseous losses. Apart from the gaseous losses the others do increase or decrease the store of the soil nitrogen. So if ΔN has negative sign it means decreasing N-store in the soil and if ΔN is positive, the soil becomes richer in nitrogen. In our experiment the nitrogen content of the willow leaves were did not calculated in the nitrogen budget, due to the difficulty of estimate the leaf mass and partly because it can not be considered as real loss as leaves stay at the soil surface for the degradation processes, hence in our case the ΔN values also expresses some nitrogen in falling leaves at autumn. However, according to Szalai (1968) the leaves contain small amount of nitrogen. The sodium budget calculation was according to the same method above (without the fertilizer).

Table 2. Nitrogen budget of the willow lysimeter experiment in 2015 and 2016

| 2015 | | | | | | | | |
|---|---------------------|--------------------|-------------------|--------------------|----------------------|---------------------|----------------------|----------------------|
| | W15 | W30 | W60 | HG60 | K15 | K30 | K60 | Control |
| Input | | | | | | | | |
| Fertilizer (kg/ha) | 40.0 | 40.0 | 40.0 | 40.0 | 40.0 | 40.0 | 40.0 | 40.0 |
| N content of the irrigation water (kg/ha) | 43.3 | 86.7 | 173.3 | 76.3 | 1.3 | 2.6 | 5.2 | 0.0 |
| N content of the rainwater (kg/ha) | 6.2 | 6.2 | 6.2 | 6.2 | 6.2 | 6.2 | 6.2 | 6.3 |
| Output | | | | | | | | |
| N content of the willow stem (kg/ha) | 90.4 | 122.8 | 171.2 | 101.8 | 58.0 | 80.1 | 95.6 | 35.7 |
| N leaching (kg/ha) | 6.1 | 5.6 | 15.9 | 0.6 | 1.5 | 0.6 | 0.3 | 56.9 |
| Δ N (kg/ha) | -7.0 ^{abc} | 4.5 ^{abc} | 32.5 ^c | 20.1 ^{bc} | -11.9 ^{abc} | -31.8 ^{ab} | -44.5 ^a | -46.3 ^a |
| 2016 | | | | | | | | |
| | W15 | W30 | W60 | HG60 | K15 | K30 | K60 | Control |
| Input | | | | | | | | |
| Fertilizer (kg/ha) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| N content of the irrigation water (kg/ha) | 20.8 | 41.6 | 83.1 | 38.2 | 0.6 | 1.2 | 2.3 | 0.0 |
| N content of the rainwater (kg/ha) | 5.9 | 5.9 | 5.9 | 5.9 | 5.9 | 5.9 | 5.9 | 5.9 |
| Output | | | | | | | | |
| N content of the willow stem (kg/ha) | 72,8 | 92,4 | 95,6 | 53,2 | 46,6 | 46,6 | 41,6 | 28,5 |
| N leaching (kg/ha) | 0.4 | 5.5 | 12.3 | 1.1 | 0.9 | 0.4 | 0.3 | 8.1 |
| Δ N (kg/ha) | -46,1 ^a | -44,9 ^a | -6,5 ^c | -9,1 ^{bc} | -40,1 ^{ab} | -39,5 ^{ab} | -33,4 ^{abc} | -22,6 ^{abc} |

Remark: ANOVA was used to determine the significant difference between the treatments. The ^{a, b, c} indexes means the homogenous subsets of the Tukey's post-hoc test. The negative Δ N values mean the soil (and the willow leaves) contribution to the nitrogen budget.

The statistical calculation was performed in SPSS 22.0 Statistics Software. ANOVA and Tukey's Test was used to determine significant difference between Δ N values and the homogenous subset means no significant difference between the treatments. To determine significant different between Δ Na values Non-parametric, Kruskal-Wallis test was used in both years.

Results

Nitrogen budget

To calculate the N budget three input factor were determined. The N content of the fertilizer, irrigation water and rainwater build up the all input N (Table 2).

The wastewater contained considerably more inorganic-N (2015: 23.1 mg/l, 2016: 22.2 mg/l) than the diluted and improved irrigation water type (2015, 2016: 10.6 mg/l) and the irrigation water from the River Körös (2015: 0.64 mg/l, 2016: 0.67 mg/l). In the wastewater the 99.8% of inorganic-N was in ammonium form. Nitrogen (N) and phosphorous (P) in metabolic waste produced by fish are the origin of most dissolved N and P waste resulting from intensive aquaculture operations (Lazarri and Baldisserotto 2008, Tóth et al. 2016). The ammonia production

by fish is primarily dependent on the protein intake and metabolic efficiency of the fish, which is species-specific and is affected by waterborne ammonia levels (Dosdat et al. 2003). In case of the River Körös, the ammonium : nitrate : nitrite rate was 43:50:7. All of them was under the limit values of Decree No. 10 of 2010 (VIII. 18.) VM of the Ministry of Rural Development hence the oxbow lake can be consider as good quality surface water body. The input N content from the irrigation water was depended on the irrigation water amount (Table 1, Table 2).

The inorganic-N concentration of the rainwater was 1.25 mg/l. According to Gelencsér et al. (2012) the average total N concentration of the rainwater is 1.68 mg/l and the main nitrogen forms are inorganic forms. According to Csapák (2009) the ammonium concentration in the rainwater was 0.24 mg/l in $\text{NH}_4\text{-N}$, the nitrate was 0.21 mg/l in $\text{NO}_3\text{-N}$ and there were no nitrite. The input N in the budget from the precipitation was 6.2 kg/ha in 2015 and 5.9 kg/ha in 2016 (Table 2).

Because of the different weather conditions of the experimental years the applied irrigation water amount was twice more in 2015 than in 2016. In 2016 fertilizer were not applied and less irrigation water was used hence the N input in

the second year were lower than in 2015. In the treatments without waste water (K15, K30, K60 and Control) the fertilise-input was only 13-16%, but in waste and diluted water treatments the fertilise-input even was not more than 36-41% of the total input N in 2015.

Two output factor was used to determine the nitrogen budget: the N content of the willow stem and the N content of the leaching water. The mean N content of the willow stem was 0.51 m/m%, the range was 0.2-0.87 m/m%. According to our results the lowest accumulated nitrogen values of the willow stem was 29 kg N/ha and the highest was 171 kg N/ha (Table 2). According to Dimitriou and Aronsson (2004) the N-uptake of the willow was 110-115 kg N/ha in clay soil and 36-44 kg N/ha in sandy soil. Four willow clones were compared by Curneen and Gill (2014), according to their results the smallest N-uptake was 82 kg N/ha (in case of *Torhild* clone, irrigated with freshwater) and the highest was 262 kg N/ha (in case of *Sven* clone, irrigated with secondary treated wastewater). According to Galbally et al. (2013) the willow N-uptake after treatment with biosolid was 107 kg N/ha and after irrigation with distillery effluent water it was 231 kg N/ha. So our data can show the same order of magnitude then other researchers'.

According to our results the mean nitrogen losses caused by nitrate leaching was 10.9 kg N/ha in 2015 and it was 3.6 kg N/ha in 2016 (Table 2). According to willow lysimeter experiment

belongs to Aronsson and Bergström (2001) the highest N leaching was measured in the first year of the willow plantations (clay and sandy soil: 341-140 kg N/ha). After the starting year in the second (43-17 kg N/ha, respectively) and in the third year (3-1 kg N/ha, respectively) the N leaching was negligible. According to Mortensen et al. (1998) also in the first year of the plantation was the highest nitrogen leaching (130-142 kg N/ha) and it was decreased in the second (9-61 kg N/ha) and in the third (0-4 kg N/ha) years.

The subtraction of the input and the output N amount was the result of the nitrogen budget (ΔN). In 2015 in the wastewater treatments were calculated the most favourable ΔN values, in three cases (W30, W60, HG60) it was positive. The lowest values were in case of irrigation with River Körös water and in the Control treatment (mean of the ΔN values for this 4 treatment was -34 kg N/ha). In 2016 the most favourable ΔN values were also in the W60 and the HG60 treatments. All the other treatments had lower values then W60 and HG60 (Table 2).

Sodium budget

To calculate the Na budget two input factor was determined. The Na content of the irrigation water and rainwater build up the all input Na. The wastewater had considerably higher Na concentration (2015: 278 mg/l, 2016: 274 mg/l) than the diluted and improved irrigation water

Table 3. Sodium budget of the willow lysimeter experiment in 2015 and 2016

| 2015 | | | | | | | | |
|--|-------|--------|--------|-------|------|-------|-------|---------|
| | W15 | W30 | W60 | HG60 | K15 | K30 | K60 | Control |
| Input | | | | | | | | |
| Na content of the irrigation water (kg/ha) | 542.6 | 1085.2 | 2170.4 | 945.0 | 81.5 | 162.9 | 325.9 | 0.0 |
| Na content of the rainwater (kg/ha) | 14.1 | 14.1 | 14.1 | 14.1 | 14.1 | 14.1 | 14.1 | 14.2 |
| Output | | | | | | | | |
| Na content of the willow stem (kg/ha) | 1.0 | 1.5 | 1.4 | 1.2 | 0.5 | 0.7 | 0.9 | 0.3 |
| Na leaching (kg/ha) | 28.4 | 29.6 | 65.7 | 88.4 | 23.3 | 33.6 | 48.7 | 37.8 |
| Δ Na (kg/ha) | 527.3 | 1068.2 | 2117.4 | 869.5 | 71.8 | 142.7 | 290.4 | -23.9 |
| 2016 | | | | | | | | |
| | W15 | W30 | W60 | HG60 | K15 | K30 | K60 | Control |
| Input | | | | | | | | |
| Na content of the irrigation water (kg/ha) | 247.0 | 494.0 | 987.9 | 472.5 | 26.0 | 52.0 | 104.0 | 0.0 |
| Na content of the rainwater (kg/ha) | 13.4 | 13.4 | 13.4 | 13.4 | 13.4 | 13.4 | 13.4 | 13.4 |
| Output | | | | | | | | |
| Na content of the willow stem (kg/ha) | 0,8 | 1,1 | 1,0 | 0,6 | 0,4 | 0,5 | 0,4 | 0,3 |
| Na leaching (kg/ha) | 1.6 | 11.1 | 68.1 | 61.5 | 5.7 | 14.3 | 29.9 | 6.0 |
| Δ Na (kg/ha) | 258,0 | 495,2 | 932,3 | 423,8 | 33,4 | 50,7 | 87,1 | 7,2 |

type (2015, 2016: 131 mg/l) and the irrigation water from the River Körös (2015: 42 mg/l, 2016: 29 mg/l). The high sodium content of the wastewater was originated from the fishfarm water source because thermal water is used to provide the high water amount for intensive technology (Tóth et al. 2016). Because of the wastewater quality the input sodium from the irrigation water was very high (200-2200 kg/ha/year), (Table 3).

Two output factor was used to determine the Na budget: the Na content of the willow stem and the Na content of the leaching water. The mean Na content of the willow stem was 53.6 mg/kg, the range was 23.8-161 mg/kg. It means 0.3-1.5 kg/ha/year which is negligible compared to the leaching. According to Jama-Rodzská et al. (2016) most of sodium (~300 mg/kg) was found in the willow shoots during the first year of the study which decreased in the second year (~200 mg/kg). According to Stolarski et al. (2017) the average Na concentrations of the analysed willow clones were between 80-180 mg/kg. Due to the low ion uptake of the willow the output mean Na values were very low in both years varied between 0.3 and 1.5 kg Na/ha (Table 3).

According to our results the mean Na losses caused by leaching was 44.0 kg Na/ha in 2015 and it was 24.5 kg Na/ha in 2016 (Table 3). According to Stefanovits (1963) the average sodium losses by leaching on clay soil is 9-42 kg/ha/year, in our lysimeter experiment the sodium losses varied between 1 and 125 kg/ha/year. According to Sorrenti and Toselli (2016) in 18-month-lysimeter experiment with 1-year old nectarine the sodium losses by leaching was 224 kg/ha in sandy soil.

The subtraction of the input and output Na amount was the result of the sodium budget (ΔN). In 2015 in the wastewater treatments were calculated the most unfavourable ΔNa values between 527-2117 kg/ha. The prosperous sodium budget was calculated in both years in the treatments with irrigated with River Körös. The lowest ΔNa value were in case of non-irrigated treatment, in the Control treatment in both years (Table 3).

Discussion

The ΔN value of the W60 treatment in 2015 was significantly higher than the other treatments (Table 2). In this case there were the highest N input owing to the nitrogen content of the wastewater. According to our results the wastewater not only does not pose the threat to groundwater nitrate pollution but it could be useful in the irrigation management and foster the sustainable soil management practices also. However according to the European regulation the maximum permissible amount of fertilizer per hectare is 170 kg N. Therefore, the concentration of nitrogen in the wastewater should be taken into account during the application on nitrate sensitive areas (Council Directive 91/676/EEC). There were no significant differences between the ΔN values of the W15, W30, K15 treatments. In these cases the input nitrogen was approximately enough for the plants hence neither the soil nitrogen loading nor the soil exhaustion did not occur. The ΔN value of the K30, K60 and Control was the lowest. In the treatments with Körös River water (K30, K60) the input N was not enough for the soil-water-plant system because the irrigation water did not contain enough inorganic-N hence the ΔN values were negative and it means the soil exhaustion. According to Kenessey (1931) the irrigation should always coexist with nutrient supply, because the water explore the soil nutrient stock and the plants become rich in nutrients but the soil exhaustion can occur. Otherwise in case of the Control treatment the adverse nitrogen budget was caused by high nitrogen losses by leaching. According to Szalókiné and Szalóki (2003) in the soil of non-irrigated treatment, there is more nitrogen (because of the less plant N-uptake) which, during the winter period, results in higher nitrate concentration in leachate waters.

In 2016 (the third year of the plantation) the lowest value was in the treatment W15 and W30, but they were in the same homogenous subset as the K15, K30, K60 and Control (Table 2). Nitrogen balance could be realized only in W60 and HG60 treatment, where the input N and the output N was approximately equal. In 2016 the

Table 4. Subset of the Δ Na values in the different treatments

| | | 2015 | | | | | | |
|-----------------|------------|------------|------------|-------------|------------|------------|------------|-----------------|
| | <i>H15</i> | <i>H30</i> | <i>H60</i> | <i>HG60</i> | <i>K15</i> | <i>K30</i> | <i>K60</i> | <i>Kontroll</i> |
| <i>H15</i> | - | -541 | -1590 | -342 | 456 | 385 | 237 | 551* |
| <i>H30</i> | | - | -1049 | 199 | 996** | 925* | 778 | 1092*** |
| <i>H60</i> | | | - | 1248 | 2046*** | 1975** | 1827* | 2141*** |
| <i>HG60</i> | | | | - | 798* | 727 | 579 | 893** |
| <i>K15</i> | | | | | - | -71 | -219 | 96 |
| <i>K30</i> | | | | | | - | -148 | 167 |
| <i>K60</i> | | | | | | | - | 314 |
| <i>Kontroll</i> | | | | | | | | - |

| | | 2016 | | | | | | |
|-----------------|------------|------------|------------|-------------|------------|------------|------------|-----------------|
| | <i>H15</i> | <i>H30</i> | <i>H60</i> | <i>HG60</i> | <i>K15</i> | <i>K30</i> | <i>K60</i> | <i>Kontroll</i> |
| <i>H15</i> | - | -236 | -671 | -160 | 225 | 208 | 171 | 251 |
| <i>H30</i> | | - | -435 | 76 | 461** | 444 | 407 | 487*** |
| <i>H60</i> | | | - | 511 | 896*** | 878** | 842 | 922*** |
| <i>HG60</i> | | | | - | 386 | 368 | 331 | 411** |
| <i>K15</i> | | | | | - | -18 | -55 | 26 |
| <i>K30</i> | | | | | | - | -37 | 43 |
| <i>K60</i> | | | | | | | - | 80 |
| <i>Kontroll</i> | | | | | | | | - |

Remark: Results of the Non-parametric, Kurskal-Wallis test. Values (i-j) mean the subset of the columns (i) and the rows (j). (*: $p < 0.05$ **: $p < 0.01$ ***: $p < 0.001$).

higher Δ N values in the control occurred than in the previous year because of the less nitrogen loss by leaching. The lower Δ N values in the W15 and W30 was the results of the lack of nitrogen fertilizer, because the output by the willow stems and the leaching was less than in 2015. According to Arronsson and Bergström (2001) in the third year of the willow plantation the Δ N value was -39 kg N/ha (low N input and irrigation water) and it was -237 kg N/ha (high N input and irrigation water) on clay soil.

For all irrigated treatments the sodium budget was unfavourable but in case of the River Körös the values were significantly lower (Table 3, Table 4). The Na concentration of the River Körös is very low in both years and it makes it suitable for irrigation purposes. According to Ayers and Westcot (1989) above the limit of sodium (3 meq/l) in the irrigation water ion toxicity can occur and the sodium sensitive plant developing can decline. According to the recommendation of the FAO for irrigation water quality under the limit there is no harmful impact of the sodium content in case of sprinkler irrigation (Ayers and Westcot 1989). According to our results the wastewater and the diluted and improved water did not meet the recommended limit (~12 meq/l and 5.7 meq/l, respectively) but the River Körös (1.8 meq/l) is under the limit.

In all irrigated treatments with wastewater the Δ Na values were the highest which means that harmful sodium stay at the system (Table 3).

In both years neither the sodium contain of the stems nor the sodium leaching were not able to increase the output Na amount making the sodium budget more favourable. In the treatments irrigated with wastewater the sodium outputs were higher than in the treatments with River Körös however it was not enough high to reduce the sodium loading during the irrigation period owing to the wastewater quality. According to Nouri et al. (2017) green remediation is the use of vegetation to remove or contain environmental contaminants such as heavy metals, trace elements, organic compounds and radioactive compounds in soil or water and it is suggested to use for treat the soil salinity. In our case the willow did not uptake and accumulate so many sodium ion to balance the sodium budget.

In case of the HG60 treatment the wastewater was diluted and improved by gypsum hence the sodium budget were better than in the W60 (Table 3). However in the HG60 the sodium budget was not auspicious and also truly high hence the soil monitoring is exceptionally important for the sustainable, long-term irrigation. In both year, the Δ N value were significantly higher than in the K60 because the N input was higher

(from the diluted wastewater origin), (Table 2). According to our results the wastewater treatment was sufficient to reduce the sodium loading of the soil-water-plant system compared to the wastewater irrigation in both years and it was able to increase the nitrogen budget.

Summary

Aim of our study was to evaluate the impact of the wastewater irrigation on the nitrogen budget of soil-water-plant system and to assess the environmental risk of the sodium accumulation. The wastewater is originated an intensive fish farm hence it had high nitrogen content what it was able to increase the nitrogen amount

in the examined system however without supplementary fertilizer it could not able to balance the budget only just at W60 treatment. The wastewater had high environmental risk as soil sodicity according to results of the sodium budget. Even the treated wastewater require soil monitoring to ignore the sodium accumulation, however the dilution and added gypsum was able to reduce the rate of the sodium accumulation of the wastewater.

Acknowledgements

This research was supported by the Hungarian Ministry of Agriculture.

References

- Aronsson P.G., Bergström L.F. (2001): Nitrate leaching from lysimeter-grown short-rotation willow coppice in relation to N-application, irrigation and soil type. *Biomass and Bioenergy* **21**, 155-164. [https://doi.org/10.1016/S0961-9534\(01\)00022-8](https://doi.org/10.1016/S0961-9534(01)00022-8)
- Ayers R.S., Westcot D.W. (1989): *Water quality for agriculture*. FAO.Rome.
- Curneen S.J., Gill L.W. (2014): A comparison of the suitability of different willow varieties to treat on-site wastewater effluent in an Irish climate. *Journal of Environmental Management* **133**, 153-161. <https://doi.org/10.1016/j.jenvman.2013.12.004>
- Csapák A. (2009): *Települési vízgazdálkodás, lakossági csapadékvíz-gyűjtés és felhasználás*. Doktori (PhD) értekezés. ELTE TTK. Budapest.
- Dimitriou I., Aronsson P. (2004): Nitrogen leaching from short-rotation willow coppice after intensive irrigation with wastewater. *Biomass and Bioenergy* **26**: 5, 433-441. <https://doi.org/10.1016/j.biombioe.2003.08.009>
- Dosdat A., Person-Le Ruyet J., Covés D., Dutto G., Gasset E., Roux A., Lemarié G. (2003): Effect of chronic exposure to ammonia on growth, food utilisation and metabolism of the European sea bass (*Dicentrarchus labrax*). *Aquatic Living Resources* **16**: 6, 509-520. <https://doi.org/10.1016/j.aquativ.2003.08.001>
- Elgallal M., Fletcher L., Ewans B. (2016): Assessment of potential risks associated with chemicals in wastewater used for irrigation in arid and semiarid zones: A review. *Agricultural Water Management* **177**, 419-431. DOI: <http://dx.doi.org/10.1016/j.agwat.2016.08.027> <https://doi.org/10.1016/j.agwat.2016.08.027>
- Francés G.E., Quevauviller P., González E.S.M., Amelin E.V. (2017): Climate change policy and water resources in the EU and Spain. A closer look into the Water Framework Directive. *Environmental Science and Policy* **69**, 1-12. <https://doi.org/10.1016/j.envsci.2016.12.006>
- Galbally P., Ryan D., Fagan C.C., Finnan J., Grant J., McDonnell K. (2013): Biosolid and distillery effluent amendments to Irish short rotation coppiced willow plantations: impacts on groundwater quality and soil. *Agricultural Water Management* **116**, 193-203. <https://doi.org/10.1016/j.agwat.2012.07.010>
- Grónás, V., Magyari, J., Belényesi, M., Centeri, Cs. (2006): Agrár-környezetgazdálkodási programok bevezetésének hatása a kijelölt mintaterületek földhasználatára és természeti értékeinek védelmére. *Tájökológiai Lapok* **4**: 2, 277-289.
- Isebrands J.G., Richardson J. (editors) (2014): *Poplars and Willows. Trees for Society and the Environment*. FAO and CABI. Rome.
- Jama-Rodzenska A., Bocianowski J., Nowak W., Ciszek D. (2016) The influence of communal sewage on the content of macroelements in the stem of selected clones of willow (*Salix viminalis* L.). *Ecological Engineering* **87**, 212-217. <https://doi.org/10.1016/j.ecoleng.2015.11.046>
- Kenessey B. (1931): *Vízgazdálkodás. A magyar földművelésügyi miniszter kiadványai* **2**, 29-42.

- Khajanchi Lal, Minhas P.S., Yadav R.K. (2015): Long-term impact of wastewater irrigation and nutrients II. Nutrients balance, nitrate leaching and soil properties under peri-urban cropping system. *Agriculture Water Management* **156**, 110-117. <https://doi.org/10.1016/j.agwat.2015.04.001>
- Kun Á. (2018): Intenzív halnevelő telepről származó elfolyóvíz öntözésre való alkalmasságának vizsgálata. *Hidrológiai Közlöny* **98**: 1, 60-71.
- Kun Á., Bozán Cs, Oncsik B.M., Barta K. (2018): Evaluating of wastewater irrigation in Lysimeter experiment through energy willow yields and soil sodicity. *Carpathian Journal of Earth and Environmental Sciences* **13**: 1, 77-84.
- Lazzari R., Baldisserotto B. (2008): Nitrogen and phosphorus waste in fish farming. *Boletim do Instituto de Pesca* **34**: 4, 591-600.
- Ligetvári F., Juhász E., Bardóczyné Székely E. (2014): Velünk élő történelem: Szennyvíz- és szennyvíziszap-hasznosítás. *Hidrológiai Közlöny* **94**: 4, 2-9.
- Mortensen J., Nielsen K.H., Jorgensen U. (1998) Nitrate leaching during establishment of willow (*Salix Viminalis*) on two soil types and at two fertilization levels. *Biomass and Bioenergy* **15**: 6, 457-466. [https://doi.org/10.1016/S0961-9534\(98\)00056-7](https://doi.org/10.1016/S0961-9534(98)00056-7)
- Nouri H., Borujeni S.C., Nirola R., Hassanli A., Beecham S., Alaghmand S., saint C., Mulcahy D. (2017): Application of green remediation on soil salinity treatment: A review on halophytoremediation. *Process Safety and Environmental Protection* **107**, 94-107. <https://doi.org/10.1016/j.psep.2017.01.021>
- Singh A. (2015): Poor quality water utilization for agricultural production: An environmental perspective. *Land Use Policy* **43**, 259-262. <https://doi.org/10.1016/j.landusepol.2014.11.015>
- Sorrenti G., Toselli M. (2016): Soil leaching as affected by the amendment with biochar and compost. *Agriculture, Ecosystems and Environment* **226**, 56-64. <https://doi.org/10.1016/j.agee.2016.04.024>
- Stefanovits P. (1963): Magyarország talajai Mezőgazdasági Kiadó. Budapest.
- Stolarski M.J., Niksa D., Kryzaniak M. (2017): Elemental composition of willow short rotation crops biomass depending on variety and harvest cycle. *Biomass and Bioenergy* **105**, 342-350. <https://doi.org/10.1016/j.biombioe.2017.07.016>
- Szalai I. (1968): Növényélettan. Tankönyvkiadó. Budapest.
- Szalókiné Z.I., Szalóki S. (2003): Nitrátlemosódás vizsgálata liziméteres és szabadföldi tartamkísérletben. *Agrokémia és Talajtan*. **52**: 1-2, 35-52.
- Tóth F., Kerepeczki É., Berzi N.L., Gál D. (2016): Létesített vizes élőhelyek hasznosítása az intenzív haltermelés elfolyóvizének kezelésében. Kutatói utánpótlást elősegítő program I. szakmai konferenciája. Gödöllő, NAIK kiadása
- Vergine P., Salerno C., Libutti A., Beneduce L., Gatta G., Berardi G., Pollice A. (2017): Closing the water cycle in the agro-industrial sector by reusing treated wastewater for irrigation. *Journal of Cleaner Production* **164**, 587-596. <https://doi.org/10.1016/j.jclepro.2017.06.239>
- Vermes L. (2017): Vízgazdálkodásunk mostoha gyermeke – a szennyvízöntözés. *Hidrológiai Közlöny* **97**, 66-75.
- Council Directive 91/676/EEC of 12 December 1991 concerning the protection of waters against pollution caused by nitrates from agricultural sources. <https://eur-lex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:31991L0676&from=EN>
- Decree No. 10 of 2010 (VIII. 18.) VM of the Ministry of Rural Development providing limit values for water pollutants in surface waters and rules of application
- Hungarian River Basin Management Plan 2015. Available online at 12.07.2018 <https://www.vizugy.hu/index.php?module=vizstrat&programelemid=149>

OBITUARY

György FÜLEKY (1945-2018) prominent agrochemist passed away

Kálmán RAJKAI

Hungarian Academy of Sciences, Institute for Soil Science and Agricultural Chemistry,
1022 Budapest, Herman Ottó út 15.; E-mail: rajkai.kalman@agrar.mta.hu

Professor Emeritus György Füleky former Head of the Department of Soil Science and Agrochemistry, Director of the Institute of Environmental Sciences, Dean of the Agricultural and Environmental Sciences Faculty, and Deputy Rector of the Szent István University died on May 4 2018 at the age of 74.



György FÜLEKY (1945-2018)

György Füleky was born in Ekecs on February 5, 1945. He continued his primary school, high school and university studies in Budapest. He graduated from the Faculty of Chemistry and Physics at the Eötvös Lóránd University in 1968. Between 1968 and 1983, he worked as a research assistant at the Research Institute of Soil Science and Agricultural Chemistry at the Hungarian Academy of Sciences, and after becoming a candidate for agricultural sciences in 1978 as a senior lecturer. The title of his candidate thesis was “The phosphorous state and easily soluble phosphorus content of soils”. Since 1983, he worked as assistant professor at the Agricultural Chemistry Department at the Agricultural University of Gödöllő. His university positions were chronologically the followings: 1987-1990 Deputy Assistant Professor of Agricultural Sciences; from 1991 Professor at the Department of Soil Science and Agrochemistry; from 1990 to 2008 Head of the Department of Soil Science and Agrochemistry; 1991-1994 Deputy Dean of Education of the Faculty. He awarded the Széchenyi Professorate Fellowship between 1998 and 2001. He was the deputy rector of the Szent István University from 2000 to 2003. He functioned as Director of the Institute of Environmental Sciences between 2008 and 2010, and as Dean of the Faculty of Agricultural and Environmental Sciences in 2012.

György Füleky had a wide range of education subjects. Most importantly he taught subjects in graduate courses in agrochemistry, soil management, environmental effects of agriculture, geochemical circuits, land evaluation, environmental reconstruction, and environmental practices. In English language he held courses in agrochemistry, plant nutrition, and land evaluation. In the doctoral training program he was responsible for soil fertility and soil chemistry specializations.

György Füleky designed the thematic of the subjects listed above. He prepared the curriculum for agricultural engineer degree after the political change of regime in 1991 on the Agricultural University of Gödöllő. Nationally, in 1989, he prepared a new curriculum for the Soil Science – Soil Fertility Engineer vocational training what he led till 1998. In 1998 he worked on the curriculum of Soil Science for the Faculty of Civil Engineering. After preparing accreditation of the Agricultural Environmental Management vocational training program he led it till 2006.

In 1993 he developed the doctoral training program for Soil and Agrochemical Fundamentals for Environmental Management and led it till 2000. From 2000 he led the Soil Science, Agrochemistry, Environmental Chemistry subareas of the Environmental Sciences Doctoral School to 2014. Under his leadership 16 aspirants earned the PhD degree.

Additionally György Füleky worked out a curriculum for five year university and three-year college education of the Environmental Engineer vocational training. He developed the curriculum for specialization in Disaster Management of that training. Between 2002 and 2005 he participated in the work of the Agri-Bologna Committee and in the design of the BSc and MSc programs. At Szent István University, he developed curriculums for Environmental Engineer BSc and MSc programs. By the end of 2014, MSc's Disaster Management Engineer's curriculum was also prepared by him.

In 1986, he launched the journal “Bulletin of the University of Agricultural Sciences Gödöllő”, which was published from 2000 as “Bulletin of the Szent István University”. He was chairman of the editorial board of the periodical until its termination in 2012.

Prof. Füleky organized and edited the proceedings of 10 conferences entitled as “Landscape changes in the Carpathian Basin”.

He was the author and editor of “A talaj” (Soil) (1988), “Talajtan” (Soil Science) (1999), “Tápanyag-gazdálkodás” (Nutrition Management) (1999), Korszerű tápanyag-gazdálkodás” (Modern Nutrition Management) (2014) and “Tápanyag-gazdálkodás mezőgazdasági mérnökök számára” (Nutrition Management for Agricultural Engineers) (2014) books and textbooks. He did important editorial work in the production of the monograph EOLSS (UNESCO) entitled as “Cultivated plants as primarily food sources” issued in 2000.

Prof. Füleky presented different social activities, as chairman of the Hungarian Quality Compost Society, whereby he was one of the main organizers of the “Biological Waste Management Conference” organized in 2014 Gödöllő. He was a member of the MTA Soil Science, Water Management and Crop Cultivation committee and the Editorial Board of Agrochemistry and Soil Science Journal.

His research topics covered both basic and applied research and development. György Füleky's most important basic research activity focused on determination of the nutrient supply capacity of soils by plants, by chemical methods and their mathematical description. His research included determination of toxic elements in the soil by the rapid seedling biological test and the hot water percolation (HWP) chemical method. He developed a rapid methods for determining nitrogen content of damaged plant parts and fertilizers. In the framework of his applied research, he primarily intended to monitor the unfavorable or favorable effects of intensive crop production and nutrient management technologies on soil. In the field of environmental management, determination of the natural environment of the past times he concerned with the parallel use of soil and archaeological methods. With this concept he initiated a new research branch in the Hungarian soil research.

He has received numerous professional and scientific awards and publishing prizes for his outstanding work.

György Füleky was demanding and knowledgeable. He communicated with literary variety and inimitable kindness, he was shaded and always personalized. People who watched his words and understood his thoughts learned much from him. He was liked by closer and distant colleagues and students because he was straight, warm-hearted and selfless. Through his knowledge, extensive cultivation and unshakeable honesty, he gained undeniable prestige. Because of his gentle nature, helpfulness and special humor, he also had a unique love by his colleagues.

György Füleky's professional work remains active long and even will be continued by his colleagues and students in Hungary and the Carpathian Basin. We are all grateful for his oeuvre, his educational activities and opening the soil science to co-sciences of archeology and agriculture in historical times in Hungary and neighboring countries in the Carpathian Basin. All of us will miss his dear, helpful personality, cheerful humor and professional wisdom. Colleagues working on soil and agrochemistry disciplines, as well as his former students express a thankful heart and ask for eternal rest.

Source of the graphics

Front cover:

Gallo-Roman harvesting machine, called Vallus. Source: U. Troitzsch - W. Weber (1987): Die Technik : Von den Anfängen bis zur Gegenwart

Rear cover:

Portrait of Columella, in Jean de Tournes, *Insignium aliquot virorum icones*. Lugduni: Apud Ioan. Tornaesium 1559. Centre d'Études Supérieures de la Renaissance - Tours



URBÁNYI Béla, editor-in-chief

DSc /agric/, interim dean of the Faculty of Agricultural and Environmental Sciences of the Szent István University, Gödöllő, Hungary, member of the Animal Husbandry and Foraging Committee of the Hungarian Academy of Sciences. Professional fields: genetics and biotechnology on fish, general aquaculture and water toxicology.



Lucius Junius Moderatus Columella

(AD 4 – 70) is the most important writer on agriculture of the Roman empire. His *De Re Rustica* in twelve volumes has been completely preserved and forms an important source on agriculture. This book was translated to many languages and used as a basic work in agricultural education until the end of the 19th Century.